

MED for GP Model Selection

Kristyn Pantoja

1/30/2020

MMED for GPs

Sequential M-MED for GP

Space-filling input points

Close-together input points

Increasingly farther away input points

Finer Grid

MMED for GPs

Applying M-MED to Gaussian Process Model Selection

- ▶ Goal: Choose a design that will distinguish the two gaussian process models.
- ▶ Distinguishing functions vs. distributions over functions:
 - ▶ For regression models, we use $f_D(\mathbf{x}) = \text{Wasserstein}(\phi_{0,\mathbf{x}}, \phi_{1,\mathbf{x}})$. What is the distance function now? What are $\phi_{0,\mathbf{x}}, \phi_{0,\mathbf{x}}$?
 - ▶ Key Question: Do we need to consider the predictive distribution for each GP model?
 - ▶ Doing so would give us an option for $\phi_{0,\mathbf{x}}, \phi_{0,\mathbf{x}}$.
 - ▶ We would need to have at least some data in order to model each Gaussian Process (training set) and use M-MED to select points for comparing them.

Simulations Set-Up

- ▶ I consider two cases:
 - ▶ Gaussian vs. Matern kernels, where the true function is generated from the Matern kernel
 - ▶ Matern vs. Periodic kernels, where the true function is generated from the Periodic kernel
- ▶ To evaluate MED for each case, I draw uniformly selected input points for my training set, and then apply MED to the data.
- ▶ I consider two measures for comparing MED to a space-filling design:
 - ▶ ratio of RSS for each hypothesized kernel:


$$\frac{\sum_{i \in \mathbf{D}} (y_i^{\text{pred}_0} - y_i^{\text{new}})^2}{\sum_{i \in \mathbf{D}} (y_i^{\text{pred}_1} - y_i^{\text{new}})^2}$$

- ▶ likelihood ratio:


$$\frac{L(y^{\text{new}} | \xi, y^{\text{obs}}, \mathbf{X}^{\text{obs}}, \Theta = 0)}{L(y^{\text{new}} | \xi, y^{\text{obs}}, \mathbf{X}^{\text{obs}}, \Theta = 1)}$$

Gaussian vs. Matern (simulation)


Space-filling




M-MED



M-MED



Wasserstein(x)



Gaussian vs. Matern: $\log(\text{RSS0}/\text{RSS1})$

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-0.62	0.053	0.25	0.54	0.62	5.5

Space-filling

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-0.42	0.042	0.14	0.42	0.39	4.8

Percentage of simulations that resulted in M-MED evaluations that were larger than Space-filling evaluations

[1] 0.68

Gaussian vs. Matern: log ratio of predictive densities

(after removing NAs caused from non-invertible matrix)

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-Inf	-2.9e+08	-4.8e+07	-Inf	-9.9e+05	-1.3e+05

Space-filling


Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-2.7e+10	-1.2e+06	-3.7e+05	-1.1e+09	-9.6e+04	-1.8e+04

Percentage of simulations that resulted in M-MED evaluations that were smaller than Space-filling evaluations


[1] 0.875

Matern vs. Periodic (simulation)


Space-filling




M-MED



M-MED



Wasserstein(x)



Matern vs. Periodic: $\log(\text{RSS0}/\text{RSS1})$

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.56	1.3	1.8	2	2.2	3.6

Space-filling

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-0.094	0.72	1	1.1	1.4	3

Percentage of simulations that resulted in M-MED evaluations that were larger than Space-filling evaluations

```
## [1] 1
```

Matern vs. Periodic: log ratio of predictive densities

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-280	-160	-130	-140	-110	-65

Space-filling

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-Inf	-99	-85	-Inf	-60	-38

Percentage of simulations that resulted in M-MED evaluations that were smaller than Space-filling evaluations

[1] 0.92

Sequential M-MED for GP


Will a sequential design improve results?

For the sequential designs, I:


1. Start with 6 input data
2. Use SMMED to sequentially gather 15 new data points in 3 steps (5 new points at each step)
3. To compare SMMED to a space-filling design, I use the previous evaluations on the 15 new points (pretending that data was not gathered for them yet)

Gaussian vs. Matern (sequentially): Step 1


Space-filling




M-MED



M-MED




Wasserstein(x)




Step 2


Space-filling




M-MED



M-MED




Wasserstein(x)




Step 3


Space-filling




M-MED



M-MED




Wasserstein(x)




Compare to non-Sequential M-MED


M-MED




M-MED



non-Seq M-MED



Seq M-MED



Gaussian vs. Matern: $\log(\text{RSS0}/\text{RSS1})$

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-0.49	-0.082	0.27	0.48	0.64	2.8

Space-filling

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-0.53	-0.21	0.24	0.45	0.57	2.5

Percentage of simulations that resulted in M-MED evaluations that were larger than Space-filling evaluations

[1] 0.64

Gaussian vs. Matern: log ratio of predictive densities

(after removing NAs caused from non-invertible matrix)

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-8.7e+05	-1.2e+05	-5.5e+04	-1.3e+05	-3.1e+04	-5.8e+03

Space-filling


Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-9.6e+08	-1.0e+07	-2.8e+06	-6.6e+07	-3.6e+05	-2.9e+04

Percentage of simulations that resulted in M-MED evaluations that were smaller than Space-filling evaluations


[1] 0

Matern vs. Periodic (sequentially): Step 1


Space-filling




M-MED



M-MED




Wasserstein(x)




Step 2


Space-filling




M-MED



M-MED




Wasserstein(x)




Step 3


Space-filling




M-MED







M-MED



Wasserstein(x)



Compare to non-Sequential M-MED



Matern vs. Periodic: $\log(\text{RSS0}/\text{RSS1})$

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.26	1.2	1.5	1.6	2	4.7

Space-filling

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
0.35	0.9	1.3	1.3	1.7	2.5

Percentage of simulations that resulted in M-MED evaluations that were larger than Space-filling evaluations

[1] 0.56

Matern vs. Periodic: log ratio of predictive densities

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-490	-250	-160	-210	-130	-89

Space-filling

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-150	-96	-75	-77	-55	-32

Percentage of simulations that resulted in M-MED evaluations that were smaller than Space-filling evaluations

```
## [1] 1
```


Space-filling input points

Motivation for space-filling input points


Since the Gaussian process first wants to fill in areas where the uncertainty is highest generally, a space-filling design would evenly distribute those uncertainty regions so that perhaps the Gaussian Process could start to focus on different things.

Gaussian vs. Matern (sequentially): Step 1


Space-filling




M-MED



M-MED




Wasserstein(x)




Step 2


Space-filling




M-MED



M-MED




Wasserstein(x)




Step 3


Space-filling




M-MED



M-MED




Wasserstein(x)




Compare to non-Sequential M-MED


M-MED




M-MED



non-Seq M-MED



Seq M-MED



Gaussian vs. Matern: $\log(\text{RSS0}/\text{RSS1})$

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-0.36	-0.085	-0.0041	-0.00016	0.15	0.25

Space-filling

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-0.39	-0.11	-0.0053	-0.009	0.13	0.24

Percentage of simulations that resulted in M-MED evaluations that were larger than Space-filling evaluations

[1] 0.6

Gaussian vs. Matern: log ratio of predictive densities

(after removing NAs caused from non-invertible matrix)

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-4.0e+05	-1.2e+05	-2.0e+04	-7.7e+04	-7.9e+03	-1.5e+03

Space-filling

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-5.3e+05	-1.2e+05	-6.4e+04	-1.0e+05	-1.5e+04	-9.3e+02

Percentage of simulations that resulted in M-MED evaluations that were smaller than Space-filling evaluations

[1] 0.32


Close-together input points

Motivation for close-together input points


Seeing the space-filling input points seems to lend more evidence to the theory that for the Gaussian vs. Matern case, the main difference is the uncertainty quantification and hence a space-filling design is optimal and sought-after. Maybe the close-together input points will tell a similar story.

Gaussian vs. Matern (sequentially): Step 1


Space-filling




M-MED



M-MED




Wasserstein(x)




Step 2


Space-filling




M-MED



M-MED




Wasserstein(x)




Step 3


Space-filling




M-MED



M-MED




Wasserstein(x)




Compare to non-Sequential M-MED


M-MED




M-MED



non-Seq M-MED



Seq M-MED



Gaussian vs. Matern: $\log(\text{RSS0}/\text{RSS1})$

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-0.58	0.47	1.6	1.4	2.1	3

Space-filling

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-0.48	0.41	1.5	1.3	2	2.7

Percentage of simulations that resulted in M-MED evaluations that were larger than Space-filling evaluations

[1] 0.76

Gaussian vs. Matern: log ratio of predictive densities

(after removing NAs caused from non-invertible matrix)

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-2.9e+07	-2.0e+06	-9.7e+05	-3.5e+06	-1.1e+05	-1.4e+04

Space-filling

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-5.3e+05	-1.2e+05	-6.4e+04	-1.0e+05	-1.5e+04	-8.0e+02

Percentage of simulations that resulted in M-MED evaluations that were smaller than Space-filling evaluations

[1] 0.92


Increasingly farther away input points

Motivation for increasingly farther away input points


For completion

Gaussian vs. Matern (sequentially): Step 1


Space-filling




M-MED



M-MED




Wasserstein(x)




Step 2


Space-filling




M-MED



M-MED




Wasserstein(x)




Step 3


Space-filling




M-MED







M-MED



Wasserstein(x)



Compare to non-Sequential M-MED



Gaussian vs. Matern: $\log(\text{RSS0}/\text{RSS1})$

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-0.37	-0.024	0.071	0.065	0.17	0.42

Space-filling

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-0.39	-0.029	0.07	0.066	0.2	0.44

Percentage of simulations that resulted in M-MED evaluations that were larger than Space-filling evaluations

[1] 0.48

Gaussian vs. Matern: log ratio of predictive densities

(after removing NAs caused from non-invertible matrix)

M-MED

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-2.5e+06	-3.0e+05	-1.3e+05	-2.8e+05	-4.1e+04	-1.3e+03

Space-filling

Min.	1st Qu.	Median	Mean	3rd Qu.	Max.
-5.3e+05	-1.2e+05	-6.4e+04	-1.0e+05	-1.5e+04	-9.3e+02

Percentage of simulations that resulted in M-MED evaluations that were smaller than Space-filling evaluations

[1] 0.64

Finer Grid

If we choose a finer grid for x_{seq}

