


Medical Terminology 2

- Presented by:
- Soumajit Sinha
- Lecturer, Dept. Of Hospital Management
- B.Sc.(H) (Medical Physiology), MBA,
- P.G.P(HealthCare Mgt.),
DBA,MAIAHM,M.sc(Life Sciences)

Renal Procedures

Kidney dialysis

Kidney dialysis is the process of removing excess water, solutes, and toxins from the blood in people whose kidneys can no longer perform these functions naturally. This is referred to as renal replacement therapy. The first successful




NEPHRITIS

- A condition in which the tissues in the kidney become inflamed and have problems filtering waste from the blood. Nephritis may be caused by infection, inflammatory conditions (such as lupus), certain genetic conditions, and other diseases or conditions.

- Hydronephrosis is a condition of the urinary tract where one or both kidneys swell. This happens because pee (urine) doesn't fully empty from your body. Symptoms may include sudden or intense pain in your back or side, vomiting, painful urination, blood in your pee, or weakness and fever due to a urinary tract infection. There are several approaches to treating the condition, depending on the underlying cause.

BILATERAL HYDRONEPHROSIS





Benign prostatic hyperplasia (BPH) is a condition that occurs when the prostate gland enlarges, potentially slowing or blocking the urine stream. Other names for benign prostatic hyperplasia include benign prostatic hypertrophy, an enlarged prostate, and BPH. BPH occurs only in men; approximately 8 percent of men aged 31 to 40 have BPH. In men over age 80, more than 80 percent have BPH.

Gangrene is a serious condition where a loss of blood supply causes body tissue to die. It can affect any part of the body but typically starts in the toes, feet, fingers and hands.

Gangrene can occur as a result of an injury, infection or a long-term condition that affects blood circulation.

Symptoms of gangrene include:

red, purple or black skin in the affected area, which may be harder to see on black or brown skin

swelling of the skin in the affected area

either a loss of sensation or severe pain in the affected area

sores or blisters in the affected area that bleed or produce a foul-smelling pus

Gangrenous tissue

Healthy tissue



© MANDCLINIC

Diabetes

Diabetes is a condition that happens when your blood sugar (glucose) is too high. It develops when your pancreas doesn't make enough insulin or any at all, or when your body isn't responding to the effects of insulin properly. Diabetes affects people of all ages. Most forms of diabetes are chronic (lifelong), and all forms are manageable with medications and/or lifestyle changes.

Glucose (sugar) mainly comes from carbohydrates in your food and drinks. It's your body's go-to source of energy. Your blood carries glucose to all your body's cells to use for energy.

When glucose is in your bloodstream, it needs help — a “key” — to reach its final destination. This key is insulin (a hormone). If your pancreas isn't making enough insulin or your body isn't using it properly, glucose builds up in your bloodstream, causing high blood sugar (hyperglycemia).

Over time, having consistently high blood glucose can cause health problems, such as heart disease, nerve damage and eye issues.

The technical name for diabetes is diabetes mellitus. Another condition shares the term “diabetes” — diabetes insipidus — but they're distinct. They share the name “diabetes” because they both cause increased thirst and frequent urination. Diabetes insipidus is much rarer than diabetes mellitus.

There are several types of diabetes. The most common forms include:

Type 2 diabetes: With this type, your body doesn't make enough insulin and/or your body's cells don't respond normally to the insulin (insulin resistance). This is the most common type of diabetes. It mainly affects adults, but children can have it as well.

Prediabetes: This type is the stage before Type 2 diabetes. Your blood glucose levels are higher than normal but not high enough to be officially diagnosed with Type 2 diabetes.

Type 1 diabetes: This type is an autoimmune disease in which your immune system attacks and destroys insulin-producing cells in your pancreas for unknown reasons. Up to 10% of people who have diabetes have Type 1. It's usually diagnosed in children and young adults, but it can develop at any age.

Gestational diabetes: This type develops in some people during pregnancy. Gestational diabetes usually goes away after pregnancy. However, if you have gestational diabetes, you're at a higher risk of developing Type 2 diabetes later in life.

Symptoms of Diabetes



Increased thirst.



Slow-healing cuts and sores.



Fatigue.



Blurred vision.



Frequent urination.



Unexplained weight loss.

Reproductive Syndromes

Sterility

Not getting pregnant, despite having carefully timed, unprotected sex for one year.

The cause of infertility may be difficult to determine but may include inadequate levels of certain hormones in both men and women, and trouble with ovulation in women.

The main symptom is an inability to get pregnant. In many cases, there are no additional symptoms.


Many treatments significantly improve the chances of getting pregnant. They include hormone treatments, fertility drugs and surgery. In addition, assisted reproduction uses various medical techniques to fertilise an egg.

Polycystic ovary syndrome (PCOS) is a condition in which the ovaries produce an abnormal amount of androgens, male sex hormones that are usually present in women in small amounts. The name polycystic ovary syndrome describes the numerous small cysts (fluid-filled sacs) that form in the ovaries.

PCOD or PCOS is a condition that affects women's ovaries, the reproductive organs that produce progesterone and estrogen hormones that help in regulating the menstrual cycle and also produce a small amount of hormones inhibin, relaxin, and male hormones called androgens. Almost 10% of women in the world are suffering from PCOD. In compare to PCOD women with PCOS produce higher-than-normal amounts of male hormones. This hormone imbalance causes them to skip menstrual periods.



Uterine fibroids are common growths of the uterus. They often appear during the years you're usually able to get pregnant and give birth. Uterine fibroids are not cancer, and they almost never turn into cancer.



Syphilis is an infection caused by bacteria. Most often, it spreads through sexual contact. The disease starts as a sore that's often painless and typically appears on the genitals, rectum or mouth. Syphilis spreads from person to person through direct contact with these sores. It also can be passed to a baby during pregnancy and childbirth and sometimes through breastfeeding.

After the infection happens, syphilis bacteria can stay in the body for many years without causing symptoms. But the infection can become active again. Without treatment, syphilis can damage the heart, brain or other organs. It can become life-threatening.

Early syphilis can be cured, sometimes with a single shot of medicine called penicillin. That's why it's key to get a health care checkup as soon as you notice any symptoms of syphilis. All pregnant people should get tested for syphilis at their first prenatal checkup too.



What is prostatitis?

Prostatitis is a frequently painful condition that involves inflammation of the prostate and sometimes the areas around the prostate.

Scientists have identified four types of prostatitis:

1. chronic prostatitis or chronic pelvic pain syndrome
2. acute bacterial prostatitis
3. chronic bacterial prostatitis
4. asymptomatic inflammatory prostatitis

Men with asymptomatic inflammatory prostatitis do not have symptoms. A health care provider may diagnose asymptomatic inflammatory prostatitis when testing for other urinary tract or reproductive tract disorders. This type of prostatitis does not cause complications and does not need treatment.

Erectile dysfunction (impotence) is the inability to get and keep an erection firm enough for sex.


Having erection trouble from time to time isn't necessarily a cause for concern. If erectile dysfunction is an ongoing issue, however, it can cause stress, affect your self-confidence and contribute to relationship problems. Problems getting or keeping an erection can also be a sign of an underlying health condition that needs treatment and a risk factor for heart disease.

Epididymitis

Epididymitis (ep-ih-did-uh-MY-tis) is an inflammation of the coiled tube, called the epididymis, at the back of the testicle. The epididymis stores and carries sperm. Males of any age can get epididymitis.

Epididymitis is most often caused by a bacterial infection, including sexually transmitted infections (STIs), such as gonorrhea or chlamydia. Sometimes, a testicle also becomes inflamed — a condition called epididymo-orchitis.

Epididymitis is usually treated with antibiotics and measures to relieve discomfort.





Hysterectomy is surgery to remove the uterus. After a hysterectomy, you will not menstruate (have periods) or be able to get pregnant. Uterus removal is a common treatment for a variety of conditions that affect a woman's reproductive organs.

What are the different kinds of hysterectomy?


.


Total hysterectomy: Removing your uterus and cervix, but leaving your ovaries.

Supracervical hysterectomy: Removing just the upper part of your uterus while leaving your cervix.

Total hysterectomy with bilateral salpingo-oophorectomy: Removing your uterus, cervix, fallopian tubes (salpingectomy) and ovaries (oophorectomy). If you haven't experienced menopause, removing your ovaries will start menopausal symptoms.

Radical hysterectomy with bilateral salpingo-oophorectomy: The removal of your uterus, cervix, fallopian tubes, ovaries, the upper portion of your vagina and some surrounding tissue and lymph nodes. This type of hysterectomy is performed when cancer is involved.





A laparotomy is a surgical incision (cut) into the abdominal cavity. This operation is performed to examine the abdominal organs and aid diagnosis of any problems, including abdominal pain. In many cases, the problem – once identified – can be fixed during the laparotomy. In other cases, a second operation is required.

