



COSMETIC PRODUCT SAFETY REPORT

For:

FUNGOMAX

In accordance with Regulation (EC) 1223/2009 of the European Parliament and of the Council of 30 November 2009 on cosmetic products

Responsible Person: Republic of Serbia: Clicky doo, Zvezdarskih Jelki 3/2, Beograd

EU: Clicky EOOD, Traditsa 6, Sofia, Bulgaria

Date: 20.09.2025.

Document version: 1

Safety Assessor: Vesna Milanković

Part A. Cosmetic Product Safety Information

Trade name: FUNGOMAX

Volume: 50ml

Product category: Skin lotion

Manufacturer: BIO-TEO centar d.o.o., Mitropolita Mraovića 1a, 11000 Belgrade, Serbia

1) Qualitative and quantitative composition of the product

INCI	Concentration of raw material in the product (%)	Raw material composition (%)	CAS	EINECS/ELINCS	Function	Impurities and allergens
GLYCERIN	20	100	56-81-5	200-289-5	Humectans	N/A
LAVANDULA HYBRIDA OIL	0.15	100	8022-15-9 / 91722-69-9	- / 294-470-6	Active, fragrance	CAMPBOR 8.5%, LINALYL ACETATE 35%, BETA-CARYOPHYLLENE 2.5%, GERANYL ACETATE 1%, PINENE 1.5%, TERPINEOL 2%, EUGENOL, LINALOOL, LIMONENE 1,5%
ROSMARINUS OFFICINALIS LEAF OIL	0.15	100	84604-14-8 / 8000-25-7	283-291-9	Active, fragrance	LIMONENE 1.5-4%
EUGENIA CARYOPHYLLUS LEAF OIL	0.1	100	8015-97-2 / 8000-34-8 / 84961-50-2	- / - / 284-638-7	Active, fragrance	EUGENOL max 88%
ALLANTOIN	0.15	100	97-59-6	202-592-8	Active	N/A
THYMUS VULGARIS EXTRACT	79.45	10--15	84929-51-1	284-535-7	Active	Eugenol 8%, Linalol 8%,
JUGLANS REGIA LEAF EXTRACT		7--10	84012-43-1	281-688-1		

SALVIA OFFICINALIS LEAF EXTRACT		4--6	84082-79-1 / 8022-56-8	282-025-9 / -		
QUERCUS ROBUR BARK EXTRACT		2.5--4	71011-28-4	275-129-0		
ALCOHOL		20-40	64-17-5	200-578-6		
WATER		20-40	7732-18-5	231-791-2 (I)		

* EUGENOL, LINALOOL, LIMONENE must be labelled, CAMPHOR, LINALYL ACETATE, BETA-CARYOPHYLLENE, GERANYL ACETATE, PINENE, TERPINEOL – products containing them may be placed on the Union market until 31 July 2026 and made available on the Union market until 31 July 2028

2) Physical and chemical characteristics and stability of the cosmetic product

a) Physical and chemical characteristics of the raw materials

For all raw materials used for production of this cosmetic product technical documentation (SDS, TDS, CoA, etc.) provided by the supplier of cosmetic ingredients is available and is attached to this document. After evaluating the documents provided, it was concluded that all the ingredients used for production of cosmetic product FUNGOMAX met the requirements of quality and purity criteria, as well as that they contain the information needed for an adequate assessment of the safety of the cosmetic product.

INCI	Chemical name	Molekular formula	Molecular weight	Water solubility	Vapour pressure (Pa)	log Pow
GLYCERIN	Glycerol	C3H8O3	92.09	Completely miscible with water at 25 °C.	N/A	log POW = -1.75 at 25 °C
LAVANDULA HYBRIDA OIL	N/A	N/A	N/A	2.012 mg/L	358 Pa	4.8
ROSMARINUS OFFICINALIS LEAF OIL	N/A	N/A	N/A	0.088 mg/L	9 Pa	6.23
EUGENIA CARYOPHYLLUS LEAF OIL	N/A	N/A	N/A	insoluble	N/A	N/A
ALLANTOIN	1-(2,5-dioximidazolidin-4-yl)urea	C4H6N4O3	58.12	4.9 g/L	<0.001 Pa	-2.26
THYMUS VULGARIS EXTRACT	N/A	N/A	0	Insoluble	N/A	N/A
JUGLANS REGIA LEAF EXTRACT	N/A	N/A	N/A	N/A	N/A	N/A

SALVIA OFFICINALIS LEAF EXTRACT	N/A	N/A	N/A	N/A	N/A	N/A
QUERCUS ROBUR BARK EXTRACT	N/A	0	N/A	soluble	N/A	N/A
ALCOHOL	ethanol	C2H6O	46.07	789 000 mg/L, completely miscible in water	5 726 Pa	-0.35
WATER	Water	H2O	18.02 g/mol	N/A	N/A	N/A

b) Physical and chemical characteristics of the cosmetic product

Parameter	Value
Apearance	Liquid
Colour	Beige-brown
Odour	Pleasant
pH	5.17
Density	N/A
Viscosity	5.18cP

c) Cosmetic product's stability

A laboratory study of stability and compatibility was performed, and the study protocol and results are attached to this document.

Stability study protocol description:

- Accelerated test, in original packaging, for 12 weeks, on +5C, room temperature (25C) and +40C,.
- Centrifugation measurement
- Measurement of organoleptic parameters (color, odor), pH, viscosity and phase separation were measured periodically
- Microbiology testing at the beginning and at the end of the test period

Based on the results obtained, it is concluded that the product FUNGOMAX is stable and compatible with the packaging under reasonably applicable conditions of use and storage on 5C, 25C and 40C.

3) Microbiological quality

a) Microbiological quality of raw materials

Raw materials used for FUNGOMAX are followed with technical documentation (SDS, TDS, CoA, etc.) provided by the supplier which is available and attached to this document. The technical documentation was reviewed, based on which it was concluded that all the ingredients of the cosmetic product represent a satisfactory microbiological quality. The production of the cosmetic product is carried out in accordance with the principles of good manufacturing practice, which excludes the possibility of microbiological contamination during the production process.

b) Microbiological quality of the cosmetic product

Quantitative and qualitative limits are based on the European Standard EN ISO 17516:2014 Cosmetics – Microbiology – Microbiological limits. The European Standard EN ISO 17516:2014 was approved by CEN on 9 August 2014 and at present is widely used 149 by the cosmetics industry as international standard as given (SCCS Notes of Guidance 12th revision):

Types of microorganism	Products specifically intended for children under three years of age, the eye area or the mucous membranes	Other products
Total Aerobic Mesophilic Microorganisms (Bacteria plus yeast and mould)	$\leq 1 \times 10^2$ CFU per g or ml ^a	$\leq 1 \times 10^3$ CFU per g or ml ^b
<i>Escherichia coli</i>	Absence in 1 g or 1 ml	Absence in 1 g or 1 ml
<i>Pseudomonas aeruginosa</i>	Absence in 1 g or 1 ml	Absence in 1 g or 1 ml
<i>Staphylococcus aureus</i>	Absence in 1 g or 1 ml	Absence in 1 g or 1 ml
<i>Candida albicans</i>	Absence in 1 g or 1 ml	Absence in 1 g or 1 ml
Due to inherent variability of the plate count method, according to USP Chapter 61 or EP Chapter 2.6.12, Interpretation of results, results considered out of limit if a > 200 CFU/g or ml, b > 2 000 CFU/g or ml. NOTE When colonies of bacteria are detected on Sabouraud Dextrose agar, Sabouraud Dextrose agar containing antibiotics may be used.		

Microbiological testing of the cosmetic product FUNGOMAX was conducted. The results are proving the compliance of the product with the limits named above for the category „Other products“. The laboratory results report is attached to this document.

Challenge test: There is no need to conduct the efficacy of preservation system for cosmetic product based on the exclusion criteria as stated in ISO 29621:2017 Cosmetics — Microbiology — Guidelines for the risk assessment and identification of microbiologically low-risk products standard (alcohol content>20%).

4) Impurities, traces, information about the packaging material

a) Impurities and traces in the raw materials

Raw materials used for the manufacturing of the cosmetic product FUNGOMAX are well-known chemical substances and mixtures that have long been used as raw materials in the cosmetic industry.

Based on the technical documentation (SDS, TDS, CoA, etc.) provided by the suppliers of used raw materials, there are no impurities and traces for which the Margin of Safety (MoS) should be calculated.

b) Heavy metals content

Heavy metals content of the cosmetic product FUNGOMAX was determined for Pb, Cd, As, Cr, Hg, and Ni. The results are given as less than 0.5mg/kg for each component which is considered acceptable. The laboratory results report is attached to this document.

Cosmetic product FUNGOMAX is produced in Republic of Serbia where heavy metals content determination is mandatory. Results obtained are acceptable according to the limits set by the cosmetic regulation in Republic of Serbia.

c) The relevant characteristics of packaging material

Cosmetic product FUNGOMAX was packed in a glass cosmetic bottle with a dropper. The bottle is being placed in a cardboard box. Elements used for packaging described are well known and are being widely used as packaging for pharmaceutical and cosmetic products for a long time.

The packaging supplier submitted the technical documentation for each element that is attached to this document and is describing the materials used. Technical documentation provided is considered adequate for the assessment of suitability of packaging for cosmetic use.

5) Normal and reasonably foreseeable use

Cosmetic product FUNGOMAX is intended to be used as cosmetic lotion for the skin in areas that are prone to the appearance of fungus, without subsequent washing after application.

Instructions for use written on the cosmetic product:

„Drip twice a day and gently rub on the target area.“

Warnings and precautions written on the cosmetic product:

„Hypersensitivity to any of the ingredients. The product is intended for external use only. Do not apply on open wounds. Wash hands after use. Avoid product contact with eyes. If contact occurs, wash with plenty of water. In case of extended irritation, seek medical help. Keep on dry place, away from children.“

The cosmetic product FUNGOMAX is presented in a way that clearly communicates its purpose. Precise and clearly understandable instructions are given, based on which it is considered that it cannot be misused.

6) Exposure to the cosmetic product

Duration of product application	Leave-on
Body part intended for application	Skin and nails
Reasonably foreseeable application on other body parts	N/A
Duration and frequency of application	2 times a day
Relative daily exposure (mg/kg bw/day)*	48.28
Target population	Adults
Retention factor	1

*estimated according to the SCCS Notes of Guidance for the testing of cosmetic ingredients and their safety evaluation (12th revision)

7) Exposure to the cosmetic products ingredients

Systemic Exposure Dose (SED) was calculated using the formula:

$$\text{SED (mg/kg bw/day)} = A \text{ (mg/kg bw/day)} * C \text{ (\%)/100} * DA \text{ (\%)/100}$$

Where parameters are:

A (mg/kg bw/day) – Relative daily exposure

C (%) – Maximum concentration of the ingredient in the final product

DA (%) – Dermal absorption

Dermal absorption (DA) and no observed adverse effect level (NOAEL) values were determined by detailed analysis of the toxicological profiles of substances that are given in the part 8. If oral bioavailability and dermal absorption values were not known, the default value of 50% was used.

Margin of safety (MoS) was calculated using the formula:

$$\text{MoS} = \text{NOAEL} / \text{SED}.$$

INCI naziv sastojka	Maksimalna koncentracija sastojka u proizvodu (%)	SED (mg/kg bw/day)	NOAEL (mg/kg bw/day)	MoS
GLYCERIN	20	4.828	1000	736.4448
LAVANDULA HYBRIDA OIL	0.15	0.07242	160	690.417
ROSMARINUS OFFICINALIS LEAF OIL	0.15	0.07242	150	2071.251
EUGENIA CARYOPHYLLUS LEAF OIL	0.1	0.02414	2.5	103.5626
ALLANTOIN	0.15	0.07242	N/A	N/A
THYMUS VULGARIS EXTRACT	11.9175	2.876885	N/A	N/A
JUGLANS REGIA LEAF EXTRACT	7.945	3.835846	N/A	N/A
SALVIA OFFICINALIS LEAF EXTRACT	4.767	0.690452	N/A	N/A
QUERCUS ROBUR BARK EXTRACT	3.178	0.767169	100	130.3493
ALCOHOL	31.78	7.671692	1200	156.4192
WATER	31.78	15.34338	N/A	N/A

According to the guidelines of the World Health Organization and the Scientific Committee on Consumer Safety (SCCS), a Margin of Safety (MoS) value of at least 100 is considered a sufficient indicator of safe use of the substance.

8) Toxicological profile of the substances

INCI	ALCOHOL
Presence in the annexes	0
Description	According to the International Cosmetic Ingredient Dictionary and Handbook (Gottschalk and McEwen 2004), Alcohol is the name given by convention to ethanol as used in cosmetics, and Alcohol Denat. is the name given to denatured Alcohol for use in cosmetics. Alcohol Denat. is prepared by adding one or more denaturing agents to potable Alcohol, making it unfit for

	beverage or internal human medicinal use. Alcohol may be completely denatured (CD Alcohol) or specially denatured (SD Alcohol) depending on the type(s) of denaturant(s) used.
Oral toxicokinetics	Following any route of intake resulting in an elevated blood ethanol level . Inhalation: In a study to determine the fractional uptake of ethanol during inhalation exposure, human volunteers were exposed to ethanol vapour concentrations of 10 -12mg/l (6000ppm) for periods of 1 hour. Experiments were done with and without a prior dose of ethanol by the oral route. The fractional absorption was typically around 55% (variation less than 10% within and between individuals) and was found to be not detectably affected by variations in tidal volume, breathing frequency nor the existing blood ethanol concentration. The concentration of ethanol in expired air did not fall below 30%.
Inhalation toxicokinetics	In a study to determine the fractional uptake of ethanol during inhalation exposure, human volunteers were exposed to ethanol vapour concentrations of 10 -12mg/l (6000ppm) for periods of 1 hour. Experiments were done with and without a prior dose of ethanol by the oral route. The fractional absorption was typically around 55% (variation less than 10% within and between individuals) and was found to be not detectably affected by variations in tidal volume, breathing frequency nor the existing blood ethanol concentration. The concentration of ethanol in expired air did not fall below 30%.
Dermal absorption	In a study to assess the skin penetration potential of ethanol, an in vitro study was carried out using excised pig skin and radiolabelled ethanol. Ethanol penetration was greater under occlusion conditions than non-occlusive conditions, as might be expected. Absorption rates were around 21% and 1% of applied doses respectively.
Skin irritation	not irritating
Eye irritation	not irritating
Skin sensitisation	not sensitising
Acute toxicity	In a guideline acute oral toxicity study, the LD50 was determined to be 10470 mg/kgbw when dosed as a 95% solution in water. Dermal: LD50 17 100 mg/kg bw. Inhalation: In an acute inhalation study that approximated to guideline, ethanol vapour was found to be of very low acute toxicity to male and female rats. The LC50 (4hr) was established to be around the 117 -125mg/l and the LC0 around 62mg/l. It is worthy of note that the LD50 is well above the lower explosive limit (LEL).
Repeated dose studies	Oral treatment of rats for 14 weeks with 0, 5, 10 or 20 mg/Kg of a mixture containing 16.25% USP ethanol, The NOAEL for the study is determined at 10 ml/Kg. Most data available on ethanol is via the oral route of exposure. Much is at high doses which limits its value to risk assessment of ethanol as a chemical substance. From the data available, it is possible to surmise that ethanol is of repeat dose low toxicity by the oral route, with a lowest reported NOAEL in animals of 2400 mg/kg for rats.
Mutagenicity/genotoxicity	OECD Guideline 471 (Bacterial Reverse Mutation Assay): negative. In vivo mammalian germ cell study: cytogenicity / chromosome aberration: Ethanol is unlikely to be a dominant lethal mutagen, at least up to the maximum tolerated dose.
Carcinogenicity	Rats: No effect level >3000mg/kg. Mice (B6C3F1): Females: NOAEL>44000mg/kg (cancer), Males: NOAEL>4250mg/kg (based on historic control data), Males BMDL10=1400mg/kg (based on concurrent control data). Overall, there is no significant evidence to warrant a classification of ethanol

	for cancer in the context of the relevant classification and labelling regulations for chemical substances.
Reproductive toxicity	Overall, ethanol in drinking water at concentrations up to 15% (equivalent to 20.7 g/kg/day) had no demonstrable effect on fertility in this two-generation study. Groups of female Sprague-Dawley rats were given liquid diets with 15, 25, or 36% ethanol-derived calories (E15, E25, and E36 groups, respectively), or without ethanol (pair-fed isocaloric (PF15, PF25, PF36) or ad libitum (C) control) for 3 weeks prior to mating, and throughout 21 days of gestation. Prenatal ethanol exposure affected fetal skeletal ossification at exposure levels lower than those required to affect fetal body weight and length, although the significance of these changes for long-term bone health is unknown. No statistically significant effects on fetal growth or ossification were seen in the fetuses of dams ingesting about 5.2 g ethanol/kg bw/day from 3 weeks prior to mating to GD 21.
Photosensitivity	N/A
Endocrine effects	N/A
Reference	1. ECHA https://echa.europa.eu/sl/registration-dossier/-/registered-dossier/16105/7/10/2 2. OECD SIDS, Ethanol, Assessment Report https://hvpchemicals.oecd.org/ui/handler.axd?id=2602cc56-d998-4e67-bd78-5454ef3f8f9a
POD (NOAEL)	2400mg/kg bw/day

INCI	GLYCERIN
Presence in the annexes	0
Description	Glycerol is widely distributed in food as a natural constituent, and it has undergone review and approval for use as both a direct and indirect food additive and is GRAS by US FDA. Animal studies involving single or repeated exposure by various routes indicated a low order of toxicity, the possible target sites (particularly following repeated oral administration) being the kidney and gastrointestinal tract. Glycerol appears to be of generally low oral toxicity in humans.
Oral toxicokinetics	Glycerin is rapidly absorbed in the intestine and the stomach, distributed throughout the extracellular fluids through much of the body. It is mostly metabolized by the liver and kidneys with the remainder excreted in urine. Free glycerin is naturally present in humans, primarily in plasma.
Inhalation toxicokinetics	The NOAEC in rats for local and systemic effects after 13-week exposure is 662 mg/m ³ .
Dermal absorption	0.1
Skin irritation	Not dermally irritating to rabbits when applied at concentrations up to 100% to up to 30% of the body surface. Mild dermal irritant at 100% in guinea pigs.
Eye irritation	Undiluted glycerin was not irritating when administered to the eyes of human subjects. There was a strong burning and stinging sensation, with tear production but no injury was observed.
Skin sensitisation	A moisturizer containing 65.9% glycerin was not sensitizing to human subjects. Natural and synthetic glycerin were not sensitising to white male guinea pigs at 0.1%.
Acute toxicity	The reported oral LD50 of glycerin ranged from 2530-58400 mg/kg in rats, 4090-38000 mg/kg in mice, 27000 mg/kg in rabbits, and 77500 mg/kg in guinea pigs. The dermal LD50 of glycerin in rats was reported to be > 21900 mg/kg and >18700 mg/kg in rabbits. The approximate Lt50 for rats was determined to be 423 min for exposure to glycerin vapors at 11.0 mg/L. The intraperitoneal LD50 of glycerin in rats ranged from 4420-10100 mg/kg and 8600-9500 mg/kg in mice. The subcutaneous LD50 of glycerin was 100 mg/kg

	in rats and ranged from 91-10000 mg/kg in mice. The intravenous LD50 of glycerin ranged from 5200-6600 mg/kg in rats, 4250-6700 mg/kg in mice, and 53000 mg/kg in rabbits.
Repeated dose studies	In the best available dietary study, groups of 22 rats (Long-Evans)/sex/treatment received 5, 10 and 20% glycerol (natural or synthetic) in their diet (males 2000, 4000 and 8000 mg/kg bw; females 2500, 5000 and 10000 mg/kg bw) for 2 years. Although the results were not described in detail, based on this limited dietary study it can be concluded that no adverse effects were observed at up to 10,000 mg/kg bw. Dermal: The effects following repeated dermal application of glycerin was examined. There were no effects noted in rabbits dosed 8 hours/day, 5 days/week for 45 weeks with dose levels as high as 4.0 ml/kg. Using a density of 1.2611 g/cm ³ at 20 °C, a dose of 4.0 ml/kg corresponds to 5040 mg/kg/day.
Mutagenicity/genotoxicity	Glycerin was not genotoxic in multiple Ames tests using multiple strains of <i>S. typhimurium</i> at concentrations up to 50 mg/plate. It was not genotoxic in a cytogenetic assay, X-linked HGPRT, sister chromatid exchange assay, unscheduled DNA synthesis assay, and chromosome aberration test at concentrations up to 1.0 mg/mL.
Carcinogenicity	Glycerin administered in the feed of rats at doses up to 20% in feed for 1 year or up to 10 g/kg for 2 years did not increase the incidence of tumors. Orally administered glycerin, in concentrations up to 5%, had a potentiating effect on the carcinogenicity of 4NQO in mice.
Reproductive toxicity	No convincing evidence of reproductive effects have been seen in rats treated orally or dermally, or in mice fed glycerol during pregnancy. Glycerin is transferred across the placenta in small amounts. May cause some adverse reproductive effects based on animal data, but there was no evidence of teratogenicity. No effects on fertility and reproductive performance were observed in a two generation study with glycerol administered by gavage (NOAEL 2000 mg/kg bw/day).
Photosensitivity	N/A
Endocrine effects	N/A
Reference	(1) CIR Expert Panel. Safety assessment of glycerin as used in cosmetics - final report (January 14, 2015). (2) ECHA https://echa.europa.eu/sl/registration-dossier/-/registered-dossier/14481/7/6/1
POD (NOAEL)	reproduction 2000, oral 10000, dermal 5040

INCI	LAVANDULA HYBRIDA OIL
Prilozi Pravilnika o kozm.proizvodima	None
Opis supstance	Essential oil of lavender hybrid.
Toksikokinetika oral	1
Toksikokinetika inhalacija	1
Dermalna apsorpcija	1
Iritacija kože	not irritant
Iritacija oka	irritant
Senzitizacija kože	sensitizer
Akutna toksičnost	LD50>5000 mg/kg bw
Studije ponovljenih doza	In a GLP 28-day repeated dose toxicity study (similarly to OECD Guideline 407) with read-across substance coriander oil, NOAEL rats = 160 mg/kg bw/d (males and females)
Mutagenost/genotoksičnost	„Ames test: non mutagenic, Micronucleus test in human lymphocytes: non clastogenic
Karcinogenost	N/A

Reproduktivna toksičnost	In a GLP reproductive and developmental toxicity screening study (similarly to OECD Guideline 421) in rats with read-across substance coriander oil, NOAEL for maternal toxicity and developmental toxicity = 500 mg/kg bw/day
Fotosenzitivnost	N/A
Endokrina aktivnost	N/A
Reference	1. ECHA https://echa.europa.eu/registration-dossier/-/registered-dossier/21156/7/9/1
Odabrani POD (NOAEL)	160mg/kg bw/day, 28 days, rat, oral

INCI	Eugenia Caryophyllus Leaf Oil
Prilozi Pravilnika o kozm.proizvodima	None
Opis supstance	According to the European Pharmacopoeia (1/2008: 1091), Caryophylli flos consists of the whole flower buds of Syzygium aromaticum (L.) Merril et L.M. Perry (Syn. Eugenia caryophyllus (C. Spreng.) Bull. et Harr.) which were dried until they become reddish-brown. They contain not less than 150 ml/kg essential oil. Three main components account for nearly 99% of the essential oil: eugenol (75–88%), acetyl eugenol (4–15%), β -caryophyllene (5–14%). The medicinal use of Caryophylli floris aetheroleum can be traced in literature back to the 15th century. Clove essential oil acts in high concentrations as local irritant, allergic reactions may also be possible. However, when applied in diluted form, no reports on severe adverse events are published. When applied correctly in the proposed routes of administration, clove essential oil can be considered as clinically safe.
Toksikokinetika oral	N/A
Toksikokinetika inhalacija	N/A
Dermalna apsorpcija	N/A
Iritacija kože	In concentrated form, oil of clove may be irritating to mucosal tissues (Gruenwald et al 2004). In contrast to this Anton et al (2001) report that there is no skin irritation (undiluted oil) on hairless mice. Under occlusion the undiluted clove oil was moderately irritating in rabbits.
Iritacija oka	0
Senzitizacija kože	In a patch test study a 10% ethanol extract of Caryophylli flos was investigated among other herbal preparations used in the Traditional Chinese Medicine. Out of 30 patients 8 reacted positively to clove extract (Chen et al 2003).
Akutna toksičnost	LD50 1.8–3.72 g/kg
Studije ponovljenih doza	Clove essential oil in oral dosages of 35 or 70 mg per animal (rat) over 8 weeks was tolerated without signs of toxicity. Higher doses led to inactivity and weight loss. 105 mg/animal p.o. daily for 2 to 3 weeks led to serious liver and kidney damage and death of the animals. Clove oil is allowed as food additive and therefore an administration to food-producing animals is possible. The allowed daily intake for the United States is reported with 2.5 mg/kg
Mutagenost/genotoksičnost	The dry residue of aqueous and methanolic extracts showed mutagenic effects in the rec assay in Bacillus subtilis. The mutagenic activity in the Ames test on Salmonella typhimurium TA98 and TA100 was not assessable due to the antimicrobial action. An in vivo bone marrow micronucleus assay demonstrated that the administration of 0.5% and 2% of cloves in the diet of mice for 10 days neither significantly induced micronuclei nor could effectively modulate the 7, 12-dimethylbenz[a]anthracene genotoxicity in mice

Karcinogenost	Rats: eugenol was administered in the diet in a concentration of up to 6000 ppm. Considering the mean diet consumption per day the daily uptake of eugenol was for male rats about 1100 mg/kg body weight at the beginning to 245 mg/kg body weight at the end of the study; for female rats about 790 mg/kg body weight at the beginning to 237 mg/kg body weight at the end of the study. The study outcome was negative in both sexes.
Reproduktivna toksičnost	Domaracky et al (2007) investigated the effects of clove essential oil on the growth and development of mouse pre-implantation embryos in vivo. The animals received 0.25% essential oil in the commercial diet (= 375 mg/kg per day). Treatment with clove essential oil induced a significantly higher percentage of dead cells compared to the control group. Female CD-1 mice were given clove oil from day 6 to day 15 of gestation p. o. 2.2 to 215 mg/kg body weight daily. The fetuses were obtained on day 17. The use of clove oil had no apparent toxic effect on the implantation and survival of the mother and foetus. The number of malformations of the soft tissues and the skeletal system was not different from that in the control group spontaneously occurring.
Fotosenzitivnost	N/A
Endokrina aktivnost	N/A
Reference	1. EMA Assessment report on Syzygium aromaticum (L.) Merrill et L.M. Perry, flos and Syzygium aromaticum (L.) Merrill et L.M. Perry, floris aetheroleum
Odabrani POD (NOAEL)	2.5mg/kg bw/day ADI

INCI	ROSMARINUS OFFICINALIS LEAF OIL
Prilozi Pravilnika o kozm.proizvodima	None
Opis supstance	The essential oil obtained from the flowering tops and leaves of R officinalis. R officinalis (rosemary)-derived ingredients are strongly aromatic.
Toksikokinetika oral	0.5
Toksikokinetika inhalacija	1
Dermalna apsorpcija	0.5
Iritacija kože	irritating
Iritacija oka	irritating
Senzitizacija kože	sensitising
Akutna toksičnost	LD50 = 5000 mg/kg bw
Studije ponovljenih doza	Repeated dose toxicity using read-across from Eucalyptus oil (OECD TG 422): NOAEL = 300 mg/kg bw, 28 days
Mutagenost/genotoksičnost	Based on this read-across Rosemary oil is considered not mutagenic in mammalian cells.
Karcinogenost	The genotoxicity of the substance has been assessed in vitro for three genotoxicity endpoints: genotoxicity in bacteria,

	cytogenicity in a chromosomal aberration assay and genemutation test in mammalian cells, resulting in the absence of genotoxicity. Therefore no genotoxic carcinogenicity is expected,
Reproduktivna toksičnost	Repeated dose toxicity using read-across from Eucalyptus oil (OECD TG 422): NOAEL = 300 mg/kg bw, 28 days
Fotosenzitivnost	N/A
Endokrina aktivnost	N/A
Reference	1. ECHA https://echa.europa.eu/registration-dossier/-/registered-dossier/5479/7/9/1 2. CIR Expert Panel, Safety Assessment of Rosmarinus officinalis (Rosemary)-Derived Ingredients as Used in Cosmetics
Odabrani POD (NOAEL)	300mg/kg bw/day, 28 days, rat

INCI	ALLANTOIN
Prilozi Pravilnika o kozm.proizvodima	N/A
Opis supstance	Allantoin is a heterocyclic organic compound. .Allantoin, (2,5-Dioxo-4-imidazolidinyl) urea or 5-Ureidohydantoin, is a colorless crystalline powder that is soluble in water at approximately 0.6%. It is endogenous in many mammals and some plants. In most mammals, it is the product of uric acid oxidation by purine catabolism. Potential exposure routes include dermal, oral, inhalation and intravenous, with dermal and oral predominating.
Toksikokinetika oral	Absorption of Allantoin is route and vehicle dependent. Dermal absorption was as much as 16%, other routes resulted in absorption in range of 0.2% to 1.5% of the administered dose.
Toksikokinetika inhalacija	N/A
Dermalna apsorpcija	Allantoin absorption from the hydrophilic gel vehicle was 5% after 3 hr., and 6.9% after 6 hr. Similarly, absorption was 13% at 3 hr. and 15.4% at 6 hr. for the oil/water emulsion. No evidence of skin irritation or corrosion was reported.
Iritacija kože	not irritating
Iritacija oka	not irritating
Senzitizacija kože	not sensitising
Akutna toksičnost	Oral and dermal LD50 5 000 mg/kg bw
Studije ponovljenih doza	Annex XI to the REACH Regulation provides for the waiver of additional animal testing in scenarios where adequate data exist, and further animal testing is not scientifically necessary to further the safety argument for the test substance. Given the preponderance of evidence, as well as the availability of a 2-year tumorigenicity study, it is scientifically unnecessary to

	generate additional mammalian data for Allantoin. Allantoin is the end product of purine metabolism in most mammals, excluding humans and some non-human primates; - Humans lack the enzyme necessary to convert uric acid to Allantoin, where most test species have this enzyme; - Allantoin is endogenous in standard test species and thus chronic exposure is not of relevance; - Allantoin is endogenous in humans as well;
Mutagenost/genotoksičnost	Allantoin was found to be non-mutagenic in the Ames Assay in the <i>S. typhimurium</i> TA1535, TA1538, TA98, TA100 bacterial strains.
Karcinogenost	Lijinsky76fed F344 rats (males and females) rat chow with and without 0.2% allantoin with or without 0.2% sodium nitrite ad libitum for 106 weeks (n=20). The rats were necropsied and all gross lesions and major organs and tissues were examined histologically. Controls (n=24) were fed untreated chow and tap water. The study was not performed according to OECD guidelines, however, the study was conducted by a reputable U.S. Governmental laboratory and published as a peer-reviewed paper in a renowned journal. Therefore it is considered to be sound and rated reliable with restrictions. There were no significant differences in survival between the treated groups and their controls. The following table depicts the incidence of tumors in rats given feed supplemented with Allantoin for 106 weeks and in control rats:
Reproduktivna toksičnost	N/A
Fotosenzitivnost	N/A
Endokrina aktivnost	N/A
Reference	1. ECHA https://echa.europa.eu/sl/registration-dossier/-/registered-dossier/13641/7/9/1 2. CIR Expert Panel Final Report of the Safety Assessment of Allantoin and Its Related Complexes
Odabrani POD (NOAEL)	N/A

INCI	THYMUS VULGARIS EXTRACT
Prilozi Pravilnika o kozm.proizvodima	None
Opis supstance	Thymus vulgaris L. is widely used as an ingredient in cooking and in herbal medicine. The oral administration of Thyme herb preparations can be regarded as safe, especially at therapeutic doses proposed due to their long-standing use. Data from clinical trials support

	the safe use of herbal preparations of Thyme in the indication 'cough associated with cold'. The reported adverse events are mild.
Toksikokinetika oral	N/A
Toksikokinetika inhalacija	N/A
Dermalna apsorpcija	N/A
Iritacija kože	Irritant
Iritacija oka	Irritant
Senzitizacija kože	Sensitizer
Akutna toksičnost	The LD50 of the essential oil oral route is 2.84 g/kg body weight in rats
Studije ponovljenih doza	The results of the 28-day oral toxicity test suggest that the no-observed-adverse effect level (NOAEL) is greater than 250 mg/kg/day.
Mutagenost/genotoksičnost	Thyme essential oil had no mutagenic or DNA-damaging activity in either the Ames test (strains TA1535, TA1537, TA98, TA100, with and without metabolic activation, 0.25 and 0.5 µl essential oil/plate) or Bacillus subtilis rec-Assay
Karcinogenost	N/A
Reproduktivna toksičnost	N/A
Fotosenzitivnost	N/A
Endokrina aktivnost	N/A
Reference	1. EMA, Assessment report on Thymus vulgaris L., vulgaris zygis L., herba
Odabrani POD (NOAEL)	N/A

INCI	JUGLANS REGIA LEAF EXTRACT
Prilozi Pravilnika o kozm.proizvodima	None
Opis supstance	The leaves contain approximately 10% tannins of the ellagitannins type (Blumenthal 2000). According to DAC not less than 2%, calculated as pyrogallol (Wichtl, 2004). The leaves contain about 3.4% of flavonoids, especially quercetine. The most known naphthoquinone constituent is juglone (5-hydroxy-1,4-naphthoquinone) which occurs in fresh plant. There are no reports of adverse effects of walnut leaves from Member States. The available data are sufficient to include the traditional use of specified preparations of Juglans regia leaves in a monograph of the European Community. European walnut leaves fulfil the requirement of safe therapeutic use for at least 30 years .
Toksikokinetika oral	N/A
Toksikokinetika inhalacija	N/A
Dermalna apsorpcija	N/A

Iritacija kože	possible irritant
Iritacija oka	possible irritant
Senzitizacija kože	Allergic reactions following contact with walnuts are common.
Akutna toksičnost	Juglon LD50 rat 112 mg/kg
Studije ponovljenih doza	N/A
Mutagenost/genotoksičnost	The evaluation of genotoxicity of the <i>Juglans regia</i> extracts showed significant damage in the comet assay in concentrations above 250 ppm but not in the VITOTOX test. The extracts in the Ames assay showed potent antimutagenic effects against the direct acting mutagens NPD, sodium azide, and the S9-dependent mutagen 2-AF (Arora et al. 2005). The authors conclude, that the plant extract can be mutagenic as well as antimutagenic depending on the test system used. It can be concluded that despite that juglone as a quinone compound has been linked to mutagenic action, it is unstable and in dry leaves it is present only in traces. On the basis of existing findings and comparisons with other quinones, genotoxicity seems not conclusively evaluated and may be limited, weak and probably meaningless to humans following exposure to small amounts of juglone.
Karcinogenost	N/A
Reproduktivna toksičnost	0
Fotosenzitivnost	0
Endokrina aktivnost	0
Reference	1. EMA Assessment report on <i>Juglans regia</i> L., folium
Odabrani POD (NOAEL)	N/A

INCI	SALVIA OFFICINALIS LEAF EXTRACT
Prilozi Pravilnika o kozm.proizvodima	None
Opis supstance	According to the EMA monograph, the principal components of <i>Salvia officinalis</i> oil are thujone, 1,8-cineole, and camphor. The main classes of constituents in <i>Salvia officinalis</i> oil are identified as terpenoids, hydroxycinnamic acid derivatives, flavonoids, phenolic glycosides, and polysaccharides. According to the 49th Amendment of the International Fragrance Association (IFRA) standards, in the absence of analytical data, thujone is expected to occur naturally from 2-10% in <i>Salvia officinalis</i> oleoresin and from 8 – 33% in <i>Salvia officinalis</i> oil. CIR Expert Panel noted the GRAS status and historical food uses of <i>Salvia officinalis</i> (sage)-derived ingredients, especially <i>Salvia officinalis</i> leaves, and agreed that systemic exposures from food would be much higher than those from cosmetic use.

Toksikokinetika oral	N/A
Toksikokinetika inhalacija	N/A
Dermalna apsorpcija	N/A
Iritacija kože	irritating
Iritacija oka	irritating
Senzitizacija kože	sensitising
Akutna toksičnost	LD 50> 2000 mg/kg bw
Studije ponovljenih doza	In an 8-wk study using groups of 5 white rats (sex and strain not specified), a daily dose of 250 mg/kg bw Salvia officinalis oil was well tolerated when given by oral administration. Upon increase of the daily dose to 500 mg/kg bw/d, convulsions occurred in some animals. Upon increase to 1000 mg/kg bw/d, most animals died, and all animals died when the dose was increased to 1250 mg/kg bw/d (timing and duration of all 3 dose increases not provided). The no-observed adverse effect level (NOAEL) of Salvia officinalis oil was determined to be 250 mg/kg bw/d, under the conditions of this study.
Mutagenost/genotoksičnost	Ames test: non mutagenic
Karcinogenost	N/A
Reproduktivna toksičnost	Twenty-four female ICR mice received a daily dose of 0.25% Salvia officinalis oil (equivalent to 375 mg/kg/d), in rodent feed, for 14 d. A significant decrease in embryo cell distribution, according to nucleus number, was observed in dams which consumed Salvia officinalis oil (until day 4 of gestation), compared to control females fed with 1% edible soya oil. Dam weights and the proportion of dead cells in embryos were not affected. Groups of 7 female Wistar rats were administered either distilled water or 30 mg/kg bw/d hydroalcoholic extract (70% ethanol) of Salvia officinalis, orally, for 30 d. No significant differences in blood estradiol or progesterone were observed, and a decrease in the duration of estrous cycles in Salvia officinalis extract-treated rats was not statistically significant. An increased number of alveolar buds and lobules in the whole mount slides, as well as an increase in the number and diameter of ducts in the histological sections of rats treated with Salvia officinalis extract, were statistically significant.
Fotosenzitivnost	N/A
Endokrina aktivnost	N/A
Reference	1. ECHA https://echa.europa.eu/registration-dossier/-/registered-dossier/25026/7/7/1 2. CIR Expert Panel Safety Assessment of Salvia officinalis (Sage)-Derived Ingredients as Used in Cosmetics
Odabrani POD (NOAEL)	N/A

INCI	QUERCUS ROBUR BARK EXTRACT
Prilozi Pravilnika o kozm.proizvodima	None
Opis supstance	In the European Pharmacopoeia, it is described as the cut and dried bark of young branches and the lateral shoots of <i>Quercus robur</i>. It contains a minimal amount of 3% of tannins, expressed as pyrogallol, calculated with reference to the dried herbal. The oak bark contains highly variable amount of tannins (8-20%). The tannin content depends on the time of the harvest, age of the branches and on the method of assay used. Tannins are polyphenolic secondary metabolites of higher plants. They comprise either: galloyl esters and their derivatives (gallotannins, ellagitannins and complex tannins) or they are oligomeric and polymeric proanthocyanidins and can possess different interflavanyl coupling and substitution patterns. As hydrolysable tannins are considered to present greater risk to health, several cases of acute poisonings of ruminants were noted due to ingestion of fresh leaves of several oak species. The poisoning may be due to hydrolysable tannins and simple phenols. The levels of tannins and condensed tannins were 97.7 mg tannic acid equivalent and 5.8 mg catechin equivalent/g of dry leaves. The ability of tannins to form strong complexes with proteins is considered as their most important activity for both therapeutic and toxicological effects. The available data are sufficient to include the traditional use of specified preparations of oak bark in a Community herbal monograph. Oak bark fulfils the requirement of therapeutic use for at least 30 years (15 years within the European Union, Directive 2004/24/EC). It can be concluded that the benefit/risk assessment for oak bark preparations is positive for use in therapeutical dosages in specific conditions of mild diarrhoea, in minor inflammatory conditions of the oral mucosa or skin and in conditions of medically diagnosed haemorrhoids before anorectal use. Tannic acid is authorised in the EU as a food flavouring, although no data on exposure are available for the European population. The Food and Drug Administration estimates that the added tannic acid in the USA can reach 100 mg/kg in various foods
Toksikokinetika oral	The metabolism of polymeric proanthocyanidins by human colonic microflora has been investigated in vitro in anoxic conditions using ¹⁴C-labelled purified proanthocyanidin (Déprez et al. 2000). Polymers were degraded after 48 h of incubation. As metabolites, low-molecular-weight aromatic compounds: phenylacetic, phenylpropionic and phenylvaleric

	acids were recognized by gas chromatography. The results show, that proanthocyanidins can be metabolized by the colonic microflora into low-molecularweight aromatic acids. Evaluation of polyphenol bioavailability was tested in vitro in isolated segments of the rat small intestine (Carbonaro et al. 2001). As model compounds, tannic acid and catechin were used. The results indicated a significant, concentration-dependent uptake of both compounds by the small intestinal wall. Higher uptake (50%) was recorded for tannic acid compared with catechin (30%). However, only catechin demonstrated a complete transfer through the gut wall, whereas tannic acid remained in the gut wall via interaction with wall proteins. Both polyphenols were bound significantly by proteins in the intestinal lumen.
Toksikokinetika inhalacija	Nasal cancer risk associated with occupational exposure to wood dusts via inhalation cannot be applied to the oral use of oak bark preparations.
Dermalna apsorpcija	N/A
Iritacija kože	N/A
Iritacija oka	Specific IgE to Quercus ilex and Quercus robur and conjunctival challenge test to Quercus robur pollen was positive in two patients
Senzitizacija kože	Sensitisation to pollen of Quercus ilex and Quercus robur was found in two patients suffering from rhinoconjunctivitis and seasonal bronchial asthma between April and June
Akutna toksičnost	Acute toxicity (LD50) of tannic acid given intragastrically to rats was found to be 2.26 ± 0.083 g/kg body weight.
Studije ponovljenih doza	To investigate the response of pancreas to tannins and changes in activities of enzymes in the lumen of the small intestine, broiled cockerels were fed on increasing levels of tannin-containing diet (Ahmed et al. 1991). The birds were fed with a diet consisted mainly of hydrolysable gallotannins for 4 weeks at levels of 13.5, 25 and 50 g/kg. Weight of pancreas was significantly increased with increasing dose of tannins. Activity of trypsin and α -amylase was also dose-dependently increased. Activities of dipeptidase and sucrose α -glucosidase (disaccharidase) in the intestinal mucosa were inhibited. Growth of the birds fed on tannins diet was adversely affected. In two studies with New Zealand male rabbits, inclusion of a tannic acid preparation (from chestnut wood) for 21 days at 5 000 mg or 10 000 mg/kg feed showed no adverse effects on growth or feed conversion, which were significantly better than the control group that was not given tannic acid. The published studies show that levels of tannic acid preparations up to 15 000 mg/kg feed for adult ruminants and 10 000 mg/kg

	feed for laying hens do not result in adverse effects in these species. Rabbits tolerate up to 10 000 mg/kg, pigs for fattening tolerate up to 1 500 mg/kg and chickens for fattening tolerate up to 1 000 mg/kg without adverse effects. Young animals seem to be more sensitive (> 1 500 mg/kg feed was harmful to calves). In the study with weaned piglets, animals receiving 125 mg/kg showed no sign of adverse effects. In all cases, the values shown to be safe are several orders of magnitude higher than the proposed use as feed flavouring.
Mutagenost/genotoksičnost	Oak tannins are genotoxic in cultured human embryonic MRC-5 lung cells (Zhou et al. 1995). Authors evaluated the ability of solvent extracts of natural woods to induce chromosome aberrations. In experiments three concentrations per extract with or without metabolic activation and 100 metaphase cells were examined. No dose dependent activity was found with tested extracts in the presence of S9, but dose-dependent chromosomal and chromatid breaks were caused by oak wood. No metabolic activation was required for their effect. Labieniec and Gabryelak (2003) showed that tannic, ellagic and gallic acids have genotoxic and cytotoxic properties in Chinese hamster cell line B14 when using the Comet assay for detection of DNA damage. Tannic, ellagic and gallic acids were exposed at concentration range of 15 – 240 µM. Tannins decreased viability of the cells, with highest cytotoxicity at the concentration of 60 µM. The formation of DNA single-strand breaks was observed.
Karcinogenost	Nasal cancer risk associated with occupational exposure to wood dusts via inhalation cannot be applied to the oral use of oak bark preparations.
Reproduktivna toksičnost	N/A
Fotosenzitivnost	N/A
Endokrina aktivnost	N/A
Reference	1. EMA Assessment report on Quercus robur L., Quercus petraea (Matt.) Liebl., Quercus pubescens Willd., cortex 2. EFSA Scientific Opinion on the safety and efficacy of tannic acid when used as feed flavouring for all animal species
Odabrani POD (NOAEL)	Hypothetical value of 100mg/kg bw/day as worst case from FDA

9) Undesirable effects and serious undesirable effects

The responsible person is informed about the obligation to collect information on undesirable effects and serious undesirable effects, as well as to enter them in the Cosmetic product safety report.

10) Information on the cosmetic product

The chemical substances and mixtures used to produce cosmetic product FUNGOMAX are widely used as ingredients for cosmetic products, they are used in concentrations that are acceptable from the point of view of safety assessment, based on which it can be concluded that the product will not cause irritation of the skin. Eye irritation may occur in case of contact with the eyes. The product packaging clearly states the warning measure to avoid contact with the eyes. Allergens needed to be labelled are listed in part 1.

Cosmetic product DERMOTIN BIOROID doesn't contain:

- Substances and mixtures that have been tested on animals.
- Intentionally added CMR substances
- Prohibited substances listed in the annexes of the Regulation (EC) 1223/2009
- Intentionally added nanoparticles

Part B. Cosmetic Product Safety Assessment

1) Assessment conclusion

The cosmetic product FUNGOMAX is safe for human health when used under normal or reasonably foreseeable conditions, in accordance with the Cosmetic Regulation (EC) 1223/2009.

2) Labelled warnings and instructions of use

Cosmetic product FUNGOMAX is intended to be used as anal region skin care cream without subsequent washing after application.

Instructions for use written on the cosmetic product:

„Drip twice a day and gently rub on the target area.”

Warnings and precautions written on the cosmetic product:

„Hypersensitivity to any of the ingredients. The product is intended for external use only. Do not apply on open wounds. Wash hands after use. Avoid product contact with eyes. If contact occurs, wash with plenty of water. In case of extended irritation, seek medical help. Keep on dry place, away from children.
“

The cosmetic product FUNGOMAX is presented in a way that clearly communicates its purpose. Precise and clearly understandable instructions are given, based on which it is considered that it cannot be misused.

The cosmetic product FUNGOMAX has characteristics in terms of appearance, odor, packaging design, appearance and content of the labelling and volume, and other characteristics of general appearance that cannot mislead the consumer about the purpose of the product and the way of its use.

3) Reasoning

Safety assessment of the cosmetic product FUNGOMAX was done considering all available relevant data, up to date regulatory requirements and the latest scientific knowledge published. Facts relevant for the safety assessment:

- To successfully assess the safety of the product, information on the body part and body surface of product application, the amount of product applied, duration and frequency of application, normal and reasonably applicable routes of exposure to the product and substances, and the target population were considered.
- All technical documentation that defines the substances and mixtures used to produce the cosmetic product, as well as the technical documentation that defines the packaging used for the packaging of the cosmetic product, is considered adequate for evaluating the product's safety.
- All information regarding the properties of the cosmetic product, its production method and the results of the conducted experimental tests submitted by the responsible person are considered adequate for evaluating the product's safety. Any further change of any parameter relevant to the product safety assessment must be considered by the responsible person and a revision of this document carried out, of which the responsible person has been informed.
- The toxicological profiles of the substances included in the composition of the cosmetic product were generated considering the latest scientific knowledge.

- All substances for which there is a legal concentration limit, and which are listed in relevant annexes of the Regulation (EC) 1223/2009 meet the legal requirements and for them the Margin of Safety (MoS) was not calculated.
- For substances for which there is no legal concentration limit and there are available data on conducted repeated dose studies, based on which it is possible to obtain the value of the No Observed Adverse Effects Level are expected (NOAEL), the Margin of Safety (MoS) was calculated, which for all given substances is greater than 100 which is considered acceptable.
- Based on the data on the potential of ingredients substances to cause skin irritation and sensitization, it was concluded that the product FUNGOMAX is not expected to cause skin irritation. It is possible to expect that the product in case of contact with the mucous membrane of the eye will lead to slight eye irritation, which is why it is clearly indicated on the product packaging that contact with the eyes should be avoided.
- There are no expected interactions between the ingredients of the product, as well as interactions between the ingredients of the product and the packaging during the period of 24 months after production, which would lead to negative outcomes regarding the stability and efficiency of the product, which was confirmed by stability and compatibility studies.
- The method of production and packaging of the cosmetic product excludes the possibility of occurrence of microbiological contamination that would lead to endangering human health from the use of the cosmetic product, which was confirmed by the conducted stability and compatibility studies.
- Cosmetic product doesn't contain prohibited substances listed in the Annex II of the Regulation (EC) 1223/2009.
- Based on the overall appearance of the product and the content listed on the packaging, the purpose of the product as skin lotion is clear. Instructions for use and warning measures are clearly indicated on the packaging.

4) Assessor's credentials and approval of part B

Safety Assessor: Vesna Milanković, FVRM Consulting, Koste Jovanovića 21, 11000 Belgrade, Serbia

Academic and expert qualifications:

- M.Sc. Chemical Engineering, 300 ECTS (Faculty of Technology, Univeristy of Novi Sad)
- Specialist Academic Studies, Cosmetology, 60 ECTS (Faculty of Pharmacy, University of Belgrade)
- Safety Assessmet of Cosmetics in the EU, 6 ECTS (Vrije University Brussel)
- Chemical Advisor License, Belgrade.

The conclusion of the safety assessment given in part B of this document about the cosmetic product FUNGOMAX is correct and the explanation provided is adequate. The assessment was done based on the

documentation submitted by the responsible person, up to the current date of updated regulatory requirements and up to the current date of published scientific findings.

Signature:



Vesna Milanković