Foundations of Machine Learning Al2000 and Al5000

FoML-02 Probability - Bayes Theorem and Independence

> <u>Dr. Konda Reddy Mopuri</u> Department of AI, IIT Hyderabad July-Nov 2025





So far in FoML

• What is ML?





So far in FoML

- What is ML?
- Learning Paradigms









 Provides a consistent framework for the quantification and manipulation of "Uncertainty"





- Provides a consistent framework for the quantification and manipulation of "Uncertainty"
- Where does this 'Uncertainty' come from?





Uncertainty in ML

Measurement Noise





Uncertainty in ML

- Measurement Noise
- Finite size of the datasets





Frequentist Interpretation





- Frequentist Interpretation
 - Fraction of times the event occurs





Bayesian Approach





- Bayesian Approach
 - o Quantification of plausibility or strength of the belief of an event





- Bayesian Approach
 - o Quantification of plausibility or strength of the belief of an event
 - Modeling based approach





- Bayesian Approach
 - o Quantification of plausibility or strength of the belief of an event
 - Modeling based approach
 - Plays a central role in this course





Random Variable

• Stochastic variable sampled from a set of possible outcomes





Random Variable


- Stochastic variable sampled from a set of possible outcomes
- Discrete or Continuous






Random Variable

- Stochastic variable sampled from a set of possible outcomes
- Discrete or Continuous
- Probability distribution $\rho(X)$










Random Variable - Example (discrete)

Throwing a dice







Random Variable - Example (discrete)

• Flipping a coin $\{H, 7\}$






- X
- Y






- N trails: sample both







Joint probability









Joint probability

$$p(X = x_i, Y = y_j) = (ij)$$





• If I am interested only on X








- If I am interested only on X
- Marginal probability of X

$$p(X = x_i) = \frac{\zeta_i}{N}$$

$$\zeta_i = \frac{\zeta_i}{N}$$








Sum rule of Probability

$$p(X = x_i) = \sum_{j=1}^{3} p(X = x_i, Y = y_j)$$








Conditional Probability

Conditional probability of Y given X

$$p(Y = y_j(X = x_i)) =$$








Product Rule of probability

$$p(Y = y_j/X = x_i) =$$








Product Rule of probability


$$p(Y = y_j/X = x_i) =$$



$$p(Y = y_j, X = x_i) = p(Y = y_j/X = x_i) \cdot p(X = x_i)$$
















- X
- Y
- 60 trails sample both






• Marginal distribution $\rho(Y)$
















• Marginal distribution $\rho(X)$


















Conditional distribution of X








Example: Marginal & Conditional distributions







Example: Marginal & Conditional distributions












ρ(X): Probability density over X






- ρ(X): Probability density over X
- Probability of x falling in (x, x+dx)
- Probability over a finite interval (a, b)



P(n)





Data-driven Intelligence
& Learning Lab

- Non-negativity
- Normalization





Change of variables






- Change of variables
- $\bullet \quad \times = O(\lambda)$







- Change of variables
- $\bullet \quad \times = O(\lambda) \quad \sim$
- Probabilities in (x, x+dx) must be transformed to (y, y+dy)


$$P(y) dy = P(x) dx$$

$$P(x) - P(y) dy$$



Data-driven Intelligence & Learning Lab

Cumulative distribution function





భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్ भारतीय प्रौद्योगिकी संस्थान हैदराबाद Indian Institute of Technology Hyderabad DiL Data-driven Intelligence & Learning Lab

Rules of Probability Theory

	Discrete	Continuous
Additivity	$p(X \in A) = \sum_{x \in A} p(x)$	pla) dr
Positivity	$p(x) \ge 0$	$p(x) \geq 0$
Normalization	2 p(n)=/	$\int_{\mathcal{X}} p(x) dx = 1$
Sum Rule	$p(x) = \sum_{y \in Y} p(x, y)$)(n)= [p(my) dy
Product Rule	$p(x,y) = p(x/y) \cdot p(y)$	$p(x,y) = p(x/y) \cdot p(y)$





Bayes theorem





Bayes Theorem

Product rule

$$p(x,y) = p(x/y) \cdot p(y)$$








Bayes Theorem

- Product rule
- Symmetry property
- Bayes rule
- Denominator



Data-driven Intelligence & Learning Lab



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్ भारतीय प्रौद्योगिकी संस्थान हैदराबाद

Indian Institute of Technology Hyderabad





$$p(y/x) = \frac{p(x/y) \cdot p(y)}{p(x)}$$

- Prior probability
- Posterior probability of Y
- Likelihood of X = x given Y = y
- Evidence for X = x






Example






Random variables








- Random variables
 - o Box B
 - o Fruit F







Prior Box distribution








Prior Box distribution

0

Conditional of F given B

Marginal of F








- Marginals ρ(F=a) = 11/20 & ρ(F=0) = 9/20
- Posterior probability of Box given observed fruit

$$p(B = r/F = o) =$$







Independence





Independent Random variable

 Two random variables X and Y are independent iff measuring X gives no information about Y (and vice versa)





Next Expectation, Variance, and Gaussian Distribution



