

Foundations of Machine Learning AI2000 and AI5000

FoML-02

Probability - Bayes Theorem and Independence

Dr. Konda Reddy Mopuri

Department of AI, IIT Hyderabad

July-Nov 2025



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



So far in FoML

- What is ML?



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



So far in FoML

- What is ML?
- Learning Paradigms



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Probability Theory



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Probability Theory

- Provides a consistent framework for the quantification and manipulation of “Uncertainty”



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Probability Theory

- Provides a consistent framework for the quantification and manipulation of “Uncertainty”
- Where does this ‘Uncertainty’ come from?



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Uncertainty in ML

- Measurement Noise



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Uncertainty in ML

- Measurement Noise
- Finite size of the datasets



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Probability Theory

- Frequentist Interpretation



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Probability Theory

- Frequentist Interpretation
 - Fraction of times the event occurs



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Probability Theory

- Bayesian Approach



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Probability Theory

- Bayesian Approach
 - Quantification of plausibility or strength of the belief of an event



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Probability Theory

- Bayesian Approach
 - Quantification of plausibility or strength of the belief of an event
 - Modeling based approach



Probability Theory

- Bayesian Approach
 - Quantification of plausibility or strength of the belief of an event
 - Modeling based approach
 - Plays a central role in this course



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Random Variable

- Stochastic variable sampled from a set of possible outcomes



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Random Variable

- Stochastic variable sampled from a set of possible outcomes
- Discrete or Continuous




భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad




Random Variable

- Stochastic variable sampled from a set of possible outcomes
- Discrete or Continuous
- Probability distribution $p(X)$



Random Variable - Example (discrete)

- Throwing a dice



Random Variable - Example (discrete)

- Flipping a coin $\{H, T\}$




Two Discrete Random Variables

- X
- Y


Two Discrete Random Variables

- $X = \{x_1, \dots, x_6\}$
- $Y = \{y_1, \dots, y_3\}$
- N trials: sample both



Two Discrete Random Variables


- Joint probability



Two Discrete Random Variables


- Joint probability

$$p(X = x_i, Y = y_j) = \frac{n_{ij}}{N} \quad \forall x_i, y_j$$



Two Discrete Random Variables


- If I am interested only on X



Two Discrete Random Variables


- If I am interested only on X
- Marginal probability of X

$$p(X = x_i) = \frac{c_i}{N}$$
$$c_i = \sum_{j=1}^3 n_{ij}$$



Sum rule of Probability


$$p(X = x_i) = \sum_{j=1}^3 p(X = x_i, Y = y_j)$$



Conditional Probability


- Conditional probability of Y given X

$$p(Y = y_j | X = x_i) = \frac{n_{ij}}{c_i}$$




Product Rule of probability

$$p(Y = y_j / X = x_i) =$$



Product Rule of probability


$$p(Y = y_j / X = x_i) =$$




$$p(Y = y_j, X = x_i) = p(Y = y_j / X = x_i) \cdot p(X = x_i)$$



Example: Marginal & Conditional distributions




Example: Marginal & Conditional distributions



- X
- Y
- 60 trials - sample both



Example: Marginal & Conditional distributions




- Marginal distribution $p(Y)$


$$P(Y) = \sum_{x \in X} p(x, y)$$




Example: Marginal & Conditional distributions




Example: Marginal & Conditional distributions




- Marginal distribution $p(X)$



Example: Marginal & Conditional distributions



Example: Marginal & Conditional distributions




- Conditional distribution of X

$$p(Y/x=x_i)$$

$$p(X/Y=y_j)$$



Example: Marginal & Conditional distributions



Example: Marginal & Conditional distributions

$$\sum_{y \in Y} p(Y = y_i / X = x_i) = ?$$

Handwritten notes and arrows indicate the relationship between the printed formula and the handwritten formula below it.

$$\sum_{y \in Y} p(Y = y_i / X = x_j) =$$

The handwritten formula is underlined. An arrow points from the x_i in the printed formula to the x_j in the handwritten formula. Another arrow points from the $y \in Y$ in the printed formula to the $y \in Y$ in the handwritten formula. A third arrow points from the $= ?$ in the printed formula to the $=$ in the handwritten formula. There is also a handwritten x to the right of the handwritten formula.



Continuous Random Variable



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Continuous Random Variable

- $p(X)$: Probability density over X




భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Continuous Random Variable

- $p(X)$: Probability density over X
- Probability of x falling in $(x, x+dx)$
- Probability over a finite interval (a, b)

$$p(x) \geq 0$$
$$\int_a^b p(x) dx$$
$$\int_a^b p(x) dx =$$



Continuous Random Variable

- Non-negativity
- Normalization



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Continuous Random Variable

- Change of variables



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad




Continuous Random Variable

- Change of variables
- $x = g(y)$



Continuous Random Variable

- Change of variables
- $x = g(y)$ ✓
- Probabilities in $(x, x+dx)$ must be transformed to $(y, y+dy)$



$$P_y(y) dy = P_x(x) dx$$

$$P_x(x) = P_y(y) \left| \frac{dy}{dx} \right|$$




Continuous Random Variable

- Cumulative distribution function

$$P(x) = P(X \leq x)$$

$$P(x) = \int_{-\infty}^x p(x) dx$$

$$p(x) = \frac{d}{dx} P(x)$$



Rules of Probability Theory

	Discrete	Continuous
Additivity	$p(X \in A) = \sum_{x \in A} p(x)$	$\int_{x \in X} p(x) dx$
Positivity	$p(x) \geq 0$	$p(x) \geq 0$
Normalization	$\sum_{x \in X} p(x) = 1$	$\int_{\mathcal{X}} p(x) dx = 1$
Sum Rule	$p(x) = \sum_{y \in Y} p(x, y)$	$p(x) = \int_{\mathcal{Y}} p(x, y) dy$
Product Rule	$p(x, y) = p(x/y) \cdot p(y)$	$p(x, y) = p(x/y) \cdot p(y)$



Bayes theorem



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Bayes Theorem

- Product rule

$$p(x, y) = p(x/y) \cdot p(y) \quad \checkmark$$



Bayes Theorem

- Product rule
- Symmetry property
- Bayes rule
- Denominator

$$p(x, y) = p(x/y) \cdot p(y)$$

$$p(y, x) = p(y/x) \cdot p(x)$$

$$p(y/x) = \frac{p(x/y) p(y)}{p(x)}$$

$$\sum_y p(y/x) = 1 = \frac{1}{p(x)} \sum_y p(x/y) p(y)$$

$$\sum_y p(x, y) = p(x) = \sum_y p(y/x) p(x)$$



Bayes Theorem

$$p(\hat{x}/y)$$

$$p(y/x) = \frac{p(x/y) \cdot p(y)}{p(x)}$$

- Prior probability
- Posterior probability of Y
- Likelihood of $X = x$ given $Y = y$
- Evidence for $X = x$



Example




భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad

DiL

Data-driven Intelligence
& Learning Lab

Boxes and Fruits

- Random variables




భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Boxes and Fruits

- Random variables
 - Box - B
 - Fruit - F




భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Boxes and Fruits

- Prior Box distribution



Boxes and Fruits

- Prior Box distribution

$$P(B=r) = 7/10 \quad P(B=b) = 3/10$$

- Conditional of F given B


$$P(F=a/B=r) = 2/8 = 1/4 \quad P(F=a/B=b) = 2/4$$

$$P(F=0/B=r) = 3/4 \quad P(F=0/B=b) = 1/4$$

- Marginal of F

$$P(F=a) = \sum_B P(B,F) = \sum_B P(F/B) P(B) = \frac{4}{10} \cdot \frac{1}{4} + \frac{6}{10} \cdot \frac{3}{4} = \frac{11}{20}$$

$$P(F=0) = \sum_B P(B,F) = \frac{3}{10} \cdot \frac{3}{4} + \frac{2}{10} \cdot \frac{1}{4} = \frac{9}{20}$$




0.4

0.6



Boxes and Fruits

- Marginals $p(F=a) = 11/20$ & $p(F=0) = 9/20$
- Posterior probability of Box given observed fruit



0.4

0.6

$$p(B = r / F = o) =$$

$$\frac{p(F=0 | B=r) \cdot P(B=r)}{P(F=0)} = \frac{\frac{3}{4} \cdot \frac{1}{10}}{\frac{9}{20}} = \frac{3}{10} \cdot \frac{20}{9} = \frac{2}{3} \approx 67\%$$

$$P(B=r) = 40\%$$



Independence



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad



Independent Random variable

- Two random variables X and Y are independent iff measuring X gives no information about Y (and vice versa)

$$\begin{aligned} P(Y/X) &= P(Y) \checkmark & P(X/Y) &= P(X) \\ \frac{P(X,Y)}{P(X)} &= P(Y) \\ P(X,Y) &= \underbrace{P(X)} \underbrace{P(Y)} \end{aligned}$$



Next Expectation, Variance, and Gaussian Distribution



భారతీయ సాంకేతిక విజ్ఞాన సంస్థ హైదరాబాద్
भारतीय प्रौद्योगिकी संस्थान हैदराबाद
Indian Institute of Technology Hyderabad

