

## CS 315, Day 37 – Problem Day

**1.** What is an optimal Huffman code for the following set of frequencies, based on the first 8 non-zero (sometimes 0 is considered the first) Fibonacci numbers? Can you generalize your answer to find the optimal code when the frequencies are the first  $n$  Fibonacci numbers?

a:1    b:1    c:2    d:3    e:5    f:8    g:13    h:21

**2.** You have a deck of cards that consists of one ace, two deuces, three 3's, and on up to nine 9's.

How many cards are there?


What is their sum?

Design a strategy that minimizes the expected number of questions asked in the following game: Someone draws a card from the shuffled deck, which you have to identify by asking questions answerable with yes or no.


**3.** Determine a **Longest Common Subsequence** (LCS) of  $\langle 1, 0, 0, 1 \rangle$  and  $\langle 0, 1, 0, 1 \rangle$ .

**4.** Construct the (1) string matching automaton and (2) the state-transition table for the pattern  $P = aabab$  over the alphabet  $\{a, b\}$ .


**1. initially**




**coalesce the two smallest nodes**





**coalesce the next two smallest nodes**




**coalesce the next two smallest nodes**




**coalesce the next two smallest nodes**




**coalesce the next two smallest nodes**



coalesce the next two smallest nodes



The final tree is:



Can you generalize your answer to find the optimal code when the frequencies are the first  $n$  Fibonacci numbers?

**h 0**

**g 10**

**f 110**

**e 1110**

**d 11110**

**c 111110**

**b 1111110**

**a 1111111**


In general, if the letters are  $x_1$  through  $x_n$ , the  $i$ -th Fibonacci number is  $F(i)$ , and  $x_i$  appears with frequency  $F(i)$ , then the code for letter  $x_i$  will consist of  $n - i$  1's followed by one 0, except for the case of  $x_1$ , the least frequent letter, whose code will be  $n - 1$  1's.

2. How many cards are there?  $1 + 2 + \dots + 9 = \frac{9(10)}{2} = 45$

What is their sum?  $1^2 + 2^2 + \dots + 9^2 = \frac{9(9+1)(2 \cdot 9 + 1)}{6} = 285$

To decide which question to ask, you'd build a Huffman tree containing the numbers. Then you'd ask: "Is the chosen less than 9?" If it is, move down to the left subtree and repeat. Otherwise, move to the right subtree and repeat.

1:1    2:2    3:3    4:4    5:5    6:6    7:7    8:8    9:9




3. Determine a **Longest Common Subsequence** (LCS) of  $\langle 1, 0, 0, 1 \rangle$  and  $\langle 0, 1, 0, 1 \rangle$ .

	j	0	1	2	3	4
i		<u>y</u> i	0	1	0	1
0	<u>x</u> i	0	0	0	0	0
1	1	0	↑ 0	↖ 1	← 1	↖ 1
2	0	0	↖ 1	↑ 1	↖ 2	← 2
3	0	0	↖ 1	↑ 1	↖ 2	↑ 2
4	1	0	↑ 1	↖ 2	↑ 2	↖ 3

The LCS is 101, which can be found by starting lower right and using the character corresponding to any cell with a diagonal arrow.

4. Construct the (1) string matching automaton and (2) the state-transition table for the pattern  $P = aabab$  over the alphabet  $\{a, b\}$ .



$\sigma(P_i a) = \text{"length of the longest prefix of } P \text{ which is a suffix of } P_i a \text{"}$

$\delta(\text{state}, \text{input char}) \rightarrow$

ex:  $\delta(\underline{1}, a) = \sigma(P_1 a) = \sigma(aa) = \text{"length of the longest prefix of } P \text{ which is a suffix of 'aa'"} = 2$

State	Input		P
	a	b	
0	$\delta(\underline{0}, a) = \sigma(P_0 a) = \sigma(a) = \underline{1}$	$\delta(\underline{0}, b) = \sigma(P_0 b) = \sigma(b) = \underline{0}$	a
1	$\delta(\underline{1}, a) = \sigma(P_1 a) = \sigma(aa) = \underline{2}$	$\delta(\underline{1}, b) = \sigma(P_1 b) = \sigma(ab) = \underline{0}$	a
2	$\delta(\underline{2}, a) = \sigma(P_2 a) = \sigma(\underline{aaa}) = \underline{2}$	$\delta(\underline{2}, b) = \sigma(P_2 b) = \sigma(\underline{aab}) = \underline{3}$	b
3	$\delta(\underline{3}, a) = \sigma(P_3 a) = \sigma(\underline{aaba}) = \underline{4}$	$\delta(\underline{3}, b) = \sigma(P_3 b) = \sigma(\underline{aabb}) = \underline{0}$	a
4	$\delta(\underline{4}, a) = \sigma(P_4 a) = \sigma(\underline{aabaa}) = \underline{2}$	$\delta(\underline{4}, b) = \sigma(P_4 b) = \sigma(\underline{aabab}) = \underline{5}$	b
5	$\delta(\underline{5}, a) = \sigma(P_5 a) = \sigma(\underline{aababa}) = \underline{1}$	$\delta(\underline{5}, b) = \sigma(P_5 b) = \sigma(\underline{aababb}) = \underline{0}$	

Notes:

$w \sqsubset x \rightarrow w \text{ is a prefix of } x$

$w \sqsupset x \rightarrow w \text{ is a suffix of } x$