


Dealing with data

Customer



1

Active carts



0

Checking out



0

Purchased

✕ Close custom settings

Customer behavior settings

Page views settings

Total Sessions settings

Polling interval: 10000 (ms)



Increment range: 1-2




```
import { useSprings, animated, config } from "react-spring";
```

```
const radiusSprings = useSprings(  
  data.length,  
  data.map((dataPoint, index) => ({  
    radius: radiusScale(dataPoint.count),  
    config: config.molasses,  
  })))  
);
```



Dealing with data restrictions

How can we deal with data restrictions?

How can we deal with data restrictions?

How can we deal with data restrictions?

How can we deal with data restrictions?

How can we deal with data restrictions?

How can we deal with data restrictions?



How can we deal with data restrictions?

How can we deal with data restrictions?

How can we deal with data restrictions?

How can we deal with data restrictions?

How can we deal with data restrictions?

How can we deal with data restrictions?

How can we deal with data restrictions?

How can we deal with data restrictions?



In order to adjust the characteristics of a transition, we have three variables we can tinker with: **mass**, **tension**, and **friction**. Let's look at them each in turn.

Mass

Mass refers to the heft of the thing we're moving. A heavier object will move more slowly, but it also has more inertia.

These two springs are identical, aside from their mass:



Your intuition should come in handy here. Imagine how heavier items would respond when being hung on a spring.


Tension

Tension refers to how tightly-wound the spring is*. The tighter the spring, the more energy is released, leading to a snappy, bouncy animation:



spring-physics

Customer



1

Active carts



0

Checking out



0

Purchased

✕ Close custom settings

Customer behavior settings

Page views settings

Total Sessions settings


Polling interval: 10000 (ms)



Increment range: 1-2



Customer



1

Active carts



0

Checking out



0

Purchased

✕ Close custom settings

Customer behavior settings

Page views settings

Total Sessions settings

Polling interval: 10000 (ms)



Increment range: 1-2



In order to adjust the characteristics of a transition, we have three variables we can tinker with: **mass**, **tension**, and **friction**. Let's look at them each in turn.

Mass

Mass refers to the heft of the thing we're moving. A heavier object will move more slowly, but it also has more inertia.


These two springs are identical, aside from their mass:



Your intuition should come in handy here. Imagine how heavier items would respond when being hung on a spring.

Tension

Tension refers to how tightly-wound the spring is*. The tighter the spring, the more energy is released, leading to a snappy, bouncy animation:




In order to adjust the characteristics of a transition, we have three variables we can tinker with: **mass**, **tension**, and **friction**. Let's look at them each in turn.

Mass

Mass refers to the heft of the thing we're moving. A heavier object will move more slowly, but it also has more inertia.

These two springs are identical, aside from their mass:



Your intuition should come in handy here. Imagine how heavier items would respond when being hung on a spring.

Tension

Tension refers to how tightly-wound the spring is*. The tighter the spring, the more energy is released, leading to a snappy, bouncy animation:



bit.ly/
spring-physics

In order to adjust the characteristics of a transition, we have three variables we can tinker with: **mass**, **tension**, and **friction**. Let's look at them each in turn.

Mass

Mass refers to the heft of the thing we're moving. A heavier object will move more slowly, but it also has more inertia.


These two springs are identical, aside from their mass:



Your intuition should come in handy here. Imagine how heavier items would respond when being hung on a spring.

Tension

Tension refers to how tightly-wound the spring is*. The tighter the spring, the more energy is released, leading to a snappy, bouncy animation:




Animating with react-spring

```
import { useSprings, animated, config } from "react-spring";
```

```
const radiusSprings = useSprings(
  data.length,
  data.map((dataPoint, index) => ({
    radius: radiusScale(dataPoint.count),
    config: config.molasses,
  })))
);
```

```
{data.map((dataPoint, index) => (
  <circle
    cx={xScale(index)}
    r={radiusScale(dataPoint.value)}
    // ...
  />
))}
```



✕ Close custom settings

Customer behavior settings Page views settings Total Sessions settings

Polling interval: 10000 (ms)

Increment range: 1-2