

# Combinatorial Mathematics

Mong-Jen Kao (高孟駿)


Monday 18:30 – 20:20

# Outline

- Hall's Matching Theorem
- König-Egeváry Theorem
- The Maximum Matching Problem
  - A Generic Algorithm and the Berge's Theorem
  - The Augmenting Path Problem in Bipartite Graphs
    - A simple DFS-based recursive algorithm

# Matching in Bipartite Graphs

- Let  $G = (V, E)$  be a bipartite graph with partite sets  $A$  and  $B$ .
- An edge subset  $M \subseteq E$  is called a matching for  $G$ , if each vertex in  $V$  is incident to at most one edge in  $M$ .
  - i.e., the endpoints of the edges in  $M$  are disjoint.





The same definition applies to general graphs, too.

# Matching in Bipartite Graphs

- Let  $G = (V, E)$  be a bipartite graph with partite sets  $A$  and  $B$ .
- Let  $M$  be a matching for  $G$ .
  - For any  $u, v \in V$ ,  
we say that  $u$  is matched to  $v$  by  $M$  (and vice versa),  
if  $(u, v) \in M$ .
  - For any  $U \subseteq A$ , we say that  $M$  matches  $U$ , or,  
 $M$  is a matching from  $U$  to  $B$ , or,  $M$  is a matching for  $U$ ,  
if  $M$  matches every vertex in  $U$  to some vertex in  $B$ .

# Matching in Bipartite Graphs

- Let  $G = (V, E)$  be a bipartite graph with partite sets  $A$  and  $B$ .
- Let  $M$  be a matching for  $G$ .
  - For any  $U \subseteq A$ , we say that  $M$  is a matching for  $U$ , if  $M$  matches every vertex in  $U$  to some vertex in  $B$ .



There is not enough candidates to be matched to for  $A$ .

# Hall's Matching Condition

The necessary and sufficient condition for a matching in bipartite graphs to exist.


### Theorem 5.1 (Hall's Theorem).

Let  $G = (V, E)$  be a bipartite graph with partite sets  $A$  and  $B$ .

There exists a matching  $M$  for  $A$

***if and only if***

$$|N(U)| \geq |U| \quad \text{for all } U \subseteq A. \quad (*)$$



i.e., there is always a sufficient number of candidates to be matched to.

$$|N(U)| \geq |U|, \text{ for any } U \subseteq A.$$

### Theorem 5.1 (Hall's Theorem).

Let  $G = (V, E)$  be a bipartite graph with partite sets  $A$  and  $B$ .

There exists a matching  $M$  for  $A$

***if and only if***

$$|N(U)| \geq |U| \quad \text{for all } U \subseteq A. \quad (*)$$


#### ■ Proof.

– The direction  $(\implies)$  is clear.

■  $M$  matches each vertex in  $U$  to a distinct vertex in  $B$ .

■ Hence,  $|N(U)| \geq |U|$ .





■ Proof. (continue)

- We prove the direction  $(\Leftarrow)$  by induction on the size of  $|A|$ , which we denote by  $m$ .
- The case  $m = 1$  holds trivially.
- Assume that the statement  $(\Leftarrow)$  holds for any  $A$  with  $|A| < m$ .

## ■ Proof. (continue)

- Assume that the statement ( $\Leftarrow$ ) holds when the number of vertices in the left partite set is  $< m$ .
- To prove for  $|A| = m$ , we distinguish the following two cases.

1. For any  $U \subset A$ ,  
we always have  $|N(U)| > |U|$ .

We always have more candidates than we need.

2. For some  $U \subset A$ ,  
 $|N(U)| = |U|$ .

The number of candidates for some subset is tight.

We always have more candidates than we need.

- We distinguish following two cases.

1. For any  $U \subset A$ , we always have  $|N(U)| > |U|$ .

- Pick an arbitrary  $u \in A$  and any  $v \in N(u)$ .  
Match  $u$  to  $v$  and remove  $v$  from the graph.

At most one vertex is removed from  $N(U)$ .

- Then, it follows that,  
for any  $U \subseteq A - \{u\}$ , we still have  $|N(U)| \geq |U|$ .
  - By the induction hypothesis,  
there exists a matching from  $A - \{u\}$  to  $B - \{v\}$ .
  - Hence, we obtain a matching for  $A$ .

- We distinguish following two cases.

2. For some  $U \subset A$ ,  $|N(U)| = |U|$ .

- By the induction hypothesis,  
there exists a matching  $M_1$  from  $U$  to  $N(U)$ .

Remove  $N(U)$  from the graph.

- Then, we claim that,  
for any  $U' \subseteq A - U$ , we still have  $|N(U')| \geq |U'|$ .

The number of candidates  
for some subset is tight.

- We distinguish following two cases.

The number of candidates for some subset is tight.

2. For some  $U \subset A$ ,  $|N(U)| = |U|$ .

- Remove  $N(U)$  from the graph.

- Then, we claim that,  
for any  $U' \subseteq A - U$ , we always have  $|N(U')| \geq |U'|$ .

- If not, then before  $N(U)$  is removed, we have

$$|N(U' \cup U)| \leq |N(U')| + |N(U)| < |U'| + |U|,$$

which is a contradiction.

- We distinguish following two cases.

2. For some  $U \subset A$ ,  $|N(U)| = |U|$ .

- By the induction hypothesis,  
there exists a matching  $M_1$  from  $U$  to  $N(U)$ .

Remove  $N(U)$  from the graph.

- Then, we claim that,  
for any  $U' \subseteq A - U$ , we always have  $|N(U')| \geq |U'|$ .
- By induction hypothesis, there exists a matching  $M_2$  for  $A - U$ .
- Together, we obtain a matching for  $A$ .

The number of candidates  
for some subset is tight.

Application -

System of Distinct Representatives

# Distinct Representative of Sets in a Family

- Let  $F = \{S_1, S_2, \dots, S_m\}$  be a set family.
- The elements  $x_1, x_2, \dots, x_m$  is called a set of distinct representatives for  $F$ , if the following two conditions hold.
  - $x_i \in S_i$  for all  $1 \leq i \leq m$ .
  - The elements  $x_1, x_2, \dots, x_m$  are distinct, i.e.,  $x_i \neq x_j$  for all  $i \neq j$ .



**Corollary.**

The set family  $S_1, S_2, \dots, S_m$  has a set of distinct representatives  
***if and only if***


$$\left| \bigcup_{i \in I} S_i \right| \geq |I| \quad \text{for all } I \subseteq \{1, 2, \dots, m\}.$$

- Construct a bipartite graph for the set family, and this corollary follows directly from the Hall's theorem.

# Matching v.s. Vertex Cover

Weak-duality between matching and vertex cover.

# Vertex Cover of a Graph



- Let  $G = (V, E)$  be a graph.
- A **vertex cover** of  $G$  is a subset  $U \subseteq V$  of vertices such that, every edge  $e \in E$  has at least one endpoint in  $U$ .
  - Intuitively, we use the vertices in  $U$  to cover the edges in  $E$ .



# Matching v.s. Vertex Cover

- Let  $G = (V, E)$  be a graph,
  - $M \subseteq E$  be a matching, and
  - $C \subseteq V$  be a vertex cover for  $G$ .

- It follows that


$$|M| \leq |C|.$$

- The endpoints of the edges in  $M$  are distinct.
- It takes at least one vertex to cover each edge in  $M$ , i.e., at least one endpoint of each edge has to be selected in  $C$ .



The matching  $M$

# Matching v.s. Vertex Cover



The matching  $M$

- Let  $G = (V, E)$  be a graph,  
 $M \subseteq E$  be a matching, and  $C \subseteq V$  be a vertex cover for  $G$ .
- Then, it follows that  $|M| \leq |C|$ .
  - This property is called the weak-duality between the matching and vertex cover.
  - It implies that, in any graph, the size of maximum matching is at most the size of minimum vertex cover.

# The König-Egeváry Theorem


**In bipartite graphs**, the size of the ***maximum matching*** is equal to the size of the ***minimum vertex cover***.

### Theorem 5.5 (König-Egeváry 1931).

In a bipartite graph, the size of ***maximum matching*** is equal to the size of ***minimum vertex cover***.

#### Proof.

- Let  $G$  be a bipartite graph with partite sets  $U$  and  $V$ .
- Let  $M$  be a *maximum matching* and  $C$  be a *minimum vertex cover* for  $G$ , respectively.
- It suffices to prove that  $|M| \geq |C|$ .




The matching  $M$

**Theorem 5.5 (König-Egeváry 1931).**

In a bipartite graph, the size of ***maximum matching*** is equal to the size of ***minimum vertex cover***.

**Proof.**


- It suffices to prove that  $|M| \geq |C|$ .
  - Let  $A := U \cap C$  and  $B := V \cap C$ .
  - We will prove that, there exists a matching  $M_A$  that matches all the vertices in  $A$  to the vertices in  $V \setminus B$ .





## Proof.

- It suffices to prove that  $|M| \geq |C|$ .
- Let  $A := U \cap C$  and  $B := V \cap C$ .
  - We will prove that, there exists a matching  $M_A$  that matches all the vertices in  $A$  to the vertices in  $V \setminus B$ .
  - **If the above is true**, then by a similar argument, there exists a matching  $M_B$  for  $B$  to  $U \setminus A$ .
  - The endpoints of the edges in  $M_A \cup M_B$  are distinct.
    - So,  $M_A \cup M_B$  is a matching of size  $|A| + |B| = |C|$ .
    - Hence, this will prove that  $|M| \geq |A| + |B| = |C|$ .




It suffices to prove that, there exists a matching  $M_A$  that matches all the vertices in  $A$  to the vertices in  $V \setminus B$ .

- Suppose that there exists no such matching.
- Then, by Hall's matching theorem, there exists some  $D \subseteq A$ , such that


$$|N(D) \cap (V \setminus B)| < |D|.$$

- Indeed, if  $|N(D) \cap (V \setminus B)| \geq |D|$  holds for all  $D \subseteq A$ , then there exists a matching from  $A$  to  $V \setminus B$ .
- Since there is no such matching, there must be such a  $D \subseteq A$  with  $|N(D) \cap (V \setminus B)| < |D|$ .



It suffices to prove that, there exists a matching  $M_A$  that matches all the vertices in  $A$  to the vertices in  $V \setminus B$ .

- If not, there exists some  $D \subseteq A$ , such that
 
$$|N(D) \cap (V \setminus B)| < |D|.$$
- Let  $\tilde{D} := N(D) \cap (V \setminus B)$ , then  $|\tilde{D}| < |D|$ .
- We claim that,
  - $(A \setminus D) \cup \tilde{D} \cup B$  is a valid vertex cover for  $G$ .
  - If this is true, we obtain a vertex cover with size smaller than  $|A| + |B| = |C|$ , a contradiction.




$D$  is replaceable by  $\tilde{D}$ .

It suffices to verify that,

$(A \setminus D) \cup \tilde{D} \cup B$  is a valid vertex cover for  $G$ .

- Let  $\tilde{D} := N(D) \cap (V \setminus B)$ .
- There are four categories of edges in  $G$ .
  - $E_{A,B}, E_{U \setminus A, B}$  --- covered by  $B$ .
  - $E_{A \setminus D, V \setminus B}$  --- covered by  $A \setminus D$ .
  - $E_{D, \tilde{D}}$  --- covered by  $\tilde{D}$ .
- All the edges are covered.



Since  $C = A \cup B$  is a vertex cover, there is not edge between  $U \setminus A$  and  $V \setminus B$ .

# The Maximum Matching Problem

To compute a maximum-size matching for the input graph.

# The Maximum Matching Problem

- Input :


- A graph  $G = (V, E)$ .

- Output :

- A matching  $M \subseteq E$  that has the maximum size among all possible matchings.

# Alternating Path & Augmenting Path


- Let  $M$  be a matching for a graph  $G$ .
  - An  $M$ -alternating path is a path that alternates between edges in  $M$  and edges not in  $M$ .
  - An  **$M$ -augmenting path** is an  $M$ -alternating path that both starts and ends at unmatched vertices.



Both  $v_1, v_2, v_3$  and  $v_2, v_3, v_4, v_5$  are  $M$ -alternating paths.




$v_1, v_2, v_3, v_4, v_5, v_6$  is an  $M$ -augmenting path.

# Observation



- We can see that,  
each  $M$ -augmenting path is a way to enlarge the size of  $M$  by 1.
  - This is done by swapping the status of the edges on the path.
    - Matched edges  $\Rightarrow$  *unmatched*
    - Unmatched edges  $\Rightarrow$  *matched*

So, this is still a valid matching  
with size increased by 1.




$v_1$  and  $v_6$  were unmatched.

All internal vertices are matched only by edges on the path.



# Observation



- We can see that,  
each  $M$ -augmenting path  $P$  is a way to enlarge the size of  $M$  by 1.
- $M' := (M \setminus P) \cup (P \setminus M)$  is a valid matching with  $|M'| = |M| + 1$ .

$M \Delta P$  : the edges that appear exactly once in  $M$  and  $P$ .

# A Simple Greedy Algorithm

- The observation suggests the following greedy algorithm.
  - Let  $G = (V, E)$  be the input graph.

1.  $M \leftarrow \emptyset$ .
2. Repeat until there is no  $M$ -augmenting path in  $G$ .
  - a. Compute an  $M$ -augmenting path  $P$ .
  - b. Set  $M \leftarrow (M \setminus P) \cup (P \setminus M)$ .
3. Output  $M$ .

1.  $M \leftarrow \emptyset$ .
2. Repeat until there is no  $M$ -augmenting path in  $G$ .
  - a. Find an  $M$ -augmenting path  $P$ .
  - b. Set  $M \leftarrow (M \setminus P) \cup (P \setminus M)$ .
3. Output  $M$ .

The philosophy behind the algorithm is very simple :

“Make the current matching larger until no augmenting path exists.”

■ A direct question is that,

**“Does it always output a maximum matching?”**

**Theorem 1. (Berge 1957).**

A matching  $M$  in a graph  $G$  is a maximum matching if and only if  $G$  has no  $M$ -augmenting path.

- Theorem 1 assures the correctness of the greedy algorithm.
  - When there is no  $M$ -augmenting path,  $M$  is guaranteed to be maximum.
- We begin with some definition & helper lemma.

# Symmetric Difference

- Let  $G = (V, E)$  be a graph, and  $A, B \subseteq E$  be two edge sets.

- The symmetric difference of  $A$  and  $B$  is defined as




$$A \triangle B := (A \setminus B) \cup (B \setminus A) .$$

- i.e., the set of edges that appear exactly once in  $A$  and  $B$ .

## Lemma 2.

Let  $M, M'$  be matchings for a graph  $G$ . Then, every component of  $M \triangle M'$  is either a *path* or a *cycle with an even length*.

- Let  $F := M \triangle M'$ .
  - Each vertex in  $G$  is incident to at most two edges in  $F$ .
  - Hence, each component in  $F$  is either a path or a cycle.
- Consider any cycle in  $F$ .
  - The cycle alternates between edges in  $M$  and  $M'$ .
  - It must have an even length.



### Theorem 1. (Berge 1957).

A matching  $M$  in a graph  $G$  is a maximum matching if and only if  $G$  has no  $M$ -augmenting path.

- Let us prove Theorem 1.
  - The direction ( $\Rightarrow$ ) is clear.
  - For the direction ( $\Leftarrow$ ),  
we prove the contrapositive statement.
    - We show that, if  $M'$  is a matching with  $|M'| > |M|$ ,  
then  $G$  must have an  $M$ -augmenting path.

It suffices to prove that, if  $M'$  is a matching with  $|M'| > |M|$ , then  $G$  must have an  $M$ -augmenting path.

- Let  $F := M \triangle M'$ .
  - By Lemma 2,  $F$  is a union of paths and even cycles.
- Since  $|M'| > |M|$ , there must be a component in  $F$  that has more edges from  $M'$  than  $M$ .
  - The component must be a path.  
Furthermore, it must start and ends with edges in  $M'$ .
  - The path is then an  $M$ -augmenting path.