

Clinical Study Data Reviewer's Guide

Takeda Pharmaceutical Company Limited.

Study TAK003-DEN-324

cSDRG Template Version 2018-11-01

Clinical Study Data Reviewer's Guide

Contents

1.	Introduction.....	4
1.1	Purpose	4
1.2	Acronyms	4
1.3	Study Data Standards and Dictionary Inventory	4
2.	Protocol Description	5
2.1	Protocol Number and Title	5
2.2	Protocol Design	5
2.3	Trial Design Datasets.....	6
2.3.1	TA - Trial Arms	6
2.3.2	TE - Trial Elements.....	7
2.3.3	TI - Trial Inclusion/Exclusion Criteria	7
2.3.4	TS - Trial Summary Information	7
2.3.5	TV - Trial Visits.....	7
3.	Subject Data Description	8
3.1	Overview	8
3.2	Traceability Flow Diagram.....	9
3.3	Annotated CRFs	9
3.4	SDTM Subject Domains.....	12
3.4.1	AE - Adverse Events.....	12
3.4.2	CE - Clinical Events.....	13
3.4.3	CM - Concomitant/Prior Medications	13
3.4.4	DD - Death Details	14
3.4.5	DM - Demographics	14
3.4.6	DS - Disposition.....	15
3.4.7	DV - Protocol Deviations.....	15
3.4.8	EC - Exposure as Collected	15
3.4.9	FA - Findings About Events or Interventions	15
3.4.10	IS - Immunogenicity Specimen Assessments	16
3.4.11	MH - Medical History.....	16
3.4.12	VS - Vital Signs	16
4.	Data Conformance Summary.....	17
4.1	Conformance Inputs	17

4.2	Issues Summary	17
4.3	Additional Conformance Details	20
Appendix I: Inclusion/Exclusion Criteria		20

1. Introduction

1.1 Purpose

This document provides context for tabulation datasets and terminology that benefit from additional explanation beyond the Data Definitions document (define.xml). In addition, this document provides a summary of SDTM conformance findings.

1.2 Acronyms

Acronym	Translation
IB	Investigator's Brochure
IMP	Investigational Medicinal Product
IRT	Interactive Response Technology
JE	Japanese Encephalitis
LAR	Legally Acceptable Representative
M	Month
MAAE	Medically Attended Adverse Event
MNT	Microneutralization Test
TDV	Takeda's Dengue Tetravalent Vaccine (Live, Attenuated)
V	Visit
Y	Years

1.3 Study Data Standards and Dictionary Inventory

Standard or Dictionary	Versions Used
SDTM	SDTM v1.7/SDTMIG v3.3
Controlled Terminology	•CDISC SDTM Controlled Terminology, 2024-03-29 •Sponsor-defined terms appended to CDISC SDTM Terminology 2024-03-29
Data Definitions	Define-XML v2.1
Medications Dictionary	WHODD GLOBALB3Sep25
Medical Events Dictionary	MedDRA v28.1
Other standards (optional)	SNOMED 2024-09-01, UNII 2024-09-27, MED-RT 2023-09-05

2. Protocol Description

2.1 Protocol Number and Title

Protocol Number: TAK003-DEN-324

Protocol Title: A Randomized, Double-Blind, Placebo-Controlled (Participants aged 18 to 60 years) and Open-Label (Participants aged 4 to 17 years), Phase 2/3 Trial to Evaluate the Immunogenicity and Safety of 2 doses of a Dengue Tetravalent Vaccine (Live, Attenuated) (TDV) Administered Subcutaneously to Healthy Adults, Adolescents, and Children in Japan

Protocol Versions: Original Protocol (Version 1.0): 26 Sep 2024
Amendment 1 (Version 2.0): 28 Nov 2024

Protocol Amendment 1 (Version 2.0, 28-Nov-2024) updated eligibility criteria through changes to Exclusion Criteria (Section 7.1.2), adding contraception/sperm donation requirements for non-sterilized male participants. It also added reporting and follow-up requirements for pregnancy, including pregnancy in partners of male participants (Sections 9.1.9/9.1.10). These changes affected safety/eligibility-related data collection, but did not change endpoints, statistical methods, or sample size; therefore, no material impact on efficacy/immunogenicity data interpretation is expected.

2.2 Protocol Design

This is a phase 2/3, randomized, double-blind, placebo-controlled trial in 176 healthy participants aged 18 to 60 years and open-label trial in 10 healthy participants aged 4 to 17 years conducted in Japan to investigate the immunogenicity and safety of TDV administered as 2 doses given 3 months apart (on Day 1 [Month 0] and Day 90 [Month 3], respectively) via subcutaneous injection. Participants aged 18 to 60 years will be randomized (3:1 ratio) to TDV (n=132) or placebo (n=44). Participants aged 4 to 17 years will receive TDV (n=10).

Cohort 1: Participants aged 18 to 60 years (TDV [Group 1], n=132; Placebo [Group 2], n=44).


Cohort 2: Participants aged 4 to 17 years (TDV [Group 3], n=10).

Cohort 1 and 2 will be enrolled in parallel.

All participants will be followed-up for 6 months post second investigational medicinal product administration on Day 90 (M3) until Day 270 (Month 9), so the trial duration will be approximately 270 days (9 months) for each participant.

Immunogenicity evaluation: Neutralizing antibodies against the 4 dengue serotypes (MNT) were assessed using blood samples collected at pre-dose 1, 1 month post-dose 1, pre-dose 2, 1 month post-dose 2, and 6 months post-dose 2.

Safety evaluation: Safety was evaluated by collection of solicited local AEs (7 days post-dose) and solicited systemic AEs (14 days post-dose) after each IMP administration (Day 1 and Day 90), using participant diary records. Unsolicited AEs were collected for 28 days after each dose and recorded in eCRF. SAEs, medically attended AEs (MAAEs), and AEs leading to participant discontinuation were collected from post-first dose through end of trial. AEs leading to IMP discontinuation were collected from post-first dose until second IMP administration (Day 90).

Figure 1.a Study Schema

2.3 Trial Design Datasets

Are Trial Design datasets included in the submission?

Yes.

Trial Design datasets used in this study are as shown in below table.

Dataset	Dataset Label
TA	Trial Arms
TE	Trial Elements
TI	Trial Inclusion/Exclusion Criteria
TS	Trial Summary Information
TV	Trial Visits

2.3.1 TA - Trial Arms

TA is provided to describe planned trial arms across two cohorts enrolled in parallel with different blinding schemes. ARMCD values are cohort-specific: C1TAK003 (Cohort 1: TAK-003), C1PLACEBO (Cohort 1: Placebo), and C2TAK003 (Cohort 2: TAK-003). Cohort 1 (participants aged 18–60 years) was randomized 3:1 to TAK-003 or placebo and conducted as double-blind; Cohort 2 (participants aged 4–17 years) was open-label and received TAK-003 only. No crossover or open-label extension period was implemented.

2.3.2 TE - Trial Elements

TE defines four planned elements: SCRN (Screening), C1TAK (TAK-003 Treatment for Cohort 1), C1PLA (Placebo for Cohort 1), and C2TAK (TAK-003 Treatment for Cohort 2).

Screening begins at informed consent and continues until first-dose eligibility is confirmed. Treatment elements are cohort/arm-specific and reflect the planned 2-dose regimen (Day 1 and Day 90), with planned element duration represented as 270 days.

2.3.3 TI - Trial Inclusion/Exclusion Criteria

Exclusion criteria are represented with IECAT (INCLUSION/EXCLUSION), and protocol version is identified using TIVERS.

The trial inclusion/exclusion criteria are not fully described in the TI domain. Full text of the criteria can be found in [Appendix I: Inclusion/Exclusion Criteria](#).

2.3.4 TS - Trial Summary Information

Cohort-specific TS records are represented using TSGRPID (C1 and C2) for parameters such as INTMODEL, TBLIND, TCNTRL, and RANDQT. Specifically, Cohort 1 is parallel, randomized, double-blind, placebo-controlled; Cohort 2 is single-group, open-label, with no control.

Two data cutoff records are included in TS (DCUTDESC/DCUTDTC): DATABASE LOCK (2026-04-07) and KEY OPEN (2026-06-10), reflecting staged data availability associated with key release and integration of final analysis data. KEY OPEN corresponds to availability of unblinded treatment assignment information and immunogenicity data for final analysis.

TSVAL1 is used to store continuation text when a parameter value exceeds the length limit of TSVAL (e.g., long text values).

2.3.5 TV - Trial Visits

TV includes planned visits Visit 1, Visit 2, Visit 3, Visit 4, Visit 5, and Early Termination. Visit 1 also covers screening activities (including informed consent) prior to first dose. Early Termination is represented as a separate planned visit for discontinuation assessments.

Planned visit timing is aligned to first IMP administration: Day 1 (M0), Day 30 (M1), Day 90 (M3), Day 120 (M4), and Day 270 (M9).

3. Subject Data Description

3.1 Overview

Are the submitted data taken from an ongoing study?

No.

Were the SDTM datasets used as sources for the analysis datasets?

Yes.

Do the submission datasets include screen failures?

Yes.

If yes, which datasets include screen failure data?

Dataset	Dataset Label
DM	Demographics
DS	Disposition
IE	Inclusion/Exclusion Criteria Not Met
SE	Subject Elements
SUPPDM	Supplemental Qualifiers for DM
SV	Subject Visits

Were any domains planned, but not submitted because no data were collected?

No.

Are the submitted data a subset of collected data?

No.

Is adjudication data present?


No.

Additional Content of Interest

Baseline was represented using the Last Observation Before Exposure Flag (--LOBXFL) rather than the Baseline Flag (--BLFL) in all Findings domains.

Reference start date (RFSTDTC) and Date/Time of First Study Treatment (RFXSTDTC) in DM domain are both set to the date of first evaluable dose.

3.2 Traceability Flow Diagram



3.3 Annotated CRFs

Collected fields and pages that have not been tabulated have been annotated as "Not Submitted". Takeda Pharmaceutical Company Limited. collects certain data elements to facilitate operational processes including data cleaning and dynamically creating additional forms in the electronic data capture system. All fields and pages that have been annotated as "Not Submitted" meet this criterion.

Explanation of data fields [Not Submitted]

aCRF page Number(s)	Data Collection Field	Explanation of why [NOT SUBMITTED]
3	Form: AE/SAE Specify Disability/Incapacity	This collected field was pre-specified as out of SDTM tabulation scope in the study mapping design.
4, 5	Form: AE/SAE Narrative, Narrative 2, Narrative 3, Narrative 4	This collected field was pre-specified as out of SDTM tabulation scope in the study mapping design.
6	Form: ANY ADVERSE EVENTS?	This field is used for operational purpose only.
7	Form: ANY CONCOMITANT MEDICATIONS?	This field is used for operational purpose only.
8	Form: ANY MEDICAL HISTORY?	This field is used for operational purpose only.
9	Form: ANY PRIOR INTERVENTIONS?	This field is used for operational purpose only.
10	Form: ANY PRIOR/CONCOMITANT LICENSED VACCINE ADMINISTRATION?	This field is used for operational purpose only.
11	Form: CENTRAL LAB Check here if Sample Collection was Done/Not Done	This field is used for operational purpose only.

aCRF page Number(s)	Data Collection Field	Explanation of why [NOT SUBMITTED]
18	Form: DATE OF VISIT ADHOC Type of Visit	This field is used for operational purpose only.
88	Form: DOSING INJECTION Dose Form	These are system-calculated (protocol specified) fields in EDC and are not source-collected CRF data fields for SDTM tabulation.
88	Form: DOSING INJECTION Route of Administration	These are system-calculated (protocol specified) fields in EDC and are not source-collected CRF data fields for SDTM tabulation.
88	Form: DOSING INJECTION Location of Dose Administration	These are system-calculated (protocol specified) fields in EDC and are not source-collected CRF data fields for SDTM tabulation.
90	Form: END OF STUDY Did the subject prematurely discontinue from the study?	This field is used for operational purpose only.
91	Form: END OF STUDY TREATMENT Did the subject prematurely discontinue from study treatment?	This field is used for operational purpose only.
91	Form: END OF STUDY TREATMENT Has the subject's treatment been unblinded?	This field is used for operational purpose only.
92	Form: INCLUSION/EXCLUSION CRITERIA Did the subject meet all of the eligibility criteria?	This field is used for operational purpose only.
101	Form: MEDICATION HISTORY Related Medical History (Primary)	This collected field was pre-specified as out of SDTM tabulation scope in the study mapping design.
101	Form: MEDICATION HISTORY Related Medical History (#2, if applicable)	This collected field was pre-specified as out of SDTM tabulation scope in the study mapping design.
101	Form: MEDICATION HISTORY Related Medical History (#3, if applicable)	This collected field was pre-specified as out of SDTM tabulation scope in the study mapping design.

aCRF page Number(s)	Data Collection Field	Explanation of why [NOT SUBMITTED]
102	Form: POST VACCINATION GENERAL SYMPTOMS (6 YEARS OLD OR MORE) Was assessment performed at the planned time point?	This field is used for operational purpose only.
105	Form: POST VACCINATION GENERAL SYMPTOMS (LESS THAN 6 YEARS OLD) Was assessment performed at the planned time point?	This field is used for operational purpose only.
108	Form: POST VACCINATION REACTIONS AT INJECTION SITE (6 YEARS OLD OR MORE) Was assessment performed at the planned time point?	This field is used for operational purpose only.
109	Form: POST VACCINATION REACTIONS AT INJECTION SITE (LESS THAN 6 YEARS OLD) Was assessment performed at the planned time point?	This field is used for operational purpose only.
110	Form: PREGNANCY TEST Check if the Pregnancy Test was Done/Not Done	This field is used for operational purpose only.
118	Form: SCREENING DISPOSITION Did the subject experience any serious pretreatment events?	This field is used for operational purpose only.
119	Form: VITAL SIGNS Indicate if Vital Signs Assessment was Performed	This field is used for operational purpose only.
120	Form: VITAL SIGNS ONLY Indicate if Vital Signs Assessment was Performed	This field is used for operational purpose only.
121	Form: VITAL SIGNS_DAY 90 Indicate if Vital Signs Assessment was Performed	This field is used for operational purpose only.

3.4 SDTM Subject Domains

Dataset - Dataset Label	Efficacy	Safety	Other	Custom	SUPP--	Related Using RELREC
AE - Adverse Events		X			X	CM
CE - Clinical Events		X			X	FA, VS
CM - Concomitant/Prior Medications			X		X	AE, MH
DD - Death Details		X				
DM - Demographics			X		X	
DS - Disposition			X			
DV - Protocol Deviations			X		X	
EC - Exposure as Collected			X		X	EX
EX - Exposure			X			EC
FA - Findings About Events or Interventions		X			X	CE
IE - Inclusion/Exclusion Criteria Not Met			X			
IS - Immunogenicity Specimen Assessments	X				X	
LB - Laboratory Test Results		X				
MH - Medical History			X		X	CM
RP - Reproductive System Findings			X			
SE - Subject Elements			X			
SV - Subject Visits			X			
VS - Vital Signs		X			X	CE

3.4.1 AE - Adverse Events

A relationship has been defined between concomitant/prior medications and adverse events.

The following variables are stored in SUPPAE. Following QNAM = AEDISCDT, AEHOSPDT, ACNDNC, ACNDW, ACNNA, ACNDD were planned, but not submitted because no data was collected.

QNAM	Description
ACNDD	Dose Delayed
ACNDNC	Dose Not Changed
ACNDW	Drug Withdrawn
ACNNA	Not Applicable
AEDIS	Caused Study Discontinuation
AEDISCDT	Date of Discharge
AEHOSPD	Date of Hospitalization
AEPTRTPT	Did This Event Occur Prior to The First
AERELPR	Relationship to Procedure
AESLCTD	Is This a Solicited Adverse Event?
AEVCDVER	Version of The Coding Dictionary
MAAE	Is This a Medically Attended AE?

3.4.2 CE - Clinical Events

A relationship has been defined between clinical events and findings about events or interventions.

A relationship has been defined between clinical events and vital signs.

The following variables are stored in SUPPCE:

QNAM	Description
CEEVAL	Rater ID
CESSCAT	CE 2nd Sub-Category
RELPR	Relationship to Study Procedure

3.4.3 CM - Concomitant/Prior Medications

A relationship has been defined between concomitant/prior medications and adverse events.

A relationship has been defined between concomitant/prior medications and medical history.

The following variables are stored in SUPPCM:

QNAM	Description
ATCCODE1	WHO Anatomical Main Group Code

QNAM	Description
ATCCODE2	WHO Therapeutic Subgroup Code
ATCCODE3	WHO Pharmacological Subgroup Code
ATCCODE4	WHO Chemical Subgroup Code
ATCTEXT1	WHO Anatomical Main Group
ATCTEXT2	WHO Therapeutic Subgroup
ATCTEXT3	WHO Pharmacological Subgroup
ATCTEXT4	WHO Chemical Subgroup
CMDECOD1	Standardized Medication Name 1
CMVCDVER	Version of the Coding Dictionary
TRTMAN	Treatment Manufacturer
VACDOC	Supported by Vaccination Card/Oth Docmnt
VACHISTR	Source of Vaccination History

3.4.4 DD - Death Details

DDSPID is used to distinguish multiple General Cause of Death entries where DDTESTCD = GENCDTH. DDDTC is set to null because the CRF did not collect the collection date of death detail.

3.4.5 DM - Demographics

At the unblinding stage, treatment assignment information was made available from external IRT data. ARMCD in DM was derived using cohort information and the external treatment assignment.

The following variables are stored in SUPPDM:

QNAM	Description
AGERAND	Age at Randomization
COHORT	Cohort
PLGCNTO	Parent/Legal Guardian Consent Obtained
PLGCNTO2	Second Parent/Legal Guardian Consent
PROTVRSN	Protocol Version
RACESASN	Race Asian Sub-Category
SPASENTO	Study Participation Assent Obtained

3.4.6 DS - Disposition

Following QNAM = PROTVRSN has been planned, but not submitted because no data was collected.

QNAM	Description
PROTVRSN	Protocol Version

3.4.7 DV - Protocol Deviations

The following variables are stored in SUPPDV:

QNAM	Description
DVCLASI	Classification of Protocol Deviation
DVTERM1	Protocol Deviation Term1

3.4.8 EC - Exposure as Collected

A relationship has been defined between the exposure as collected and the exposure. The observations are related by ECLNKID and EXLNKID.

The following variables are stored in SUPPEC. Following QNAM = ECACN, ECDEL, ECDELOTH were planned, but not submitted because no data was collected.

QNAM	Description
ECACN	Action Taken with Study Treatment
ECDEL	Reason for Dose Delayed
ECDELOTH	Dose Delayed Other Specify
ECREASOC	Reason for Occur Value

3.4.9 FA - Findings About Events or Interventions

A relationship has been defined between findings about events or interventions and clinical events.

FA is used to capture reactogenicity findings about clinical events, derived from post-vaccination assessment forms and diary forms, with age-stratified forms for participants ≥ 6 years and < 6 years of age.

The following variables are stored in SUPPFA.

QNAM	Description
FAREASOC	Reason Medication Taken
FASSCAT	FA 2nd Sub-Category

3.4.10 IS - Immunogenicity Specimen Assessments

The following variables are stored in SUPPIS:

QNAM	Description
ISLLOD	Lower Limit of Detection
ISTSTOPO	Test Operational Objective
ISULOD	Upper Limit of Detection

3.4.11 MH - Medical History

A relationship has been defined between medical history and concomitant/prior medications.

The following variable is stored in SUPPMH.

QNAM	Description
MHVCDVER	Version of the Coding Dictionary

3.4.12 VS - Vital Signs

A relationship has been defined between vital signs and clinical events.

The following variables are stored in SUPPVS.

QNAM	Description
VSCOLSRT	Collected Summary Result Type
VSSCAT	VS 2nd Sub-Category

4. Data Conformance Summary

4.1 Conformance Inputs

Was a validator used to evaluate conformance?

Yes

If yes, specify the version(s) of the validation rules:

Pinnacle 21 Enterprise version 6.1.1

Validation Engine version PMDA 2411.1, PMDA Validation Rule: v6.0

Were sponsor-defined validation rules used to evaluate conformance?

No

Were the SDTM datasets evaluated in relation to define.xml?

Yes

Was define.xml evaluated?

Yes

Provide any additional compliance evaluation information:

No

4.2 Issues Summary

Check ID	Diagnostic Message	Severity	Dataset	Count (Issue Rate)	Explanation
SD0058	Variable appears in dataset, but is not in SDTM model	Error	AE	1 (2.50%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	CE	1 (4.76%)	The SUBJID was added to all domains to uniquely identify subjects per screening.

Check ID	Diagnostic Message	Severity	Dataset	Count (Issue Rate)	Explanation
SD0058	Variable appears in dataset, but is not in SDTM model	Error	CM	1 (3.57%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	DD	1 (7.69%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	DS	1 (6.67%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	DV	1 (5.56%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	EC	1 (3.70%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	EX	1 (4.35%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	FA	1 (3.13%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	IE	1 (6.25%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	IS	1 (2.86%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	LB	1 (3.45%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	MH	1 (3.57%)	The SUBJID was added to all domains to uniquely identify subjects per screening.

Check ID	Diagnostic Message	Severity	Dataset	Count (Issue Rate)	Explanation
SD0058	Variable appears in dataset, but is not in SDTM model	Error	RELREC	1 (12.50%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	RP	1 (5.56%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	SE	1 (7.69%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	SV	1 (7.14%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD0058	Variable appears in dataset, but is not in SDTM model	Error	VS	1 (3.45%)	The SUBJID was added to all domains to uniquely identify subjects per screening.
SD1317	DSSTDTC for DEATH record is not the same as DM.DTHDTC	Error	DS	1 (100.00%)	DSSTDTC is the collected date of end of study due to death.
SD2244	Invalid TSVCDREF value for FCNTRY	Error	TS	1 (100.00%)	For TSPARMCD=FCNTRY [Planned Country of Investigational Sites], the appropriate code reference for TSVAL="JPN" would, strictly speaking, be more clearly specified as "ISO 3166-1 alpha-3" rather than "ISO 3166". However, "ISO 3166-1 alpha-3" is not included in CL.C66788.DICTNAM in either our internal Takeda SDTM CT or the latest SDTM CT (2026-03-27). Therefore, in accordance with these CTs, we will keep TSVCDREF as "ISO 3166" and will not change it.

Check ID	Diagnostic Message	Severity	Dataset	Count (Issue Rate)	Explanation
DD0132	Invalid use of def:ExtendedValue attribute	Error	DEFINE	1 (100.00%)	The value "DOSE DELAYED" was added in accordance with the company standards.
DD0137	Invalid variable label value	Error	DEFINE	3 (100.00%)	The labels were assigned according to company standards.

4.3 Additional Conformance Details

There are no additional details to be documented.

Appendix I: Inclusion/Exclusion Criteria

Protocol/ Amendment Version	Category	IETESTCD	Full Text of Criterion
ORIGINAL	EXCLUSION	E01V010	Participant has contraindication(s), warning(s) and/or precaution(s) applicable to vaccination with TDV as specified in the IB.
ORIGINAL	EXCLUSION	E02V010	Participant has a known hypersensitivity or allergy to any of the IMP components (including excipients of the IMP).
ORIGINAL	EXCLUSION	E03V010	Participant has behavioral or cognitive impairment or psychiatric disease that, in the opinion of the investigator, may interfere with the participant's ability to participate in the trial.
ORIGINAL	EXCLUSION	E04V010	Participant has a history of progressive or severe neurologic disorder, seizure disorder or neuro-inflammatory disease (eg, Guillain-Barré syndrome).

Protocol/ Amendment Version	Category	IETESTCD	Full Text of Criterion
ORIGINAL	EXCLUSION	E05V010	Participant has a clinically significant active infection (as assessed by the investigator) or body temperature $>38.0^{\circ}\text{C}$ ($>100.4^{\circ}\text{F}$) within 3 days of intended IMP administration on Day 1 (M0). Note: In principle, oral temperature should be measured for body temperature. In cases where it is difficult to measure oral temperature, such as in young children, underarm (axillary) temperature may be used instead.
ORIGINAL	EXCLUSION	E06V010	Participant has an illness, or history of any illness that, in the opinion of the investigator, might interfere with the results of the trial or pose additional risk to the participant due to involvement in this trial.
ORIGINAL	EXCLUSION	E07V010	Participant has a known or suspected impairment/alteration of immune function, including: a. Chronic administration of oral and/or parenteral steroids at doses considered sufficiently immunosuppressive (eg, ≥ 2 mg/kg body weight/day prednisone/prednisolone [or equivalent] for ≥ 14 consecutive days, or, ≥ 20 mg/day prednisone/prednisolone [or equivalent] for ≥ 14 consecutive days) within 60 days prior to Day 1 (M0). Note: Use of corticosteroids by inhaled, intranasal, intraarticular, bursal, tendon injection, or topical routes is allowed. b. Receipt of blood, immunoglobulins, blood products, and/or plasma derivatives within the 90 days prior to Day 1 (M0). c. Receipt of immunostimulants within 60 days prior to Day 1 (M0). d. Immunosuppressive therapy such as anti-cancer chemotherapy or radiation therapy within 6 months prior to Day 1 (M0). e. Reported or known symptomatic HIV infection or asymptomatic HIV infection when accompanied by evidence of impaired immune function. f. Reported or known Hepatitis B and/or Hepatitis C virus infection. g. Genetic immunodeficiency.
ORIGINAL	EXCLUSION	E08V010	Participant has known or suspected abnormalities of splenic or thymic function.
ORIGINAL	EXCLUSION	E09V010	Participant has a known bleeding diathesis, or any condition that may be associated with a prolonged bleeding time.
ORIGINAL	EXCLUSION	E10V010	Participant has a serious chronic or progressive disease deemed to be preclusive to trial entry, ie, not medically stable according to the judgment of the investigator.

Protocol/ Amendment Version	Category	IETESTCD	Full Text of Criterion
ORIGINAL	EXCLUSION	E11V010	Participant is participating in any clinical trial with another investigational product within 30 days prior to Day 1 (M0) or plans to participate in another clinical trial at any time during the conduct of this trial.
ORIGINAL	EXCLUSION	E12V010	Participant has previously received a vaccination against flavivirus other than Japanese encephalitis (JE) (investigational or licensed).
ORIGINAL	EXCLUSION	E13V010	Participant who received any other vaccines within 14 days (for inactivated vaccines) or 28 days (for live vaccines) prior to Day 1 (M0) or who are planning to receive any vaccine other than IMP within 28 days of IMP administration.
ORIGINAL	EXCLUSION	E14V010	Participant who received a coronavirus vaccine within 14 days prior to Day 1 (M0).
ORIGINAL	EXCLUSION	E15V010	Participant who received a vaccine authorized for emergency use within 28 days prior to Day 1 (M0).
ORIGINAL	EXCLUSION	E16V010	Participant who received any JE vaccines within 28 days prior to Day 1 (M0) or who are planning to receive any JE vaccines during the trial period.
ORIGINAL	EXCLUSION	E17V010	Previous participation in any clinical trial of a dengue or other flavivirus candidate vaccine, except for participants who received placebo in those trials.
ORIGINAL	EXCLUSION	E18V010	Participant with body mass index (BMI) greater than or equal to 35 kg/m ² (=weight in kg/[height in meters] ²) on Day 1 (M0).
ORIGINAL	EXCLUSION	E19V010	Participant who intends to travel to dengue endemic areas during the trial period.
ORIGINAL	EXCLUSION	E20V010	Participant with documented or suspected disease caused by a flavivirus and participants with a history of prolonged (≥ 1 year) habitation in a dengue endemic area.
ORIGINAL	EXCLUSION	E21V010	Participant with history of substance or alcohol abuse within the past 2 years prior to Day 1 (M0).
ORIGINAL	EXCLUSION	E22V010	Female participants who are pregnant (ie, a positive or indeterminate pregnancy test) or breastfeeding.

Protocol/ Amendment Version	Category	IETESTCD	Full Text of Criterion
ORIGINAL	EXCLUSION	E23V010	Females of childbearing potential who are sexually active and who have not used any of the acceptable contraceptive methods for at least 2 months prior to Day 1 (M0). a. "Childbearing potential" is defined as status post onset of menarche and not meeting any of the following conditions: menopausal for at least 2 years, status after bilateral tubal ligation for at least 1 year, status after bilateral oophorectomy, or status after hysterectomy. b. "Acceptable contraceptive methods" are defined as one or more of the following: · Hormonal contraceptive. · Barrier method (condom or diaphragm) every time during intercourse. · Intrauterine device. · Monogamous relationship with a vasectomized partner. The partner must have been vasectomized for at least 6 months prior to the participant's trial enrollment.
ORIGINAL	EXCLUSION	E24V010	Females of "childbearing potential" who are sexually active and who refuse to use an "acceptable contraceptive method" up to 6 weeks post second IMP administration on Day 90 (M3). In addition, they must be advised not to donate ova during this period.
ORIGINAL	EXCLUSION	E25V010	A first degree relative is involved in the conduct of this trial.
ORIGINAL	INCLUSION	I01V010	Participant is aged ≥ 4 to ≤ 60 years at the time of signing the informed consent/pediatric assent form.
ORIGINAL	INCLUSION	I02V010	Participant is Japanese male or female.
ORIGINAL	INCLUSION	I03V010	Participant is in good health at the time of entry into the trial as determined by medical history, physical examination (including vital signs) and clinical judgment of the investigator.
ORIGINAL	INCLUSION	I04V010	Participant and/or the participant's legally acceptable representative (LAR) who have signed and dated a written, informed consent/pediatric assent form, and any required privacy authorization prior to the initiation of any trial procedures, and after the nature of the trial has been explained.
ORIGINAL	INCLUSION	I05V010	Participant can comply with trial procedures and is available for the duration of follow-up.

Protocol/ Amendment Version	Category	IETESTCD	Full Text of Criterion
AMENDMENT 1	EXCLUSION	E01V020	Participant has contraindication(s), warning(s) and/or precaution(s) applicable to vaccination with TDV as specified in the IB.
AMENDMENT 1	EXCLUSION	E02V020	Participant has a known hypersensitivity or allergy to any of the IMP components (including excipients of the IMP).
AMENDMENT 1	EXCLUSION	E03V020	Participant has behavioral or cognitive impairment or psychiatric disease that, in the opinion of the investigator, may interfere with the participant's ability to participate in the trial.
AMENDMENT 1	EXCLUSION	E04V020	Participant has a history of progressive or severe neurologic disorder, seizure disorder or neuro-inflammatory disease (eg, Guillain-Barré syndrome).
AMENDMENT 1	EXCLUSION	E05V020	Participant has a clinically significant active infection (as assessed by the investigator) or body temperature >38.0°C (>100.4°F) within 3 days of intended IMP administration on Day 1 (M0). Note: In principle, oral temperature should be measured for body temperature. In cases where it is difficult to measure oral temperature, such as in young children, underarm (axillary) temperature may be used instead.
AMENDMENT 1	EXCLUSION	E06V020	Participant has an illness, or history of any illness that, in the opinion of the investigator, might interfere with the results of the trial or pose additional risk to the participant due to involvement in this trial.

Protocol/ Amendment Version	Category	IETESTCD	Full Text of Criterion
AMENDMENT 1	EXCLUSION	E07V020	Participant has a known or suspected impairment/alteration of immune function, including: a. Chronic administration of oral and/or parenteral steroids at doses considered sufficiently immunosuppressive (eg, ≥ 2 mg/kg body weight/day prednisone/prednisolone [or equivalent] for ≥ 14 consecutive days, or, ≥ 20 mg/day prednisone/prednisolone [or equivalent] for ≥ 14 consecutive days) within 60 days prior to Day 1 (M0). Note: Use of corticosteroids by inhaled, intranasal, intraarticular, bursal, tendon injection, or topical routes is allowed. b. Receipt of blood, immunoglobulins, blood products, and/or plasma derivatives within the 90 days prior to Day 1 (M0). c. Receipt of immunostimulants within 60 days prior to Day 1 (M0). d. Immunosuppressive therapy such as anti-cancer chemotherapy or radiation therapy within 6 months prior to Day 1 (M0). e. Reported or known symptomatic HIV infection or asymptomatic HIV infection when accompanied by evidence of impaired immune function. f. Reported or known Hepatitis B and/or Hepatitis C virus infection. g. Genetic immunodeficiency.
AMENDMENT 1	EXCLUSION	E08V020	Participant has known or suspected abnormalities of splenic or thymic function.
AMENDMENT 1	EXCLUSION	E09V020	Participant has a known bleeding diathesis, or any condition that may be associated with a prolonged bleeding time.
AMENDMENT 1	EXCLUSION	E10V020	Participant has a serious chronic or progressive disease deemed to be preclusive to trial entry, ie, not medically stable according to the judgment of the investigator.
AMENDMENT 1	EXCLUSION	E11V020	Participant is participating in any clinical trial with another investigational product within 30 days prior to Day 1 (M0) or plans to participate in another clinical trial at any time during the conduct of this trial.
AMENDMENT 1	EXCLUSION	E12V020	Participant has previously received a vaccination against flavivirus other than Japanese encephalitis (JE) (investigational or licensed).
AMENDMENT 1	EXCLUSION	E13V020	Participant who received any other vaccines within 14 days (for inactivated vaccines) or 28 days (for live vaccines) prior to Day 1 (M0) or who are planning to receive any vaccine other than IMP within 28 days of IMP administration.

Protocol/ Amendment Version	Category	IETESTCD	Full Text of Criterion
AMENDMENT 1	EXCLUSION	E14V020	Participant who received a coronavirus vaccine within 14 days prior to Day 1 (M0).
AMENDMENT 1	EXCLUSION	E15V020	Participant who received a vaccine authorized for emergency use within 28 days prior to Day 1 (M0).
AMENDMENT 1	EXCLUSION	E16V020	Participant who received any JE vaccines within 28 days prior to Day 1 (M0) or who are planning to receive any JE vaccines during the trial period.
AMENDMENT 1	EXCLUSION	E17V020	Previous participation in any clinical trial of a dengue or other flavivirus candidate vaccine, except for participants who received placebo in those trials.
AMENDMENT 1	EXCLUSION	E18V020	Participant with body mass index (BMI) greater than or equal to 35 kg/m ² (=weight in kg/[height in meters] ²) on Day 1 (M0).
AMENDMENT 1	EXCLUSION	E19V020	Participant who intends to travel to dengue endemic areas during the trial period.
AMENDMENT 1	EXCLUSION	E20V020	Participant with documented or suspected disease caused by a flavivirus and participants with a history of prolonged (≥ 1 year) habitation in a dengue endemic area.
AMENDMENT 1	EXCLUSION	E21V020	Participant with history of substance or alcohol abuse within the past 2 years prior to Day 1 (M0).
AMENDMENT 1	EXCLUSION	E22V020	Female participants who are pregnant (ie, a positive or indeterminate pregnancy test) or breastfeeding.
AMENDMENT 1	EXCLUSION	E23V020	Females of childbearing potential who are sexually active and who have not used any of the acceptable contraceptive methods for at least 2 months prior to Day 1 (M0). a. "Childbearing potential" is defined as status post onset of menarche and not meeting any of the following conditions: menopausal for at least 2 years, status after bilateral tubal ligation for at least 1 year, status after bilateral oophorectomy, or status after hysterectomy. b. "Acceptable contraceptive methods" are defined as one or more of the following: · Hormonal contraceptive. · Barrier method (condom or diaphragm) every time during intercourse. · Intrauterine device. · Monogamous relationship with a vasectomized partner. The partner must have been vasectomized for at least 6 months prior to the participant's trial enrollment.

Protocol/ Amendment Version	Category	IETESTCD	Full Text of Criterion
AMENDMENT 1	EXCLUSION	E24V020	Females of “childbearing potential” or non-sterilized males, who refuse to use an “acceptable contraceptive method” up to 6 weeks post second IMP administration on Day 90 (M3). In addition, they must be advised not to donate ova or sperm during this period.
AMENDMENT 1	EXCLUSION	E25V020	A first degree relative is involved in the conduct of this trial.
AMENDMENT 1	INCLUSION	I01V020	Participant is aged ≥ 4 to ≤ 60 years at the time of signing the informed consent/pediatric assent form.
AMENDMENT 1	INCLUSION	I02V020	Participant is Japanese male or female.
AMENDMENT 1	INCLUSION	I03V020	Participant is in good health at the time of entry into the trial as determined by medical history, physical examination (including vital signs) and clinical judgment of the investigator.
AMENDMENT 1	INCLUSION	I04V020	Participant and/or the participant's legally acceptable representative (LAR) who have signed and dated a written, informed consent/pediatric assent form, and any required privacy authorization prior to the initiation of any trial procedures, and after the nature of the trial has been explained.
AMENDMENT 1	INCLUSION	I05V020	Participant can comply with trial procedures and is available for the duration of follow-up.