

LECTURE 7

LECTURE OUTLINE

- Preservation of closure under partial minimization
 - Hyperplanes
 - Hyperplane separation
 - Nonvertical hyperplanes
 - Min common and max crossing problems
-

- Question: If $F(x, z)$ is closed, is $f(x) = \inf_z F(x, z)$ closed? Can be addressed by using the relation

$$P(\text{epi}(F)) \subset \text{epi}(f) \subset \text{cl}\left(P(\text{epi}(F))\right),$$

where $P(\cdot)$ denotes projection on the space of (x, w) , i.e., $P(x, z, w) = (x, w)$.

- Closedness of f is guaranteed if the closure of $\text{epi}(F)$ is preserved under the projection operation $P(\cdot)$.

PARTIAL MINIMIZATION THEOREM

- Let $F : \Re^{n+m} \mapsto (-\infty, \infty]$ be a closed proper convex function, and consider $f(x) = \inf_{z \in \Re^m} F(x, z)$.
- If there exist $\bar{x} \in \Re^n$, $\bar{\gamma} \in \Re$ such that the set


$$\{z \mid F(\bar{x}, z) \leq \bar{\gamma}\}$$

is nonempty and compact, then f is convex, closed, and proper. Also, for each $x \in \text{dom}(f)$, the set of minima of $F(x, \cdot)$ is nonempty and compact.


Proof: (Outline) By the hypothesis, there is no nonzero y such that $(0, y, 0) \in R_{\text{epi}(F)}$. Also, all the nonempty level sets

$$\{z \mid F(x, z) \leq \gamma\}, \quad x \in \Re^n, \quad \gamma \in \Re,$$

have the same recession cone, which by the hypothesis, is equal to $\{0\}$.





HYPERPLANES



- A *hyperplane* is a set of the form $\{x \mid a'x = b\}$, where a is nonzero vector in \mathbb{R}^n and b is a scalar.
- We say that two sets C_1 and C_2 are *separated* by a hyperplane $H = \{x \mid a'x = b\}$ if each lies in a different closed halfspace associated with H , i.e.,
either $a'x_1 \leq b \leq a'x_2, \quad \forall x_1 \in C_1, \forall x_2 \in C_2,$
or $a'x_2 \leq b \leq a'x_1, \quad \forall x_1 \in C_1, \forall x_2 \in C_2.$
- If \bar{x} belongs to the closure of a set C , a hyperplane that separates C and the singleton set $\{\bar{x}\}$ is said be *supporting* C at \bar{x} .



VISUALIZATION

- Separating and supporting hyperplanes:




- A separating $\{x \mid a'x = b\}$ that is disjoint from C_1 and C_2 is called *strictly separating*:

$$a'x_1 < b < a'x_2, \quad \forall x_1 \in C_1, \forall x_2 \in C_2.$$



SUPPORTING HYPERPLANE THEOREM

- Let C be convex and let \bar{x} be a vector that is not an interior point of C . Then, there exists a hyperplane that passes through \bar{x} and contains C in one of its closed halfspaces.



Proof: Take a sequence $\{x_k\}$ that does not belong to $\text{cl}(C)$ and converges to \bar{x} . Let \hat{x}_k be the projection of x_k on $\text{cl}(C)$. We have for all $x \in \text{cl}(C)$

$$a'_k x \geq a'_k x_k, \quad \forall x \in \text{cl}(C), \quad \forall k = 0, 1, \dots,$$

where $a_k = (\hat{x}_k - x_k) / \|\hat{x}_k - x_k\|$. Let a be a limit point of $\{a_k\}$, and take limit as $k \rightarrow \infty$. **Q.E.D.**

SEPARATING HYPERPLANE THEOREM

- Let C_1 and C_2 be two nonempty convex subsets of \mathbb{R}^n . If C_1 and C_2 are disjoint, there exists a hyperplane that separates them, i.e., there exists a vector $a \neq 0$ such that

$$a'x_1 \leq a'x_2, \quad \forall x_1 \in C_1, \forall x_2 \in C_2.$$

Proof: Consider the convex set

$$C_1 - C_2 = \{x_2 - x_1 \mid x_1 \in C_1, x_2 \in C_2\}.$$


Since C_1 and C_2 are disjoint, the origin does not belong to $C_1 - C_2$, so by the Supporting Hyperplane Theorem, there exists a vector $a \neq 0$ such that

$$0 \leq a'x, \quad \forall x \in C_1 - C_2,$$

which is equivalent to the desired relation. **Q.E.D.**

STRICT SEPARATION THEOREM

- *Strict Separation Theorem*: Let C_1 and C_2 be two disjoint nonempty convex sets. If C_1 is closed, and C_2 is compact, there exists a hyperplane that strictly separates them.




Proof: (Outline) Consider the set $C_1 - C_2$. Since C_1 is closed and C_2 is compact, $C_1 - C_2$ is closed. Since $C_1 \cap C_2 = \emptyset$, $0 \notin C_1 - C_2$. Let $\bar{x}_1 - \bar{x}_2$ be the projection of 0 onto $C_1 - C_2$. The strictly separating hyperplane is constructed as in (b).

- *Note*: Any conditions that guarantee closedness of $C_1 - C_2$ guarantee existence of a strictly separating hyperplane. However, there may exist a strictly separating hyperplane without $C_1 - C_2$ being closed.

ADDITIONAL THEOREMS

- *Fundamental Characterization*: The closure of the convex hull of a set $C \subset \mathbb{R}^n$ is the intersection of the closed halfspaces that contain C .
- We say that a hyperplane *properly separates* C_1 and C_2 if it separates C_1 and C_2 and does not fully contain both C_1 and C_2 .




- *Proper Separation Theorem*: Let C_1 and C_2 be two nonempty convex subsets of \mathbb{R}^n . There exists a hyperplane that properly separates C_1 and C_2 if and only if

$$\text{ri}(C_1) \cap \text{ri}(C_2) = \emptyset.$$

MIN COMMON / MAX CROSSING PROBLEMS


- We introduce a pair of fundamental problems:
- Let M be a nonempty subset of \mathbb{R}^{n+1}
 - (a) *Min Common Point Problem*: Consider all vectors that are common to M and the $(n + 1)$ st axis. Find one whose $(n + 1)$ st component is minimum.
 - (b) *Max Crossing Point Problem*: Consider “non-vertical” hyperplanes that contain M in their “upper” closed halfspace. Find one whose crossing point of the $(n + 1)$ st axis is maximum.



- Need to study “nonvertical” hyperplanes.

NONVERTICAL HYPERPLANES

- A hyperplane in \Re^{n+1} with normal (μ, β) is non-vertical if $\beta \neq 0$.
- It intersects the $(n+1)$ st axis at $\xi = (\mu/\beta)' \bar{u} + \bar{w}$, where (\bar{u}, \bar{w}) is any vector on the hyperplane.



- A nonvertical hyperplane that contains the epigraph of a function in its “upper” halfspace, provides lower bounds to the function values.
- The epigraph of a proper convex function does not contain a vertical line, so it appears plausible that it is contained in the “upper” halfspace of some nonvertical hyperplane.

NONVERTICAL HYPERPLANE THEOREM

• Let C be a nonempty convex subset of \mathbb{R}^{n+1} that contains no vertical lines. Then:

- (a) C is contained in a closed halfspace of a nonvertical hyperplane, i.e., there exist $\mu \in \mathbb{R}^n$, $\beta \in \mathbb{R}$ with $\beta \neq 0$, and $\gamma \in \mathbb{R}$ such that $\mu'u + \beta w \geq \gamma$ for all $(u, w) \in C$.
- (b) If $(\bar{u}, \bar{w}) \notin \text{cl}(C)$, there exists a nonvertical hyperplane strictly separating (\bar{u}, \bar{w}) and C .

Proof: Note that $\text{cl}(C)$ contains no vert. line [since C contains no vert. line, $\text{ri}(C)$ contains no vert. line, and $\text{ri}(C)$ and $\text{cl}(C)$ have the same recession cone]. So we just consider the case: C closed.

(a) C is the intersection of the closed halfspaces containing C . If all these corresponded to vertical hyperplanes, C would contain a vertical line.

(b) There is a hyperplane strictly separating (\bar{u}, \bar{w}) and C . If it is nonvertical, we are done, so assume it is vertical. “Add” to this vertical hyperplane a small ϵ -multiple of a nonvertical hyperplane containing C in one of its halfspaces as per (a).