

## Generating Functions 3 - Extraction & Recurrences

---

We continue to work with rational generating functions  $A(x)$ , so we have

$$A(x) = \frac{p(x)}{q(x)} = \sum_{n=0}^{\infty} a_n x^n$$

Our primary interest will be in going back and forth between expressing  $A(x)$  as a quotient of polynomials and in terms of its coefficients.

## Coefficient Extraction

**Recall:** For a generating function  $A(x)$  we let  $[x^k]A(x)$  denote the coefficient of  $x^k$  in  $A(x)$ .

Coefficient extraction is the process of determining the coefficients of a generating function. The key to coefficient extraction for rational generating functions is **partial fractions**

**Example:** Find values for  $A, B, C$  so that the expression below is true, then use this to determine  $[x^n]D(x)$

$$D(x) = \frac{1}{(x-3)(x-2)^2} = \frac{A}{x-3} + \frac{B}{x-2} + \frac{C}{(x-2)^2}$$

mult + b  
 $(x-3)(x-2)^2$

$$1 = A(x-2)^2 + B(x-3)(x-2) + C(x-3)$$

$$= Ax^2 - 4Ax + 4A + Bx^2 - 5Bx + 6B + Cx - 3C$$

$$= (A+B)x^2 + (-4A-5B+C)x + (4A+6B-3C)$$

0

0

1

$$C = -1 \quad A = 1 \quad B = -1$$





This process works in general.

**Partial Fractions.** Let  $q(x)$  be a polynomial which can be factored as  $q(x) = (x - r_1)^{d_1}(x - r_2)^{d_2} \dots (x - r_k)^{d_k}$  then there exist constants so that

$$\frac{1}{q(x)} = \frac{A_{1,1}}{x - r_1} + \frac{A_{1,2}}{(x - r_1)^2} + \dots + \frac{A_{1,d_1}}{(x - r_1)^{d_1}} + \dots + \frac{A_{k,d_k}}{(x - r_k)^{d_k}}$$

**Example:** Although we will not solve it, there exist constants  $A, B, C, D, E, F$  so that the following expression is valid:

$$\frac{1}{(x - 1)(x - 2)^2(x - 3)^3} = \frac{A}{(x - 1)} + \frac{B}{(x - 2)} + \frac{C}{(x - 2)^2} + \frac{D}{(x - 3)} + \frac{E}{(x - 3)^2} + \frac{F}{(x - 3)^3}$$

**Note:** We can use this together with the formula

$$\frac{1}{(1 - x)^k} = \sum_{n=0}^{\infty} \binom{n + k - 1}{n} x^n$$

To do coefficient extraction whenever we have a rational function and we have factored the denominator.

## Recurrences

Generating functions also provide a natural setting to work with and solve recurrence relations.

**Example:** Consider the recurrence relation

$$a_0 = 0 \quad \text{and} \quad a_1 = 1$$

$$a_n - 5a_{n-1} + 6a_{n-2} = 0 \quad \text{for } n \geq 2$$

Define the generating function  $A(x) = \sum_{n=0}^{\infty} a_n x^n$  What is

$$A(x) - 5xA(x) + 6x^2A(x)?$$

Use the above expression to express  $A(x)$  as a rational function

Use coefficient extraction to find the coefficients of  $A(x)$ .