

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/258357574>

# Apatite thermochronology in modern geology

Article in Geological Society London Special Publications · September 2009

DOI: 10.1144/SP324.1

CITATIONS

81

READS

657

3 authors:



[Frank Lisker](#)

Universität Bremen

91 PUBLICATIONS 1,250 CITATIONS

[SEE PROFILE](#)



[Barbara Ventura](#)

Universität Bremen

9 PUBLICATIONS 372 CITATIONS

[SEE PROFILE](#)



[U. A. Glasmacher](#)

Universität Heidelberg

152 PUBLICATIONS 2,231 CITATIONS

[SEE PROFILE](#)

# Apatite thermochronology in modern geology

F. LISKER<sup>1\*</sup>, B. VENTURA<sup>1</sup> & U. A. GLASMACHER<sup>2</sup>

<sup>1</sup>*Fachbereich Geowissenschaften, Universität Bremen, PF 330440, 28334 Bremen, Germany*

<sup>2</sup>*Institut für Geowissenschaften, Ruprecht-Karls-Universität Heidelberg, Im Neuenheimer Feld 234, 69120 Heidelberg, Germany*

*\*Corresponding author (e-mail: flisker@uni-bremen.de)*

**Abstract:** Fission-track and (U–Th–Sm)/He thermochronology on apatites are radiometric dating methods that refer to thermal histories of rocks within the temperature range of 40°–125 °C. Their introduction into geological research contributed to the development of new concepts to interpreting time-temperature constraints and substantially improved the understanding of cooling processes within the uppermost crust. Present geological applications of apatite thermochronological methods include absolute dating of rocks and tectonic processes, investigation of denudation histories and long-term landscape evolution of various geological settings, and basin analysis.

Thermochronology may be described as the quantitative study of the thermal histories of rocks using temperature-sensitive radiometric dating methods such as <sup>40</sup>Ar/<sup>39</sup>Ar and K–Ar, fission track, and (U–Th)/He (Berger & York 1981). Amongst these different methods, apatite fission track (AFT) and apatite (U–Th–Sm)/He (AHe) are now, perhaps, the most widely used thermochronometers as they are the most sensitive to low temperatures (typically between 40 and c. 125 °C for durations of heating and cooling in excess of 10<sup>6</sup> years), ideal for investigating the tectonic and climate-driven surficial interactions that take place within the top few (<5 km) kilometres of the Earth's crust. These processes govern landscape evolution, influence climate and generate the natural resources essential to the wellbeing of mankind.

This introductory chapter provides a brief overview of apatite thermochronology and its application to geological studies. We focus on three topics: (1) methodological developments; (2) concepts and strategies for the interpretation of thermochronological data; and (3) applications to various geodynamic settings. For more detailed insights on apatite thermochronology the reader is referred to published reviews by Green *et al.* (1986, 1989b), Laslett *et al.* (1987), Duddy *et al.* (1988), Wagner & Van den Haute (1992), R. W. Brown *et al.* (1994), Gallagher *et al.* (1998), Gleadow *et al.* (2002), Ehlers & Farley (2003) and Reiners & Brandon (2006).

## Fission-track thermochronology

### Basics of the method

Fission-track thermochronology/-chronometry (for differentiation cf. Reiners *et al.* 2005) is based on

the analysis of radiation damage trails ('fission tracks') in uranium-bearing, non-conductive minerals and glasses. It is routinely applied on the minerals apatite, zircon and titanite. Fission tracks are produced continuously through geological time as a result of the spontaneous fission of <sup>238</sup>U atoms. They are submicroscopic features with an initial width of approximately 10 nm and a length of up to 20 µm (Paul & Fitzgerald 1992) that can be revealed by chemical etching. Crucially, fission tracks are semi-stable features that can self-repair (shorten and eventually disappear) by a process known as annealing at a rate that is a function of both time and temperature. The extent of any track shortening (exposure to elevated temperatures) in a sample can be quantified by examining the distribution of fission-track lengths.

The determination of a fission-track age (a number that relates to the observable track density) depends on the same general equation as any radioactive decay scheme: it requires an estimate of the relative abundance of the parent isotope and of the daughter product. However, unlike most methods of radiometric dating, it measures the effect, rather than the product, of a radioactive decay scheme, that is it refers to the number of <sup>238</sup>U atoms and the number of spontaneous fission tracks per unit volume. This fission-track density is obtained by counting the number of spontaneous tracks intersecting a polished internal surface of a mineral grain viewed under high magnification (1000×–1250×) using an optical microscope. Depending on the sample and aims of the study, a typical fission-track sample age consists of a weighted mean of 20–100 single-grain ages. Further details and background information on practical aspects of fission-track age determination are provided by, for example, Fleischer *et al.* (1975), Naeser (1979) and Donelick *et al.* (2005).

An important dimension to fission-track thermochronology is the **semi-stable nature of tracks**, whereby annealing can change the significance of a measured age. The observable density of spontaneous tracks in a sample (age) is a function of track length (probability of intersecting the plane of observation). **All newly formed tracks** in apatite have a length of approximately 16  $\mu\text{m}$  (c. 11  $\mu\text{m}$  in zircon). If a sample (e.g. a volcanic apatite) was created at 10 Ma and then resided at low temperatures (<40 °C), the population of tracks reduces in length to a mean value of approximately 15  $\mu\text{m}$  causing an insignificant (not resolvable) reduction in track density and a measured age within an error of 10 Ma. However, if, during its history, the same sample experienced elevated temperatures (but not sufficient to cause total resetting) in its history there will be significant track shortening to a level defined by the maximum heating. This will cause a reduction in observable track density and, therefore, measurable age.

### *Some milestones in the evolution of the fission-track method*


Despite early recognition of fission tracks (Baumhauer 1894; Silk & Barnes 1959), it was not until the early 1960s that their application to geological dating was first proposed (Price & Walker 1963) and subsequently developed (Wagner 1966; Gentner *et al.* 1967; Naeser 1967). Early dating studies were tasked with finding practical ways to etch tracks, measure uranium contents that mirror the dated grain and define the time–temperature stability fields of fission tracks in different uranium-bearing minerals, tektites and glasses. Studies conducted between 1970 and 1983 highlighted the fundamental issues that needed to be resolved in order to enable routine and accurate age determination. Foremost was a lack of consensus on the value of the spontaneous fission decay constant for  $^{238}\text{U}$ , to be used in the equation for age calculation. In order to circumvent this and other fundamental problems associated with the measurement of neutron fluence, Hurford & Green (1983) advanced a suggestion made by Fleischer and Hart at a meeting in Austria in 1971 for a comparative approach to AFT dating through the use of a proportionality constant. The resultant **'Zeta' calibration method** (Hurford & Green 1983) has become the standard approach to **fission-track age determination** (Hurford 1990).

Until 1980 most fission-track ages were calculated using a pooled age, based on the ratio of total counts of spontaneous and induced tracks. However, Green (1981) highlighted the need for an alternative method for calculating an age when

there is significant (or over-) dispersion within the population of single-grain ages. Green pointed out that where there is evidence for heterogeneity (extra Poissonian variation) within a dataset, detected by statistical tests such as  $\chi^2$  (Galbraith 1981), the conventional pooled age, based on the ratio of the number of spontaneous and induced tracks ( $N_s/N_i$ ), with its Poisson standard error, becomes meaningless. An alternative approach based on the mean of ratios of individual track densities ( $\rho_s/\rho_i$ ) was used to give a larger estimate of the error to allow for an extra-Poisson component in the dispersion of single-grain ages, but this approach implied a sample should have a single age. In reality, there are a number of different causes of heterogeneity within a dataset (beyond a bad experiment), such as variable responses to partial resetting due to variations in apatite grain composition (see later) and/or a range of provenance ages. Thus, it is important to assess to what the over-dispersion is due rather than make to allowance for it with larger errors.

In 1984 Hurford *et al.* proposed the use of **probability density diagrams**, a type of continuous histogram that plots each grain-age error as a Gaussian density function, as a way of visualizing a mixed age dataset. However, this type of approach can obscure useful information by inappropriately weighting it with poor information (i.e. an overlap effect associated with broad, imprecise, peaks). To overcome this problem Galbraith developed the **radial plot** (Galbraith 1988, 1990) (Fig. 1), which is now routinely used across the chronological community. Coupled to this development Galbraith & Laslett (1993) also produced the widely adopted random effects model that gives a central age estimate of the population of grains ages with a relative standard deviation of the population of ages known as the age dispersion (normally expressed as a percentage variation).

Having established protocols to measure fission-track ages and assess data quality and structure, the next major developments were related to understanding the significance of the determined ages. Whilst track-length measurement had been used to detect track annealing more or less since the methods inception, it was not until the mid 1980s that studies demonstrated the utility of such data. Advances included moving away from using semi-tracks (projected track) to length measurement based on surface-parallel confined tracks. Although more numerous, semi-tracks contain less information and have significant sources of bias, particularly towards longer lengths (Laslett *et al.* 1994). A key paper by Gleadow *et al.* (1986b) demonstrated the utility of confined track measurement and laid the foundation for data interpretation based on thermal history.



**Fig. 1.** Radial plots were designed to graphically display single-grain age estimates, taking into account different standard errors (Galbraith 1988, 1990; example from Ventura *et al.* 2009). Single-grain ages ( $z$ ) with standard error,  $\sigma$ , are plotted (point  $x, y$ ) according to  $x$  (precision)  $= 1/\sigma$  and  $y$  (standard estimate)  $= (z - z_0)/\sigma$ , where  $z_0$  is the central age. The error attached to each point is standardized on the  $y$ -scale. The value of the age ( $z$ ) and the  $2\sigma$  uncertainty can be read off the  $z$ -scale by extrapolating lines from point 0, 0 through the plotted age (point  $x, y$ ) and projecting onto the radial age axis.

Contrary to other radiometric dating methods where the measured age equates to the time a sampled cooled below its closure temperature, the fission-track age records cooling through a temperature interval between total resetting of fission tracks and relative stability, known as the **partial annealing zone or PAZ** (Wagner 1979). The temperatures relating to the PAZ were defined by systematic investigation of the annealing of fission tracks across laboratory and geological timescales, monitored by changes in confined track-length distribution (Green & Durrani 1977; Gleadow & Duddy 1981; Laslett *et al.* 1982, 1987; Gleadow *et al.* 1986a; Green *et al.* 1986, 1989a, b; Duddy *et al.* 1988; Green 1988; Crowley *et al.* 1991; Carlson *et al.* 1999; Barbarand *et al.* 2003). Quantification of the time–temperature conditions that control the annealing of fission tracks in apatite provide the means to interpret fission track-ages by linking the level of track-length shortening and density (age) reduction to sample thermal history (e.g. Gleadow *et al.* 1986b). Typical temperature ranges for the PAZ for heating durations of  $10^7$  Ma are 60–110 °C for fluorapatite (e.g. Gleadow & Duddy 1981), 170–330 °C for zircon (Zaun & Wagner 1985; Yamada *et al.* 1995) and 265–310 °C for titanite (Coyle & Wagner 1998). Above the upper-temperature values fission tracks undergo total annealing, which removes all traces of fission tracks from the host crystal lattice.

Annealing studies recognized that **track fading** in apatite is not only governed by **temperature**, but also by **heating duration**, **chemical composition** and **crystallographic orientation**. Early descriptions of time-dependence on annealing were presented as Arrhenius plots, which generally had a fanning like form. Green *et al.* (1985) noted that **fanning plots** were probably the result of variable apatite grain composition, such that in effect these fanning plots were no more than a series of overlapping parallel plots. A series of isothermal fission-track annealing studies on apatites (Green *et al.* 1986; Crowley *et al.* 1990; Carlson *et al.* 1999; Barbarand *et al.* 2003) showed that substitution of fluoride and hydroxide ions by chloride ions appears to induce the greatest effects, although other substitutions on the halogen site can also have an influence but have proven much harder to deconvolve from laboratory timescale track-annealing experiments (O'Sullivan & Parrish 1995; Carlson *et al.* 1999). As a consequence of these studies, AFT data interpretation generally takes into account grain composition either by direct measurement of the Cl-content [generally by electron probe micro-analyser (EPMA)] or by assessing grain bulk composition by measuring the solubility of apatite through the  $c$ -axis parallel length of track etch pits ( $D_{\text{par}}$  of Donelick 1991; Burtner *et al.* 1994; cf. also Murrell *et al.* 2009). Green & Durrani (1977) first described the influence of the crystallographic orientation of spontaneous fission tracks on the annealing of those tracks. Tracks orthogonal to the  $c$ -axis anneal more rapidly than tracks parallel to the  $c$ -axis (Green 1988). This anisotropy increases with annealing (Green 1981; Laslett *et al.* 1984; Donelick *et al.* 1990; Galbraith *et al.* 1990; Donelick 1991).

The developments outlined above (and many others not cited) have helped establish AFT analysis as a routinely used tool in geological studies, examples of which are published in this Special Publication. Methodological developments continue and it is now possible, following marked improvements in the precision of laser ablation inductively coupled plasma mass spectrometry (LA-ICP-MS) technology, to make direct measurements of uranium content within the counted grains of apatite, providing a fast and precise alternative to neutron irradiation (e.g. Hasebe *et al.* 2004, 2009). A further advance is the arrival of a functioning system for automatically determining the fission-track density and measuring fission-track lengths (Gleadow *et al.* 2009). This will not only increase speed and reliability of data acquisition, but also provides new options, especially with respect to the size (quality) of density and track length datasets.

## (U–Th–Sm)/He thermochronology

The last decade has seen the extension of low-temperature studies to include AHe dating. **Radioactive decay of the elements uranium and thorium to stable helium** represents one of the earliest radiometric schemes available to geologists to investigate the ages of rocks and minerals (Rutherford 1905). Despite initial promise (e.g. Keevil 1943), the (U–Th)/He method was discounted as a dating technique until Lippolt *et al.* (1982) proposed its application as a low-temperature thermochronological tool. Zeitler *et al.* (1987) followed Lippolt *et al.*'s suggestion that, rather than defining a sample's age (implicitly its formation age), (U–Th)/He data provide useful constraints on the sample's thermal history, similar to AFT analysis, and further developed the technique. Fundamental experimental work by Farley and co-workers at Caltech through the 1990s resulted in the development of accurate and precise instrumentation for the extraction and measurement of helium. In practice, **an AHe age is obtained by measuring radiogenic  $^4\text{He}$  trapped in apatite grains by laser or furnace outgassing, and then measuring the relative amounts of uranium and thorium in the sample by solution ICP-MS.** A careful selection of inclusion and crack-free idiomorphic apatite crystals with homogeneous U, Th and Sm distribution is essential for this procedure.

AHe thermochronology **relies on the accumulation of  $^4\text{He}$  during the  $\alpha$ -disintegration of  $^{238}\text{U}$ ,  $^{235}\text{U}$ ,  $^{232}\text{Th}$ , their daughter products and  $^{147}\text{Sm}$ .** The **closure temperature ( $T_C$ )** of mineral grains is dependent on **activation energy, a geometry factor for the crystal shape, thermal diffusivity ( $D_0$ ), the length of the average diffusion pathway from the interior to the surface of the grain and the cooling rate at closure temperature.** In addition,  $^4\text{He}$  diffusion in apatite is impeded by radiation-induced damage to the apatite structure. Therefore, the kinetics is an evolving function of time. The  $^4\text{He}$  production–diffusion model predicts that the effective  $^4\text{He}$  closure temperature of apatite will vary with cooling rate and effective U- and Th-concentration, and may differ from the commonly assumed  $T_C$  of  $75^\circ\text{C Ma}^{-1}$  by up to  $\pm 15^\circ\text{C}$  (e.g. Farley *et al.* 1996; Wolf *et al.* 1996, 1998; Farley 2000, 2002; Shuster *et al.* 2006; Flowers *et al.* 2009; Shuster & Farley 2009). During radioactive decay alpha ( $\alpha$ )-particles are emitted with high kinetic energy and travel significant distances. This poses a complication for the He dating method, as  $\alpha$ -particles may be ejected out of the crystal being dated or injected from the surrounding mineral grains.

Farley *et al.* (1996) and Reiners *et al.* (2004) pointed out that implantation or ejection of  $\alpha$ -particles may generally obscure (U–Th–Sm)/

He dating, whereas He implantation needs to be considered if dealing with low U–Th–Sm phases. Therefore, Farley *et al.* (1996) propose a correction for these effects either numerically by correcting the measured He ages against the specific  $\alpha$  retentivity ( $F_T$ ) or by removal of the outer rim of the crystal (by chemical dissolution or mechanical abrasion) prior to dating. **While correcting for  $\alpha$  ejection** has become a routinely performed practice in recent years, Spiegel *et al.* (2009) demonstrate the potential for possible overcorrection, indicating that abrading the outer rim of a crystal may be favourable. However, mechanical abrasion may lead to erroneous ages when crystals are strongly zoned with respect to uranium and thorium (Farley *et al.* 1996). In this case, a U–Th–Sm zonation-dependent  $\alpha$  correction, as proposed by Hourigan *et al.* (2005), needs to be applied.

Usually, an apatite  $T_C$  in the range of  $70 \pm 7^\circ\text{C}$  (for a monotonic cooling rate of  $10^\circ\text{C Ma}^{-1}$ , a subgrain domain size  $>60\ \mu\text{m}$ , an activation energy of about  $36\ \text{kcal mol}^{-1}$  and a  $\log(D_0)$  of  $7.7 \pm 0.6\ \text{cm}^2\ \text{s}^{-1}$ ), and a He partial retention zone of between  $40$  and  $75^\circ\text{C}$ , are assumed. The He production–diffusion model relies on homogenous distribution of U, Th and Sm in secular equilibrium, He loss confined to volume diffusion, and spherical diffusion geometry. Meesters & Dunai (2002a, b) generalized a production–diffusion equation to diffusion domains of various shapes and arbitrary cooling histories. Their set of equations allows  $\alpha$  ejection corrected ages to be calculated and accounts for the non-homogeneous distribution of U, Th and Sm.

Preliminary studies on the applicability of (U–Th)/He thermochronology on zircon and titanite suggest closure temperatures about  $80$  and  $130^\circ\text{C}$  higher than for apatite. These systems have the potential to close the gap between the various mineral diffusion temperatures of the  $^{40}\text{Ar}/^{39}\text{Ar}$  thermochronological system, and the zircon and titanite fission-track annealing temperatures on one side, and AFT and AHe on the other side (e.g. Farley 2002; Reiners & Brandon 2006; Dobson *et al.* 2009). Moreover, (U–Th)/He analysis of magnetite ( $T_C$ :  $200^\circ\text{C}/10\ \text{Ma}$ ) opens new perspectives for dating volcanic rocks (Blackburn *et al.* 2007).

## Modelling of low-temperature thermochronological data

Recovery of thermal history information from AFT and AHe datasets is contingent upon a quantitative understanding of how the combination of time and temperature controls fission-track annealing and helium diffusion. Early attempts to use AFT data to track cooling were based on qualitative

indicators (e.g. Wagner & Reimer 1972; Gleadow *et al.* 1986a, b): samples with a single population of long track lengths indicate rapid cooling in contrast to slower and/or stepped cooling paths, which result in shorter and/or more complex length distributions. However, this type of approach limits interpretation to relative rates of cooling and provides no constraints on timing or temperatures. With the advent of annealing studies, algorithms were formulated to describe the time–temperature dependencies of track annealing. The models based on laboratory experiments were extrapolated to geological timescales and verified against well-constrained data from the geological record. The first and, previously, most widely used model (Laslett *et al.* 1987) describes the annealing behaviour of a single type of apatite (Durango). With the realization that annealing is also influenced by grain composition, new experiments extended the annealing database. Carlson *et al.* (1999) published the first multi-kinetic annealing model. In 2007 this model was updated to include the dataset of Barbarand *et al.* (2003) and improved data-fitting techniques. The new multi-kinetic model, now in wide use, is based on 579 experiments and 26 compositionally different types of apatite (Ketcham *et al.* 2007).

Equipped with quantitative descriptions of track annealing it is possible to extract sample thermal histories by using forward or data inversion modelling techniques. Early programs focused on forward modelling data to check annealing models against well-constrained geological examples (e.g. Green *et al.* 1989b) and as a guide to the interpretation of real samples (e.g. Willett 1992). However, forward modelling is not a very efficient means of finding solutions for unknown or poorly constrained samples and, since there is no unique solution to a given dataset, such an approach is open to user bias. Data-driven inverse modelling helps to reduce this bias. Most commonly adopted and publicly available AFT modelling programs are Monte Trax (Gallagher 1995, designed for Apple Macintosh), AFTINV (Issler 1996; Willet 1997), AFT-Solve (Ketcham *et al.* 2000) and HeFTy (Ketcham 2005, all Windows). These programs differ in modelling approach (Monte Carlo and/or genetic algorithm), annealing models, input parameters and statistical tests to evaluate the level of fit between model results and observed data (cf. Ehlers *et al.* 2005; Ketcham 2005). Some of the modelling programs refer only to AFT data, whilst others derive cooling histories from combined AFT, AHe and VR data (e.g. HeFTy). DECOMP (Meesters & Dunai 2002a, b) is a popular program to model thermal histories from AHe data. Most recent developments in modelling include a strategy for modelling sample thermal histories jointly (Gallagher *et al.* 2005) and the use of Markov Chain Monte

Carlo (MCMC) methods to address the problem of characterizing uncertainties in modelled thermal histories in two and three dimensions (2D and 3D) (e.g. Stephenson *et al.* 2006).

### Some concepts in apatite thermochronology

Methodological advancements were accompanied by the development of new strategies for the interpretation of thermochronological data, as the derivation of time–temperature constraints and the conversion of temperature information into geological and geomorphological processes.

#### *Fission-track age types*

An important part of fission-track thermochronology relates to using the distribution of measured track lengths in a sample to determine whether the measured age directly records an ‘event’ or a more complex thermal history. Early on in the AFT methods developmental history Günther Wagner suggested a classification of AFT ‘ages’ as *event ages*, *cooling ages* and *mixed ages* based on sample thermal history (Wagner 1972).


According to this concept, an *event age* refers to a rock that cools rapidly through the PAZ (e.g. a volcanic rock) and resides at low, near-surface, temperatures thereafter. The AFT age is essentially identical to the age of entrance into the PAZ, and is associated with a narrow distribution of track lengths about a mean value of 15  $\mu\text{m}$ . Slow linear cooling of a sample through the PAZ produces a *cooling age* that is significantly younger than the entrance into the PAZ, with a broader and shorter track-length distribution. Such a pattern is relatively common in old basement terrains that have undergone cooling over very long periods of geological time. *Mixed ages* refer to an at least two-stage cooling history, in which the first generation of tracks resides at relative higher temperatures within the PAZ, prior to final cooling to lower temperatures. The resulting track-length distribution in such cases is typically bimodal with a short peak representing the higher temperature tracks and a second, long peak being added after final cooling. Crucially, both *cooling* and *mixed ages* have no direct significance in terms of the timing of any geological event.

#### *Uplift, exhumation and denudation*

With the availability of low-temperature chronological data, the resolution of cooling events improved considerably, and it became possible to analyse increasingly younger and shallower processes. Early AFT studies usually interpreted







**Fig. 3.** The concept of an exhumed PAZ (adapted from Fitzgerald *et al.* 1995; Gallagher *et al.* 1998). The left panel illustrates the pre-denudation apatite fission-track age crustal profile, with the initial age as  $t_0$ . Denudation at time  $t_1$  exposes different levels of this pre-cooling profile, while the deeper samples begin to retain tracks (central panel). The right panel shows the expected trend in the fission-track data with respect to elevation, that is age increases. The length distribution has two components: tracks formed prior to cooling (dark shading) and those formed after cooling (light shading). The latter are all long, and the composite length distribution depends on the relative proportion of these two components. Only the data below the break in slope (marked by an asterisk) provide the timing for the onset of the cooling/denudation event.

Another advantage of analysing vertical AFT profiles is the possibility of calculating the palaeo-geothermal gradient that existed prior to the onset of cooling (Gleadow & Duddy 1981; Bray *et al.* 1992). The palaeo-geothermal gradient can be determined by weighted least-squares regression of modelled maximum palaeotemperatures against sample

elevation. Palaeo-geothermal gradients are not only invaluable for estimating burial depths and amounts of denudation, but also bear important constraints on modes and mechanisms of the related processes (e.g. rifting). A comparison of the palaeo-geothermal gradient with a recent one allows the cause of high palaeotemperatures, and the cause of



the subsequent cooling to the present temperatures, to be determined.

### *Thermochronological data patterns*

Thermochronological transects (horizontal profiles) across geological units or structures, or regional patterns of the AFT parameters, often allow exhumation trends to be detected or distinct igneous or tectonic events to be verified. Before modelling programs were available, the characteristic relationship between AFT ages, track-length distributions and thermal history was used to qualitatively constrain amount and timing of cooling (denudation) on a regional scale. A plot of AFT age against mean track length of an area that has undergone cooling shows a characteristic 'banana' or 'boomerang' shape that results from the mixing of two cooling components (Green 1986). In such plots, the longest mean track lengths are preserved in samples that experienced rapid cooling without subsequent reheating. Between the two end points of a boomerang there exists a series of transitional bimodal track-length distributions where the abundance of inherited annealed tracks becomes reduced in favour of newly formed tracks while the AFT ages of the samples decrease.

### *Apatite thermochronology and topography*

*Topographic effects and fast rates of cooling.* The temptation to collect vertical profiles in active mountain belts where there is considerable relief can introduce a number of interpretative problems for thermochronological data, especially if measured ages are young (<5 Ma). Young AFT (or AHe) ages require rapid rates of cooling and high rates of rock uplift and exhumation, but the calculation of true exhumation rates in such young samples is problematic owing to the combined influences of topographic wavelength and thermal advection. High rates of rock uplift cause perturbation of upper-crust thermal structure as heat is advected at rates that exceed normal heat loss by conduction, driving isotherms closer to the Earth's surface (e.g. modelling studies of Stüwe *et al.* 1994; Mancktelow & Grasemann 1997; cf. also Wang & Zhou 2009). Depending on wavelength and amplitude (height), the underlying thermal structure can undulate with the topography, causing the distance between closure isotherm and surface to vary with sample location. For example, beneath valley floors (lowest elevation samples) the depth to the PAZ will be less than beneath topographic ridges, that is the geothermal gradients will vary with location. Failure to take this effect into account when constructing age–elevation plots to determine exhumation rates can lead to incorrect

interpretations. Detailed thermal modelling studies by Braun (2002) show how variable thermal structure can bias the interpretation of low-temperature thermochronological data.

*Landscape evolution modelling.* Much effort has been devoted recently to understand the coupling between tectonic and surface processes in the formation of recent topography (e.g. Braun 2002; Burbank & Anderson 2001; Burbank 2002; Ehlers & Farley 2003; Braun *et al.* 2006). Quantification of the rate at which landforms adapt to a changing tectonic, heat flow and climate environment (i.e. 'dynamic topography evolution') are performed by combining geomorphological analytical work, low-temperature thermochronological data and 3D thermokinematic modelling. Thermokinematic modelling with the 3D finite-element computer code Pecube (Braun 2003, 2005) predicts time–temperature ( $t$ – $T$ ) paths for all rock particles that, at the end of the computations, occupy the locations of the nodes at the surface of the finite-element mesh. From the  $t$ – $T$  paths, apparent AHe and AFT ages are generated by varying topography, erosion rates, uplift rates and heat flow values. Thus, Pecube allows an overall uplift rate to be created or a block of infinite space, which is bordered by normal faults and/or thrusts, to be defined. When movement is localized at the faults or thrusts, computer-code-generated age data are subsequently compared with the determined real thermochronological age data. As a result, it is now possible to match age data with geomorphological results and cosmogenic nuclide-based age dating in order to test landscape evolution models by processing different timescale resolutions.

### *Detrital thermochronology*

The 'standard' approach to deriving an AFT age is typically based on the measurement of the track density (age) for 20–30 (more if track densities are low) single-grain ages. Detrital thermochronology requires 50–120 grains per sample to enable statistical deconvolution into source-age components. Depending on the objectives of the study, up to 117 grains should be dated if the analyst wants to ensure that no fraction of the dated population comprising more than 0.05 of the total is missed at the 95% confidence level (Vermeesch 2004). Once the desired numbers of grains have been counted, the dataset can be divided into principal age components. Binomfit is a freeware program written by Mark Brandon that calculates ages and uncertainties for mixed distributions of fission-track grain ages. It uses an algorithm based on the decomposition method of Galbraith & Green (1990). Age components are compared with the

age patterns of the hinterland and correlated with specific source areas (e.g. Brandon 1996; Bernet & Spiegel 2004; Bernet & Garver 2005).


Detrital thermochronology is routinely used in *provenance analysis*, and for denudation and landscape evolution studies (e.g. Hurford & Carter 1991; Garver *et al.* 1999; Bernet *et al.* 2006). Specific tectonic and geomorphological applications include dating of hinterland uplift to reconstruct early exhumation rates in active orogenic belts, determining sediment-source regions, and reconstructing (palaeo-) drainage systems. A particularly useful interpretative method is to plot the *lag time* (or erosion–transport interval) between sediment deposition age and youngest detrital exhumation age. This provides key information on exhumation history, and conceptually it is possible to use the lag time to monitor the evolution of a mountain belt as it passes from growth stages (decreasing lag times) into topographic and exhumation steady state (constant lag time) into orogenic decay (increasing lag time). Other provenance applications include constraining minimum depositional ages of sediments and the correlation of stratigraphic horizons.

Most sediment is derived from the erosion of pre-existing rocks, and therefore detrital apatite and zircon grains may contain tracks that accumulated in the original source rock. Weathering and physical erosion do not affect the retention and stability of fission tracks (Gleadow & Lovering 1974), but the preservation of provenance-related tracks does depend on the temperature history experienced by eroding source regions and subsequently by the

sediment as the basin evolves (Fig. 4). Whereas zircons provide excellent provenance indicators (e.g. Hurford & Carter 1991; Garver & Brandon 1994; Garver *et al.* 1999), the use of apatite for source-rock information is restricted to shallow basins (typically less than *c.* 2 km of burial) and drainage systems (e.g. Corrigan & Crowley 1992; Lonergan & Johnson 1998; Malusà *et al.* 2009). Detrital analyses often combine thermochronological and geochronological data (U–Pb dating of zircon,  $^{40}\text{Ar}/^{39}\text{Ar}$  dating of white mica), heavy mineral assemblages, grain size/shapes (zircon typology), and isotope signatures of detrital minerals (e.g. Dunkl *et al.* 2001; Carter & Foster 2009). The amount of provenance information can be increased by using double or triple dating techniques that involve more than one dating method on the same mineral grain, for example combined U–Pb or  $^{40}\text{Ar}/^{39}\text{Ar}$  and AFT dating (Carter & Moss 1999; Carrapa *et al.* 2009) and/or AHe dating (Rahl *et al.* 2003). A detrital dating technique that utilizes petrographic information is fission-track analysis of single pebbles or pebble populations using conglomerates from foreland basin deposits (Spiegel *et al.* 2001; Dunkl *et al.* 2009).

## Applications

Application of low-temperature thermochronology is not confined to the obvious, immediate purpose of dating rock formations, as this is rarely possible due to thermal resetting. Much more powerful



**Fig. 4.** Cartoon to show the relationship between the progressive exhumation of a source region and deposition of the eroded exhumation record in an adjacent sedimentary basin (modified after Garver *et al.* 1999). The time slices ( $t_1$ – $t_3$ ) correspond to three progressive and continuous intervals of erosion and deposition. Subsequently, a ‘stratigraphy’ of apatite fission-track ages develops within the basin, which is the inverse of the source-region exhumation age trend. It should be noted that with progressive burial, sediments may be heated sufficiently to cause annealing of the inherited AFT provenance records. Rarely is such burial-related heating sufficient to reset fission tracks in detrital zircons and titanites. Also, note how topography may cause the upwarping of the lower temperature isotherms.

(and more widely used) are the applications that exploit thermal resetting to reconstruct rock exhumation histories. The **main fields of application of AFT and AHe** thermochronology include provenance studies, thermal history analysis of sedimentary basins, the evolution of orogenic mountain belts and applications in non-orogenic settings. Recent work in these areas has focused on the **coupling between climate and tectonics** (e.g. Koons 1989; Willet 1999; Beaumont *et al.* 2000), and thermochronological datasets have been used to evaluate the role of climatically driven erosion as a component of exhumation (Blythe & Kleinspehn 1998; Reiners & Brandon 2006).

Fission-track and (U–Th–Sm)/He data are often combined with other sources of data such as thermochronology and geochronological dating to constrain higher temperature histories and/or rock formation age (e.g.  $^{40}\text{Ar}/^{39}\text{Ar}$  and K–Ar: Daszinnies *et al.* 2009; De Grave *et al.* 2009), and terrestrial cosmogenic nuclide dating to compare with more recent erosion rates (e.g. Cockburn *et al.* 2000; Kuhlemann *et al.* 2009). In addition, interpretations increasingly integrate data from stratigraphic archives, geomorphology, structural geology, remote sensing, petrology, fluid inclusion analysis, vitrinite reflectance, clay mineralogy, conodont colour alteration, zircon typology and seismic data. These multi-method approaches strengthen the interpretation of AFT and AHe data, and help to extend geological histories through time.

### *Absolute dating*

Owing to fission-track annealing and He diffusion at relatively low temperatures, geological events can be dated in well-defined settings, when rocks very rapidly passed the PAZ and/or partial retention zone and resided at the surface or at a very shallow-crustal level thereafter. Rapid cooling to low temperatures mainly occurs subsequent to volcanic or hydrothermal activity (Duddy *et al.* 1998), dyke emplacement, faulting and friction or meteorite impacts (e.g. Miller & Wagner 1979). For these cases, thermochronological ages are more or less identical to those obtained by conventional radiometric techniques, and can be used directly as discrete time constraints. Fission-track analysis is also used as a conventional method for dating glasses (e.g. Fleischer & Price 1964; Bigazzi & De Michele 1996), and has been applied to date stone tools and fossils (e.g. Morwood *et al.* 1998). Moreover, both fission-track and (U–Th)/He techniques have been used successfully to date the formation of supergene minerals such as some phosphates or vanadates, hematite, goethite, limonite, manganese oxides and carbonate minerals (e.g. Bender 1973; Lippolt *et al.* 1995; Shuster *et al.* 2005; Boni *et al.* 2007; Copeland *et al.* 2007).

### *Denudation and long-term landscape evolution studies*

Denudation and long-term landscape evolution studies represent the most common and broadest field of applied low-temperature thermochronology. Studies range from compressional to extensional settings and ‘stable’ cratonic interiors.

*Orogenic belts.* Orogenic mountain ranges are characterized by substantial relief and immense uplift/denudation rates, resulting in large-scale advective transfer of heat and increased thermal gradient. The obvious correlation between exhumation and cooling predetermined this setting for an early application of thermochronological research, and expanded the scope of the method(s) from a purely ‘age determination’ approach to a unique thermotectonic tool. Wagner (1968) and Wagner & Reimer (1972) first used AFT data to provide estimates of the time and rates at which rocks approach the surface and cool as a result of ‘uplift and erosion’. At present, low-temperature thermochronology is the most efficient method to quantify denudation rates on geological timescales. Changes in erosion rates with time can be constrained using multiple chronometers with different closure temperatures on the same rock sample, or from the distribution of cooling ages from a single system along a vertical transect. Spatial–temporal patterns of thermochronometrically determined erosion rates help to constrain the flow of material through orogenic wedges, orogenic growth and decay cycles, palaeorelief, and relationships with structural, geomorphological or climatic features (cf. Reiners & Brandon 2006). Subsequent to the first apatite thermochronological works in the Alps (e.g. Schaer *et al.* 1975; Wagner *et al.* 1979; Grundmann & Morteau 1985; Hurford 1986; Hurford *et al.* 1991), most of the world’s young orogenic belts were studied (e.g. Parrish 1983; Seward & Tulloch 1991; Corrigan & Crowley 1992; Hendrix *et al.* 1994; Blythe *et al.* 1996; O’Sullivan & Currie 1996; Sorkhabi *et al.* 1996; Dunkl & Demény 1997; Kamp 1997; Sanders *et al.* 1999; Fayon *et al.* 2001; Spiegel *et al.* 2001; Glasmacher *et al.* 2002b; Reiners *et al.* 2002; Thomson 2002; Willet *et al.* 2003; Van der Beek *et al.* 2006; Gibson *et al.* 2007; Vincent *et al.* 2007; Glotzbach *et al.* 2008, 2009; del Río *et al.* 2009; Ruiz *et al.* 2009).

*Continental rifts and passive continental margins.* Continental rifts are elongated tectonic depressions that result from extension and crustal thinning caused either by a regional extensional stress field or in response to asthenospheric upwelling (cf. Olsen 1995; Ziegler & Cloething 2004). Continental separation and the onset of sea-floor spreading mark the transition from a continental rift into a passive

margin. Consequently, the general morphotectonic evolution and appearance of rifts and passive margins is similar. In detail, the style of lithospheric extension, geometry of rifting, rate and amount of extension are influenced by local crust rheology coupled to erosion, which in turn is influenced by pre-existing morphology, rift-related magmatism, local drainage and climate (e.g. Gilchrist & Summerfield 1990; Kooi & Beaumont 1994; Olsen 1995; O'Sullivan & Brown 1998; Ziegler & Cloething 2004). Deconvolving these different controls has been the focus of AFT and AHe studies. Apatite thermochronological studies have been reported from most of the world's continental-size rift structures and highly extended terranes (for an overview cf. Stockli 2005) including the West Antarctic Rift System (e.g. Gleadow *et al.* 1984; Fitzgerald & Gleadow 1988; Fitzgerald 1992, 1994; Foster & Gleadow 1992a; Balestrieri *et al.* 1994; Fitzgerald & Stump 1997; Lisker 2002; Fitzgerald *et al.* 2006), the East African Rift System (e.g. Kohn & Eyal 1981; Omar *et al.* 1989; Foster & Gleadow 1993, 1996; Van Der Beek *et al.* 1998; Feinstein *et al.* 1996; Kohn *et al.* 1997; Balestrieri *et al.* 2005; Spiegel *et al.* 2007), and the Basin and Range Province (e.g. Armstrong *et al.* 2003; House *et al.* 2003; Colgan *et al.* 2006), as well as many other continental extension zones (e.g. Rohrmann *et al.* 1994; Van der Beek *et al.* 1996; Lisker & Fachmann 2001; Emmel *et al.* 2009). Two studies on the influence of heat-flow variation across major rifts were published by Gallagher *et al.* (1994) and Lisker *et al.* (2003).

Gallagher & Brown (1997) demonstrated how AFT data are used to study the evolution of rift-margin topography (linked to isostatic-flexural responses to erosional unloading) and to test the different geomorphic models of passive margin evolution. Most passive rift margins of the major continents have been extensively studied by apatite thermochronology, with a wealth of papers detailing different aspects of their evolution. These studies include Moore *et al.* (1986), Brown *et al.* (1990, 2002), Dumitru (1991), Kalaswad *et al.* (1993), Gallagher *et al.* (1994), Gallagher & Brown (1999), Carter *et al.* (2000), Johnson & Gallagher (2000), O'Sullivan *et al.* (2000), Persano *et al.* (2002), Gunnell *et al.* (2003), Seward *et al.* (2004), Japsen *et al.* (2006), Raab *et al.* (2005), Emmel *et al.* (2006, 2007) and Kounov *et al.* (2009). Transform margins have seen comparatively little study, with published work confined to the Ghana margin (Clift *et al.* 1997; Bouillin *et al.* 1997; Bigot-Cormier *et al.* 2005).

**Cratons.** Cratonic interiors were traditionally considered as tectonically and thermally stable features. They are characterized by extensive surfaces of minimal relief that infer low rates of erosion.


However, thermochronological studies conducted during the last two decades on different cratons revealed that throughout the Phanerozoic many of these ancient terrains have experienced discrete episodes of kilometre-scale crustal erosion (e.g. Crowley *et al.* 1986; Wagner 1990; Noble *et al.* 1997; Harman *et al.* 1998; Cederbom *et al.* 2000; Bojar *et al.* 2002; Glasmacher *et al.* 2002a; Kohn *et al.* 2002; Belton *et al.* 2004; Lorencak *et al.* 2004; Soderlund *et al.* 2005; Flowers *et al.* 2006). On the other hand, cratonic areas often show apparent inconsistencies between AFT and AHe ages. These *crossover ages* are discussed controversially as either resulting from non-thermal radiation-enhanced annealing of fission tracks (e.g. Hendriks & Redfield 2005) or reflecting the dependency of He retention properties from the chemical composition of apatites (e.g. Green *et al.* 2006; Kohn *et al.* 2009).

### Basin analysis

Sedimentary basins are major archives of geological history related either directly to the evolution of the basin itself (e.g. subsidence history, tectonics, climate, sea-level change) and/ or indirectly to the geological history of the sediment-source regions (e.g. provenance information, palaeo-basement, geomorphology, tectonics). Traditionally, the burial and deposition history of basin sediments is determined by investigation of the sedimentology and deposition ages derived from biostratigraphy, combined with structural and geophysical datasets. However, establishing the timing of maximum depths of burial, and the timing and extent of uplifts, is often hard to recover owing to stratigraphic gaps, poor biostratigraphic control and sediment types. In this regard, AFT chronology has proven a valuable tool.

Thermochronology becomes most useful when a basin has undergone a period of uplift or inversion, during which a section of strata has been removed by erosion (e.g. Kamp & Green 1990; Bray *et al.* 1992; Green *et al.* 1995). Specific information gained from thermochronological data include estimates of maximum palaeotemperatures, calculations of palaeogeothermal gradients and palaeo-heat flow, style (fast/slow) and time of cooling from maximum palaeotemperatures, characterization of mechanisms of heating and cooling, quantification of missing sections, stratigraphic dating, sediment provenance, and evaluation of hydrocarbon maturation (Gleadow *et al.* 1983; Green *et al.* 1989a; Kamp & Green 1990; Duane & Brown 1991; Mitchell 1997; Logan & Duddy 1998).

Particularly useful in basin analysis is the integration of thermochronological data with maximum palaeotemperature indicators (Fig. 5), like vitrinite reflectance (VR) and illite crystallinity



**Fig. 5.** Schematic to show how the thermal history of a sedimentary basin can be recovered by integration of data from multiple palaeothermometers and chronometers. Illite data constrain the burial or heating phase of a basin's thermal history, VR records maximum temperature, and combined apatite thermochronology constrains the timing of cooling and subsequent denudation (modified after Pevear 1999).

(e.g. Bray *et al.* 1992; Pagel *et al.* 1997; Duddy *et al.* 1998; Mathiesen *et al.* 2000; Ventura *et al.* 2001; Arne *et al.* 2002; Osadetz *et al.* 2002; Tingate & Duddy 2002). VR is the measure of the coalification rank of organic matter, and it is mainly dependent on temperature and time (Burnham & Sweeney 1989). VR data provide a direct estimation of maximum palaeotemperatures across the same temperature range as annealing in fission tracks in apatite, which enables the thermal history modelling of joint VR and AFT datasets to provide a more robust constraint on temperature–time histories. Basin modelling using combined thermochronological and VR constraints has become a routine, and is a valuable tool in the hydrocarbon exploration industry (Gleadow *et al.* 1983; Green *et al.* 2002; Emmerich *et al.* 2005; Underdown *et al.* 2007). Also of economic relevance is the application of apatite thermochronology to the exploration of hydrothermal ore deposits (cf. McInnes *et al.* 2005).

### *Tectonic processes*

Thermochronological methods can be used to detect tectonic activities in two ways. In areas of substantial block uplift, thermochronological ages may be

disrupted across tectonic structures. Such offsets in the palaeo-isotherm/-depth stratigraphy can be used to determine relative uplift between different blocks and the amount of throw on bounding faults (Fitzgerald & Gleadow 1988, 1990; Dumitru 1991; Foster & Gleadow 1992a, b, 1996; Fitzgerald *et al.* 1993; O'Sullivan *et al.* 1995, 2000; Johnson 1997; Rahn *et al.* 1997; Wagner *et al.* 1997; Thomson 1998; Kohn *et al.* 1999; Redfield *et al.* 2007; Ventura *et al.* 2009; Xu *et al.* 2009). Moreover, the timing of tectonic activity may be determined by the direct dating of faults, lineaments or pseudotachylites (Harman *et al.* 1998; O'Sullivan *et al.* 1998; Raab *et al.* 2002, 2009; Zwingmann & Mancktelow 2004; Tagami 2005; Timar-Geng *et al.* 2009; Yamada *et al.* 2009).

### **Summary**

As can be seen from the papers contained in this Special Publication, apatite thermochronology has grown to become a reliable and routinely used method of helping in solving a diverse range of geological problems. Although mature, the method continues to undergo developments. Over the last decade there have been considerable improvements

in AFT methodology and data interpretation, but there is still scope for further advances. Future studies to refine models of track annealing in apatite may shift away from laboratory-based experimentation to well-constrained geological experiments (e.g. Spiegel *et al.* 2007) and molecular dynamic simulations (Rabone *et al.* 2008). Further work will also need to be carried out on understanding the controls on annealing in zircons and titanites, as well as the establishment of new fission-track mineral dating systems (e.g. monazite, merrillite: cf. Wagner & Van den Haute 1992; Gleadow *et al.* 2002).

We are very grateful to A. Carter and F. Stuart for constructive reviews of the manuscript.

## References

- AHNERT, F. 2003. *Einführung in die Geomorphologie*. Ullmer, Stuttgart.
- ARMSTRONG, P. A., EHLERS, T. A., CHAPMAN, D. S., FARLEY, K. A. & KAMP, P. J. J. 2003. Exhumation of the central Wasatch Mountains, Utah: 1. Patterns and timing of exhumation deduced from low-temperature thermochronology data. *Journal of Geophysical Research*, **108** (B3), 2172; doi:10.1029/2001JB001708.
- ARNE, D. C., GRIST, A. M., ZENTILLI, M., COLLINS, M., EMBRY, A. & GENTZIS, T. 2002. Cooling of the Sverdrup Basin during Tertiary basin inversion: implications for hydrocarbon exploration. *Basin Research*, **14**, 183–206.
- BALESTRIERI, M. L., BIGAZZI, G., GHEZZO, C. & LOMBARDO, B. 1994. Fission track dating of apatites from the Granite Harbour Intrusive Suite and uplift–denudation history of the Transantarctic Mountains in the area between the Mariner and David Glaciers (northern Victoria Land, Antarctica). *Terra Antarctica*, **1**, 82–87.
- BALESTRIERI, M. L., STUART, F. M., PERSANO, C., ABBATE, E. & BIGAZZI, G. 2005. Geomorphic development of the escarpment of the Eritrean margin, southern Red Sea from combined apatite fission-track and (U–Th)/He thermochronometry. *Earth and Planetary Science Letters*, **231**, 97–110.
- BARBARAND, J., CARTER, A., WOOD, I. & HURFORD, T. 2003. Compositional and structural control of fission-track annealing in apatite. *Chemical Geology*, **198**, 107–137.
- BAUMHAUER, H. A. 1894. *Die Resultate der Aetz-methode in der kristallographischen Forschung*. W. Engelmann, Leipzig.
- BEAUMONT, C., KOOI, H. & WILLETT, S. 2000. Coupled tectonic-surface process models with applications to rifted margins and collisional orogens. In: MA, S. (ed.) *Geomorphology and Global Tectonics*. Wiley, Chichester, 29–55.
- BELTON, D. X., BROWN, R. W., KOHN, B. P., FINK, D. & FARLEY, K. A. 2004. Quantitative resolution of the debate over antiquity of the central Australian landscape: implications for the tectonic and geomorphic stability of cratonic interiors. *Earth and Planetary Science Letters*, **219**, 21–34.
- BENDER, M. 1973. Helium–uranium dating of corals. *Geochimica Cosmochimica Acta*, **37**, 1229–1247.
- BERGER, G. W. & YORK, D. 1981. Geothermometry from dating experiments. *Geochimica et Cosmochimica Acta*, **45**, 795–811.
- BERNET, M. & GARVER, J. I. 2005. Fission-track analysis of detrital zircon. *Reviews in Mineralogy and Geochemistry*, **58**, 205–237.
- BERNET, M. & SPIEGEL, C. 2004. Introduction: detrital thermochronology. In: BERNET, M. & SPIEGEL, C. (eds) *Detrital Thermochronology – Provenance Analysis, Exhumation, and Landscape Evolution of Mountain Belts*. Geological Society of America, Special Paper, **378**, 1–6.
- BERNET, M., VAN DER BEEK, P., PIK, R., HUYGHE, P., MUGNIER, J.-L., LABRIN, E. & SZULC, A. 2006. Miocene to Recent exhumation of the central Himalaya determined from combined detrital zircon fission-track and U/Pb analysis of Siwalik sediments, western Nepal. *Basin Research*, **18**, 393–412.
- BIGAZZI, G. & DE MICHELE, V. 1996. New fission-track age determinations on impact glasses. *Meteoritics and Planetary Science*, **31**, 234–236.
- BIGOT-CORMIER, F., BASILE, C., POUPEAU, G., BOUILLIN, J.-P. & LABRIN, E. 2005. Denudation of the Côte d'Ivoire–Ghana transform continental margin from apatite fission tracks. *Terra Nova*, **17**, 189–195; doi: 110.1111/j.1365-3121.2005.00605.x.
- BLACKBURN, T. J., STOCKLI, D. F. & WALKER, D. 2007. Magnetite (U–Th)/He dating and its application to the geochronology of intermediate to mafic volcanic rocks. *Earth and Planetary Science Letters*, **259**, 360–371.
- BLYTHE, A. E. & KLEINSPEHN, K. L. 1998. Tectonically versus climatically driven Cenozoic exhumation of the Eurasian plate margin, Svalbard: fission track analyses. *Tectonics*, **17**, 621–639.
- BLYTHE, A. E., BIRD, J. M. & OMAR, G. I. 1996. Deformational history of the central Brooks Range, Alaska: results from fission-track and  $^{40}\text{Ar}/^{39}\text{Ar}$  analyses. *Tectonics*, **15**, 440–455.
- BOJAR, A.-V., FRITZ, H., KARGL, S. & UNZOG, W. 2002. Phanerozoic tectonothermal history of the Arabian Nubian shield in the Eastern Desert of Egypt: evidence from fission track and paleostress data. *Journal of African Earth Sciences*, **34**, 191–202.
- BONI, M., TERRACCIANO, R., EVANS, N. J., LAUKAMP, C., SCHNEIDER, J. & BECHSTÄDT, T. 2007. Genesis of vanadium ores in the Otavi Mountainland, Namibia. *Economic Geology*, **102**, 441–469.
- BOUILLIN, J. P., POUPEAU, G., LABRIN, E., BASILE, C., SABIL, N., MASCLE, J., MASCLE, G., GILLOT, F. & RIOU, L. 1997. Fission track study: heating and denudation of marginal ridge of the Ivory Coast–Ghana transform margin. *Geo-Marine Letters*, **17**, 55–61.
- BRANDON, M. T. 1996. Probability density plot for fission-track grain-age samples. *Radiation Measurements*, **26**, 663–676.
- BRAUN, J. 2002. Estimating exhumation rate and relief evolution by spectral analysis of age–elevation datasets: *Terra Nova*, **14**, 210–214.

- BRAUN, J. 2003. Pecube: A new finite element code to solve the heat transport equation in three dimensions in the Earth's crust including the effects of a time-varying, finite amplitude surface topography. *Computer Geosciences*, **29**, 787–794.
- BRAUN, J. 2005. Quantitative constraints on the rate of landform evolution derived from low-temperature thermochronology. *Reviews in Mineralogy and Geochemistry* **58**, 351–374.
- BRAUN, J., VAN DER BEEK, P. & BATT, G. 2006. *Quantitative Thermochronology: Numerical Methods for the Interpretation of Thermochronological Data*. Cambridge University Press, Cambridge.
- BRAY, R. J., GREEN, P. F. & DUDDY, I. R. 1992. Thermal history reconstruction using apatite fission track analysis and vitrinite reflectance; a case study from the UK East Midlands and southern North Sea. In: HARDMAN, R. F. P. (ed.) *Exploration Britain; Geological Insights for the Next Decade*. Geological Society, London, Special Publications, **67**, 2–25.
- BROWN, R., GALLAGHER, K. & DUANE, M. 1994. A quantitative assessment of the effects of magmatism on the thermal history of the Karoo sedimentary sequence. *Journal of African Earth Sciences*, **18**, 227–243.
- BROWN, R., SUMMERFIELD, M. A. & GLEADOW, A. J. W. 2002. Denudational history along a transect across the Drakensberg Escarpment of southern Africa derived from apatite fission track thermochronology. *Journal of Geophysical Research*, **107**, 2350; doi:2310.1029/2001JB000745.
- BROWN, R. W., RUST, D. J., SUMMERFIELD, M. A., GLEADOW, A. J. W. & DEWIT, M. C. J. 1990. An Early Cretaceous phase of accelerated erosion on the south-western margin of Africa: evidence from apatite fission track analysis & the offshore sedimentary record. *Nuclear Tracks Radiation Measurements*, **17**, 339–350.
- BROWN, R. W., SUMMERFIELD, M. A. & GLEADOW, A. J. W. 1994. Apatite fission track analysis: its potential for the estimation of denudation rates and the assessment of models of long term landscape development. In: KIRKBY, M. J. (ed.) *Process Models and Theoretical Geomorphology*. Wiley, Chichester, 23–53.
- BURBANK, D. W. 2002. Rates of erosion and their implications for exhumation. *Mineralogical Magazine*, **66**, 25–52.
- BURBANK, D. W. & ANDERSON, R. S. 2001. *Tectonic Geomorphology*. Blackwell, Oxford.
- BURNHAM, A. K. & SWEENEY, J. J. 1989. A chemical kinetic model of vitrinite maturation and reflectance. *Geochimica et Cosmochimica Acta*, **53**, 2649–2657.
- BURTNER, R. L., NIGRINI, A. & DONELICK, R. A. 1994. Thermochronology of lower Cretaceous source rocks in the Idaho–Wyoming thrust belt. *AAPG Bulletin*, **78**, 1613–1636.
- CARLSON, W. D., DONELICK, R. A. & KETCHAM, R. A. 1999. Variability of apatite fission-track annealing kinetics; I, Experimental results. *American Mineralogist*, **84**, 1213–1223.
- CARRAPA, B., DECELLES, P. G., REINERS, P. W., GEHRELS, G. E. & SUDO, M. 2009. Apatite triple dating and white mica  $^{40}\text{Ar}/^{39}\text{Ar}$  thermochronology of syntectonic detritus in the Central Andes: a multi-phase tectonothermal history. *Geology*, **37**, 407–410.
- CARTER, A. & FOSTER, G. L. 2009. Improving constraints on apatite provenance: Nd measurement on fission-track-dated grains. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 57–72.
- CARTER, A. & MOSS, S. J. 1999. Combined detrital-zircon fission-track and U–Pb dating: a new approach to understanding hinterland evolution. *Geology*, **27**, 235–238.
- CARTER, A., ROQUES, D. & BRISTOW, C. S. 2000. Denudation history of onshore central Vietnam; constraints on the Cenozoic evolution of the western margin of the South China Sea. *Tectonophysics*, **322**, 265–277.
- CEDERBOM, C., LARSON, S. A., TULLBORG, E.-L. & STIBERG, J.-P. 2000. Fission track thermochronology applied to Phanerozoic thermotectonic events in central and southern Sweden. *Tectonophysics*, **316**, 153–167.
- CLIFT, P. D., LORENZO, J., CARTER, A., HURFORD, A. J. & PARTY, O. L. S. 1997. Transform tectonics and thermal rejuvenation on the Cote d'Ivoire-Ghana margin, west Africa. *Journal of the Geological Society, London*, **154**, 483–489.
- COCKBURN, H. A. P., BROWN, R. W., SUMMERFIELD, M. A. & SEIDL, M. A. 2000. Quantifying passive margin denudation and landscape development using a combined fission-track thermochronology and cosmogenic isotope analysis approach. *Earth and Planetary Science Letters*, **179**, 429–435.
- COLGAN, J. P., DUMITRU, T. A., MCWILLIAMS, M. & MILLER, E. L. 2006. Timing of Cenozoic volcanism and Basin and Range extension in northwestern Nevada: new constraints from the northern Pine Forest Range. *Geological Society of America Bulletin*, **118**, 126–139.
- COPELAND, P., WATSON, E. B., URIZAR, S. C., PATTERSON, D. & LAPEN, T. J. 2007. Alpha thermochronology of carbonates. *Geochimica et Cosmochimica Acta*, **71**, 4488–4511.
- CORRIGAN, J. D. & CROWLEY, K. D. 1992. Unroofing of the Himalayas: a view from apatite fission-track analysis of Bengal Fan sediments. *Geophysical Research Letters*, **19**, 2345–2348.
- COYLE, D. A. & WAGNER, G. A. 1998. Positioning the titanite fission-track partial annealing zone. *Chemical Geology*, **149**, 117–125.
- CROWLEY, K., NAESER, C. & BABEL, C. 1986. Tectonic significance of Precambrian apatite fission-track ages from the midcontinent United States. *Earth and Planetary Science Letters*, **79**, 329–336.
- CROWLEY, K. D., CAMERON, M. & SCHAEFER, R. L. 1990. Annealing of etchable fission-track damage in F-, OH-, Cl- and Sr-apatite: 1. Systematics and preliminary interpretations. *Nuclear Tracks Radiation Measurements*, **17**, 409–410.
- CROWLEY, K. D., CAMERON, M. & SCHAEFER, R. L. 1991. Experimental studies of annealing of etched fission tracks in fluorapatite. *Geochimica et Cosmochimica Acta*, **55**, 1449–1465.
- DASZINNIES, M. C., JACOBS, J., WARTHO, J.-A. & GRANTHAM, G. H. 2009. Post Pan-African



- thermo-tectonic evolution of the north Mozambican basement and its implication for the Gondwana rifting. Inferences from  $^{40}\text{Ar}/^{39}\text{Ar}$  hornblende, biotite and titanite fission-track dating. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 261–286.
- DE GRAVE, J., BUSLOV, M. M., VAN DEN HAUTE, P., METCALF, J., DEHANDSCHUTTER, B. & MCWILLIAMS, M. O. 2009. Multi-method chronometry of the Teletskoye graben and its basement, Siberian Altai Mountains: new insights on its thermo-tectonic evolution. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 237–259.
- DEL RÍO, P., BARBERO, L. & STUART, F. M. 2009. Exhumation of the Sierra de Cameros (Iberian Range, Spain): constraints from low-temperature thermochronology. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 153–165.
- DOBSON, K. J., PERSANO, C. & STUART, F. M. 2009. Quantitative constraints on mid- to shallow-crustal processes using the zircon (U–Th)/He thermochronometer. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 47–56.
- DONELICK, R. A. 1991. Crystallographic orientation dependence of mean etchable fission-track length in apatite: an empirical model and experimental observations. *American Mineralogist*, **76**, 83–91.
- DONELICK, R. A., RODEN, M. K., MOEDERS, J. D., CARPENTER, B. S. & MILLER, D. S. 1990. Etchable length reduction of induced fission tracks in apatite at room temperature,  $\sim 23^\circ\text{C}$ , crystallographic orientation effects and 'initial' mean lengths. *Nuclear Tracks Radiation Measurement*, **17**, 261–265.
- DONELICK, R. A., O'SULLIVAN, P. B. & KETCHAM, R. A. 2005. Apatite fission-track analysis. *Reviews in Mineralogy and Geochemistry*, **58**, 49–94.
- DUANE, M. & BROWN, R. 1991. Tectonic brines and sedimentary basins: further application of fission-track analysis in understanding Karoo Basin evolution (South Africa). *Basin Research*, **3**, 187–195.
- DUDDY, I. R., GREEN, P. F., HEGARTY, K. A., BRAY, R. J. & O'BRIAN, G. W. 1998. Dating and duration of hot fluid flow events determined using AFTA and vitrinite reflectance-based thermal history reconstruction. In: PARSELL, J. (ed.) *Dating and Duration of Fluid Flow and Fluid–Rock Interaction*. Geological Society, London, Special Publications, **144**, 41–51.
- DUDDY, I. R., GREEN, P. F. & LASLETT, G. M. 1988. Thermal annealing of fission tracks in apatite, 3. Variable temperature behavior. *Chemical Geology (Isotope Geoscience Section)*, **73**, 25–38.
- DUMITRU, T. A. 1991. Major Quaternary uplift along the northernmost San Andreas Fault, King Range, north-western California. *Geology*, **19**, 526–529.
- DUMITRU, T. A., HILL, K. C. ET AL. 1991. Fission track thermochronology: application to continental rifting of south-eastern Australia. *The APEA Journal*, **31**, 131–142.
- DUNKL, I. & DEMÉNY, A. 1997. Exhumation of the Rechnitz Window at the border of the Eastern Alps and Pannonian Basin during Neogene extension. *Tectonophysics*, **272**, 197–211.
- DUNKL, I., DI GIULIO, A. & KUHLEMANN, J. 2001. Combination of single-grain fission-track chronology and morphological analysis of detrital zircon crystals in provenance studies – sources of the Macigno Formation (Apennines, Italy). *Journal of Sedimentary Research*, **71**, 516–525.
- DUNKL, I., FRISCH, W., KUHLEMANN, J. & BRÜGEL, A. 2009. Pebble population dating as an additional tool for provenance studies – examples from the Eastern Alps. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 125–140.
- EHLERS, T. A. & FARLEY, K. A. 2003. Apatite (U–Th)/He thermochronometry: methods and applications to problems in tectonic and surface processes. *Earth and Planetary Science Letters*, **206**, 1–14.
- EHLERS, T. A., CHAUDHRI, T. ET AL. 2005. Computational tools for low-temperature thermochronometer interpretation. *Reviews in Mineralogy and Geochemistry*, **58**, 589–622.
- EMMEL, B., GEIGER, M. & JACOBS, J. 2006. Detrital apatite fission-track ages in Middle Jurassic strata at the rifted margin of W Madagascar – indicator for a protracted resedimentation history. *Sedimentary Geology*, **186**, 27–38.
- EMMEL, B., JACOBS, J., CROWHURST, P. & DASZINNIES, M. C. 2007. Combined apatite fission-track and single grain apatite (U–Th)/He ages from basement rocks of central Dronning Maud Land (East Antarctica) – Possible identification of thermally overprinted crustal segments. *Earth and Planetary Science Letters*, **264**, 72–88.
- EMMEL, B., JACOBS, J. & DASZINNIES, M. C. 2009. Combined titanite and apatite fission-track data from Gjelsvikfjella, East Antarctica – another piece of a concealed intracontinental Permo-Triassic Gondwana rift basin? In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 317–330.
- EMMERICH, A., GLASMACHER, U. A., BAUER, F., BECHSTÄDT, T. & ZÜLKE, R. 2005. Meso-/Cenozoic basin and carbonate platform development in the SW-Dolomites unravelled by basin modelling and apatite FT analysis: Rosengarten and Latemar (Northern Italy). *Sedimentary Geology*, **175**, 415–438.
- ENGLAND, P. & MOLNAR, P. 1990. Surface uplift, uplift of rocks, and exhumation of rocks. *Geology*, **18**, 1173–1177.

- FARLEY, K. A. 2000. Helium diffusion from apatite: general behaviour as illustrated by Durango fluorapatite. *Journal of Geophysical Research*, **105**, 2903–2914.
- FARLEY, K. A. 2002. (U–Th)/He dating: techniques, calibrations, and applications. *Reviews in Mineralogy and Geochemistry*, **47**, 819–844.
- FARLEY, K. A., WOLF, R. A. & SILVER, L. T. 1996. The effects of long-alpha-stopping distances on (U–Th)/He ages. *Geochimica et Cosmochimica Acta*, **60**, 4223–4229.
- FAYON, A. K., WHITNEY, D. L., TEYSSIER, C., GARVER, J. I. & DILEK, Y. 2001. Effects of plate convergence obliquity on timing and mechanisms of exhumation of a mid-crustal terrain, the Central Anatolian Crystalline Complex. *Earth and Planetary Science Letters*, **192**, 191–205.
- FEINSTEIN, S., KOHN, B. P., STECKLER, M. S. & EYAL, M. 1996. Thermal history of the eastern margin of the Gulf of Suez, I. reconstruction from borehole temperature and organic maturity measurements. *Tectonophysics*, **266**, 203–220.
- FITZGERALD, P. G. 1992. The Transantarctic Mountains of southern Victoria Land: the application of apatite fission track analysis to a rift shoulder uplift. *Tectonics*, **11**, 634–662.
- FITZGERALD, P. G. 1994. Thermochronologic constraints on post-Paleozoic tectonic evolution of the central Transantarctic Mountains, Antarctica. *Tectonics*, **13**, 818–836.
- FITZGERALD, P. G. & GLEADOW, A. J. W. 1988. Fission-track geochronology, tectonics and structure of the Transantarctic Mountains in Northern Victoria Land, Antarctica. *Chemical Geology (Isotope Geoscience Section)*, **73**, 169–198.
- FITZGERALD, P. G. & GLEADOW, A. J. W. 1990. New approaches in fission track geochronology as a tectonic tool: examples from the Transantarctic Mountains. *Nuclear Tracks*, **17**, 351–357.
- FITZGERALD, P. G. & STUMP, E. 1997. Cretaceous and Cenozoic episodic denudation of the Transantarctic Mountains, Antarctica: new constraints from apatite fission track thermochronology in the Scott Glacier region. *Journal of Geophysical Research*, **102**, 7747–7766.
- FITZGERALD, P. G., BALDWIN, S. L., WEBB, L. E. & O'SULLIVAN, P. B. 2006. Interpretation of (U–Th)/He single grain ages from slowly cooled crustal terranes: a case study from the Transantarctic Mountains of southern Victoria Land. *Chemical Geology*, **225**, 91–120.
- FITZGERALD, P. G., SORKHABI, R. B., REDFIELD, T. F. & STUMP, E. 1995. Uplift and denudation of the central Alaska Range: a case study in the use of apatite fission track thermochronology to determine absolute uplift parameters. *Journal of Geophysical Research*, **100**, 20,175–20,191.
- FITZGERALD, P. G., STUMP, E. & REDFIELD, T. F. 1993. Late Cenozoic uplift of Denali and its relation to relative plate motion and fault morphology. *Science*, **259**, 497–499.
- FLEISCHER, R. L. & PRICE, P. B. 1964. Glass dating by fission fragment tracks. *Journal of Geophysical Research*, **69**, 331–339.
- FLEISCHER, R. L., PRICE, P. B. & WALKER, R. M. 1975. *Nuclear Tracks in Solids. Principles and Applications*. University of California Press, Berkeley, CA.
- FLOWERS, R. M., BOWRING, S. A. & REINERS, P. W. 2006. Low long-term erosion rates and extreme continental stability documented by ancient (U–Th)/He dates. *Geology*, **34**, 925–928.
- FLOWERS, R. M., KETCHAM, R. A., SHUSTER, D. L. & FARLEY, K. A. 2009. Apatite (U–Th)/He thermochronometry using a radiation damage accumulation and annealing model. *Geochimica et Cosmochimica Acta*, **73**(8), 2347–2365; doi:10.1016/j.gca.2009.01.015.
- FOSTER, D. A. & GLEADOW, A. J. W. 1992a. Reactivated tectonic boundaries and implications for the reconstruction of southeastern Australia and northern Victoria Land, Antarctica. *Geology*, **20**, 267–270.
- FOSTER, D. A. & GLEADOW, A. J. W. 1992b. The morphotectonic evolution of rift-margin mountains in central Kenya: constraints from apatite fission-track thermochronology. *Earth and Planetary Science Letters*, **113**, 157–171.
- FOSTER, D. A. & GLEADOW, A. J. W. 1993. Episodic denudation in East Africa: a legacy of intracontinental tectonism. *Geophysical Research Letters*, **20**, 2395–2398.
- FOSTER, D. A. & GLEADOW, A. J. W. 1996. Structural framework and denudation history of the flanks of the Kenya and Anza Rifts, East Africa. *Tectonics*, **15**, 258–271.
- GALBRAITH, R. F. 1981. On statistical models for fission track counts. *Mathematical Geology*, **13**, 471–478.
- GALBRAITH, R. F. 1988. Graphical display of estimates having differing standard errors. *Technometrics*, **30**, 271–281.
- GALBRAITH, R. F. 1990. The radial plot: graphical assessment of spread in ages. *Nuclear Tracks and Radiation Measurement*, **17**, 207–214.
- GALBRAITH, R. F. & GREEN, P. F. 1990. Estimating the component ages in a finite mixture. *Nuclear Tracks and Radiation Measurements*, **17**, 197–206.
- GALBRAITH, R. F. & LASLETT, G. M. 1993. Statistical models for mixed fission track ages. *Nuclear Tracks and Radiation Measurements*, **21**, 459–470.
- GALBRAITH, R. F., LASLETT, G. M., GREEN, P. F. & DUDDY, I. R. 1990. Apatite fission track analysis: geological thermal history analysis based on a three-dimensional random process of linear radiation damage. *Philosophical Transactions of the Royal Society, A* **332**, 419–438.
- GALLAGHER, K. 1995. Evolving temperature histories from apatite fission-track data. *Earth and Planetary Science Letters*, **136**, 421–435.
- GALLAGHER, K. & BROWN, R. 1997. The onshore record of passive margin evolution. *Journal of the Geological Society, London*, **154**, 451–457.
- GALLAGHER, K. & BROWN, R. 1999. The Mesozoic denudation history of the Atlantic margins of southern African and Southeast Brazil and the relationship to offshore sedimentation. In: CAMERON, N. R., BATE, R. H. & CLURE, V. S. (eds) *The Oil and Gas Habitats of the South Atlantic*. Geological Society, London, Special Publications, **153**, 41–53.
- GALLAGHER, K., BROWN, R. & JOHNSON, C. 1998. Fission track analysis and its applications to geological

- problems. *Annual Review of Earth and Planetary Science*, **26**, 519–572.
- GALLAGHER, K., HAWKESWORTH, C. J. & MANTOVANI, M. S. M. 1994. The denudation history of the onshore continental margin of SE Brazil inferred from apatite fission track data. *Journal of Geophysical Research*, **99**, 18,117–18,145.
- GALLAGHER, K., STEPHENSON, J., BROWN, R., HOLMES, C. & FITZGERALD, P. 2005. Low temperature thermochronology and modeling strategies for multiple samples 1: Vertical profiles. *Earth and Planetary Science Letters*, **237**, 193–208.
- GARVER, J. I. & BRANDON, M. T. 1994. Fission-track ages of detrital zircons from Cretaceous strata, southern British Columbia: implications for the Baja BC hypothesis. *Tectonics*, **13**, 401–420.
- GARVER, J. I., BRANDON, M. T., RODEN-TICE, M. & KAMP, P. J. J. 1999. Erosional denudation determined by fission-track ages of detrital apatite and zircon. In: RING, U., BRANDON, M. T., WILLETT, S. & LISTER, G. (eds) *Exhumation Processes: Normal Faulting, Ductile Flow, and Erosion*. Geological Society, London, Special Publications, **154**, 283–304.
- GENTNER, W., KLEINMANN, B. & WAGNER, G. A. 1967. New K–Ar- and fission track ages of impact glasses and tektites. *Earth and Planetary Science Letters*, **2**, 83–86.
- GIBSON, M., SINCLAIR, H. D., LYNN, G. J. & STUART, F. M. 2007. Late- to post-orogenic exhumation of the Central Pyrenees revealed through combined thermochronological data and modelling. *Basin Research*, **19**, 323–334.
- GILCHRIST, A. R. & SUMMERFIELD, M. A. 1990. Differentiated denudation and flexural isostasy in formation of rifted-margin upwarps. *Nature*, **346**, 739–742.
- GLASMACHER, U. A., MANN, U. & WAGNER, G. N. A. 2002a. Thermotectonic evolution of the Barrandian, Czech Republic, as revealed by apatite fission-track analysis. *Tectonophysics*, **359**, 381–402.
- GLASMACHER, U. A., WAGNER, G. A. & PUCHKOV, V. N. 2002b. Thermo-tectonic evolution of the western Fold-and-Thrust belt, southern Urals, Russia, as revealed by apatite fission-track data. *Tectonophysics*, **354**, 25–48.
- GLEADOW, A. J. W. & DUDDY, I. R. 1981. A natural long-term track annealing experiment for apatite. *Nuclear Tracks*, **5**, 169–174.
- GLEADOW, A. J. W. & FITZGERALD, P. G. 1987. Uplift history and structure of the Transantarctic Mountains: new evidence from fission track dating of basement apatites in the Dry Valleys area, southern Victoria Land. *Earth and Planetary Science Letters*, **82**, 1–14.
- GLEADOW, A. J. W. & LOVERING, J. F. 1974. Effect of weathering on fission track dating. *Earth and Planetary Science Letters*, **22**, 163–168.
- GLEADOW, A. J. W., BELTON, D. X., KOHN, B. P. & BROWN, R. W. 2002. Fission track dating of phosphate minerals and the thermochronology of apatite. In: KOHN, M. J., RAKOVAN, J. & HUGHES, J. M. (eds) *Phosphates; Geochemical, Geobiological, and Materials Importance*. Mineralogical Society of America and Geochemical Society, Washington, DC, 579–630.
- GLEADOW, A. J. W., DUDDY, I. R., GREEN, P. F. & HEGARTY, K. A. 1986a. Fission track lengths in the apatite annealing zone and the interpretation of mixed ages. *Earth and Planetary Science Letters*, **78**, 245–254.
- GLEADOW, A. J. W., DUDDY, I. R., GREEN, P. F. & LOVERING, J. 1986b. Confined fission track lengths in apatite: a diagnostic tool for thermal history analysis. *Contribution Mineralogy Petrology*, **94**, 405–415.
- GLEADOW, A. J. W., DUDDY, I. R. & LOVERING, J. F. 1983. Fission track analysis: a new tool for the evaluation of thermal histories and hydrocarbon potential. *The APEA Journal*, **23**, 93–102.
- GLEADOW, A. J. W., GLEADOW, S., BELTON, D. X., KOHN, B. P., KROCHMAL, M. & BROWN, R. 2009. Coincidence mapping – a key strategy for the automatic counting of fission tracks in natural minerals. In: LISKE, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 25–36.
- GLEADOW, A. J. W., MCKELVEY, B. C. & FERGUSON, K. U. 1984. Uplift history of the Transantarctic Mountains in the Dry Valleys area, southern Victoria Land, Antarctica, from apatite fission track ages. *New Zealand Journal of Geology and Geophysics*, **27**, 457–464.
- GLOTZBACH, C., REINECKER, J., DANISIK, M., RAHN, M., FRISCH, W. & SPIEGEL, C. 2008. Neogene exhumation history of the Mont Blanc massif, western Alps. *Tectonics*, **27**, TC4011; doi:10.1029/2008TC002257.
- GLOTZBACH, C., SPIEGEL, C., REINECKER, J., RAHN, M. & FRISCH, W. 2009. What perturbs isotherms? An assessment using fission-track thermochronology and thermal modelling along the Gotthard transect, Central Alps. In: LISKE, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 111–124.
- GREEN, P. F. 1981. ‘Track-in track’ length measurements in annealed apatites. *Nuclear Tracks*, **5**, 121–128.
- GREEN, P. F. 1985. Comparison of zeta calibration base-lines for fission-track dating of apatite, zircon and sphene. *Chemical Geology*, **58**, 1–22.
- GREEN, P. F. 1986. On the thermo-tectonic evolution of Northern England: evidence from fission track analysis. *Geological Magazine*, **123**, 493–506.
- GREEN, P. F. 1988. The relationship between track shortening and fission track age reduction in apatite: combined influences of inherent instability, annealing anisotropy, length bias and system calibration. *Earth and Planetary Science Letters*, **89**, 335–352.
- GREEN, P. F. & DURRANI, S. A. 1977. Annealing studies of tracks in crystals. *Nuclear Track Detection*, **1**, 33–39.
- GREEN, P. F., CROWHURST, P. V., DUDDY, I. R., JAPSEN, P. & HOLFORD, S. P. 2006. Conflicting (U–Th)/He and fission track ages in apatite: enhanced He retention, not anomalous annealing behaviour. *Earth and Planetary Science Letters*, **250**, 407–427.
- GREEN, P. F., DUDDY, I. R. & BRAY, R. J. 1995. Applications of thermal history reconstruction in inverted

- basins. In: BUCHANAN, J. G. & BUCHANAN, P. G. (eds) *Basin Inversion*. Geological Society, London, Special Publications, **88**, 149–165.
- GREEN, P. F., DUDDY, I. R., GLEADOW, A. J. W. & LOVERING, J. F. 1989a. Apatite fission track analysis as a paleotemperature indicator for hydrocarbon exploration. In: NAESER, N. D. & MCCULLOH, T. H. (eds) *Thermal History of Sedimentary Basins: Methods and Case Histories*. Springer, New York, 181–195.
- GREEN, P. F., DUDDY, I. R., LASLETT, G. M., HEGARTY, K. A., GLEADOW, A. J. W. & LOVERING, J. F. 1989b. Thermal annealing of fission tracks in apatite: 4 – Quantitative modelling techniques and extension to geological timescales. *Chemical Geology (Isotope Geoscience Section)*, **79**, 155–182.
- GREEN, P. F., DUDDY, I. R., GLEADOW, A. J. W., TINGATE, P. R. & LASLETT, G. M. 1985. Fission-track annealing in apatite; track length measurements and the form of the Arrhenius plot. *Nuclear Tracks*, **10**, 323–328.
- GREEN, P. F., DUDDY, I. R., GLEADOW, A. J. W., TINGATE, P. R. & LASLETT, G. M. 1986. Thermal annealing of fission tracks in apatite, 1. A qualitative description. *Chemical Geology (Isotope Geoscience Section)*, **59**, 237–253.
- GREEN, P. F., DUDDY, I. R. & HEGARTY, K. A. 2002. Quantifying exhumation from apatite fission-track analysis and vitrinite reflectance data: precision, accuracy and latest results from the Atlantic margin of NW Europe. In: DORÉ, A. G., CARTWRIGHT, J. A., STOKER, M. S., TURNER, J. P. & WHITE, N. (eds) *Exhumation of the North Atlantic Margin: Timing, Mechanisms and Implications for Petroleum Exploration*. Geological Society, London, Special Publications, **196**, 331–354.
- GRUNDMANN, G. & MORTEANI, G. 1985. The young uplift and thermal history of the Central Eastern Alps (Austria/Italy), evidence from apatite fission track ages. *Jahrbuch der Geologischen Bundesanstalt*, **128**, 197–216.
- GUNNELL, Y., GALLAGHER, K., CARTER, A., WIDDOWSON, M. & HURFORD, A. J. 2003. Denudation history of the continental margin of western peninsular India since the early Mesozoic – reconciling apatite fission-track data with geomorphology. *Earth and Planetary Science Letters*, **215**, 187–201.
- HARMAN, R., GALLAGHER, K., BROWN, R., RAZA, A. & BIZZI, L. 1998. Accelerated denudation and tectonic/geomorphic reactivation of the cratons of northeastern Brazil during the Late Cretaceous. *Journal of Geophysical Research B*, **103**, 27,091–27,105.
- HASEBE, N., BARBARAND, J., JARVIS, K., CARTER, A. & HURFORD, A. J. 2004. Apatite fission-track chronometry using laser ablation ICP-MS. *Chemical Geology*, **207**, 135–145.
- HASEBE, N., CARTER, A., HURFORD, A. J. & ARAI, S. 2009. The effect of chemical etching on LA-ICP-MS analysis in determining uranium concentration for fission-track chronometry. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 37–46.
- HENDRIKS, B. W. H. & REDFIELD, T. F. 2005. Apatite fission track and (U–Th)/He data from Fennoscandia: An example of underestimation of fission track annealing in apatite. *Earth and Planetary Science Letters*, **236**, 443–458.
- HENDRIX, M. S., DUMITRU, T. A. & GRAHAM, S. A. 1994. Late Oligocene–early Miocene unroofing in the Chinese Tian Shan: an early effect of the India–Asia collision. *Geology*, **22**, 487–490.
- HOURIGAN, J. K., REINERS, P. W. & BRANDON, M. T. 2005. U–Th zonation-dependent alpha-ejection in (U–Th)/He chronometry. *Geochimica et Cosmochimica Acta*, **69**, 3349–3365.
- HOUSE, M. A., KELLEY, S. A. & ROY, M. 2003. Refining the footwall cooling history of a rift flank uplift, Rio Grande rift, New Mexico. *Tectonics*, **22**, 1060; doi:10.1029/2002TC001418.
- HURFORD, A. J. 1986. Cooling and uplift patterns in the Lepontine Alps South Central Switzerland and an age of vertical movement on the Insubric fault line. *Contributions to Mineralogy and Petrology*, **92**, 413–427.
- HURFORD, A. J. 1990. Standardization of fission track dating calibration: Recommendation by the Fission Track Working Group of the I.U.G.S. Subcommittee on Geochronology. *Chemical Geology*, **80**, 171–178.
- HURFORD, A. J. & CARTER, A. 1991. The role of fission track dating in discrimination of provenance. In: MORTON, A. C., TODD, S. P. & HAUGHTON, P. D. W. (eds) *Developments in Sedimentary Provenance Studies*. Geological Society, London, Special Publications, **57**, 67–78.
- HURFORD, A. J. & GREEN, P. F. 1983. The zeta age calibration of fission-track dating. *Chemical Geology*, **1**, 285–317.
- HURFORD, A. J., FITCH, F. J. & CLARKE, A. 1984. Resolution of the age structure of the detrital zircon populations of two Lower Cretaceous sandstones from the Weald of England by fission track dating. *Geological Magazine*, **121**, 269–277.
- HURFORD, A. J., HUNZIKER, J. C. & STOCKHERT, B. 1991. Constraints on the late thermotectonic evolution of the western Alps: evidence for episodic rapid uplift. *Tectonics*, **10**, 758–769.
- ISSLER, D. R. 1996. Optimizing time step size for apatite fission track annealing models. *Computers and Geosciences*, **22**, 67–74.
- JAPSEN, P., BONOW, J. M., GREEN, P. F., CHALMERS, J. A. & LIDMAR-BERGSTROM, K. 2006. Elevated, passive continental margins: Long-term highs or Neogene uplifts? New evidence from West Greenland. *Earth and Planetary Science Letters*, **248**, 330–339.
- JOHNSON, C. 1997. Resolving denudational histories in orogenic belts with apatite fission-track thermochronology and structural data: an example from southern Spain. *Geology*, **25**, 623–626.
- JOHNSON, C. & GALLAGHER, K. 2000. A preliminary Mesozoic and Cenozoic denudation history of the North East Greenland onshore margin. *Global and Planetary Change*, **24**, 261–274.
- KALASWAD, S., RODEN, M. K., MILLER, D. S. & MORISAWA, M. 1993. Evolution of the continental margin of western India: new evidence from apatite fission-track dating. *Journal of Geology*, **101**, 667–673.

- KAMP, P. J. J. 1997. Paleogeothermal gradient and deformation style, Pacific front of the Southern Alps Orogen: constraints from fission track thermochronology. *Tectonophysics*, **271**, 37–58.
- KAMP, P. J. J. & GREEN, P. F. 1990. Thermal and tectonic history of selected Taranaki Basin (New Zealand) wells assessed by apatite fission track analysis. *AAPG Bulletin*, **74**, 1401–1419.
- KEEVIL, N. B. 1943. The distribution of helium in rocks, V. *American Journal of Science*, **241**, 277–306.
- KETCHAM, R. A. 2005. Forward and inverse modeling of low-temperature thermochronometry data. *Reviews in Mineralogy and Geochemistry*, **58**, 275–314.
- KETCHAM, R. A., CARTER, A., DONELICK, R. A., BARBARAND, J. & HURFORD, A. J. 2007. Improved modelling of fission-track annealing in apatite. *American Mineralogist*, **92**, 789–798.
- KETCHAM, R. A., DONELICK, R. A. & DONELICK, M. B. 2000. AFTSolve: a program for multi-kinetic modeling of apatite fission-track data. *Geological Materials Research*, **2**, 1–32.
- KOHN, B. P. & EYAL, M. 1981. History of uplift of the crystalline basement of Sinai and its relation to the opening of the Red Sea as revealed by fission track dating of apatites. *Earth and Planetary Science Letters*, **52**, 129–141.
- KOHN, B. P., FOSTER, D. A., STECKLER, M. S. & EYAL, M. 1997. Thermal history of the eastern Gulf of Suez, II. Reconstruction from apatite fission track and  $^{40}\text{Ar}/^{39}\text{Ar}$  K-feldspar measurements. *Tectonophysics*, **283**, 219–239.
- KOHN, B. P., GLEADOW, A. J. W., BROWN, R. W., GALLAGHER, K., O'SULLIVAN, P. B. & FOSTER, D. A. 2002. Shaping the Australian crust over the last 300 million years: insights from fission track thermotectonic imaging and denudation studies of key terranes. *Australian Journal of Earth Sciences*, **49**, 697–717.
- KOHN, B. P., GLEADOW, A. J. W. & COX, S. J. D. 1999. Denudation history of the Snowy Mountains; constraints from apatite fission track thermochronology. In: KOHN, B. P. & BISHOP, P. (eds) *Long-term Landscape Evolution of the Southeastern Australian Margin; Apatite Fission Track Thermochronology and Geomorphology*. Blackwell, Melbourne, 181–198.
- KOHN, B. P., LORENCAK, M., GLEADOW, A. J. W., KOHLMANN, F., RAZA, A., OSADETZ, K. G. & SORJONEN-WARD, P. 2009. A reappraisal of low-temperature thermochronology of the eastern Fennoscandia Shield and radiation-enhanced apatite fission-track annealing. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 193–216.
- KOOI, H. & BEAUMONT, C. 1994. Escarpment evolution on high-elevation rifted margins: insights derived from a surface processes model that combines diffusion, advection, and reaction. *Journal of Geophysical Research*, **99**, 12,191–12,209.
- KOONS, P. O. 1989. The topographic evolution of collisional mountain belts: a numerical look at the southern Alps, New Zealand. *American Journal of Science*, **289**, 1041–1069.
- KOUNOV, A., VIOLA, G., DE WIT, M. & ANDREOLI, M. 2009. Denudation along the Atlantic passive margin: new insights from apatite fission-track analysis on the western coast of South Africa. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 287–306.
- KUHLEMANN, J., KRUMREI, I., DANIŠÍK, M. & VAN DER BORG, K. 2009. Weathering of granite and granitic regolith in Corsica: short-term  $^{10}\text{Be}$  versus long-term thermochronological constraints. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 217–235.
- LASLETT, G. M., GALBRAITH, R. F. & GREEN, P. F. 1994. The analysis of projected track lengths. *Radiation Measurements*, **23**, 103–123.
- LASLETT, G. M., GLEADOW, A. J. W. & DUDDY, I. R. 1984. The relationship between fission track length and track density distributions. *Nuclear Tracks*, **9**, 29–38.
- LASLETT, G. M., GREEN, P. F., DUDDY, I. R. & GLEADOW, A. J. W. 1987. Thermal annealing of fission tracks in apatite, 2. A quantitative analysis. *Chemical Geology (Isotope Geoscience Section)*, **65**, 1–13.
- LASLETT, G. M., KENDALL, W. S., GLEADOW, A. J. W. & DUDDY, I. R. 1982. Bias in measurement of fission-track length distributions. *Nuclear Tracks*, **6**, 79–85.
- LIPPOLT, H. J., BOSCHMANN KÄTHLER, W. & ARNDT, H. 1982. Helium und Uran in Schwarzwälder Bleiglanzen, ein Datierungsversuch. *Oberrheinische Geologische Abhandlungen*, **31**, 31–46.
- LIPPOLT, H. J., WERNICKE, R. S. & BÄHR, R. 1995. Paragenetic specularite and adularia (Elba, Italy): Concordant (U + Th)–He and K–Ar ages. *Earth and Planetary Science Letters*, **132**, 43–51.
- LISKER, F. 2002. Review of fission track studies in northern Victoria Land – Passive margin evolution versus uplift of the Transantarctic Mountains. *Tectonophysics*, **349**, 57–73.
- LISKER, F. & FACHMANN, S. 2001. Phanerozoic history of the Mahanadi region, India. *Journal of Geophysical Research*, **B106**, 22,027–22,050.
- LISKER, F., BROWN, R. & FABEL, D. 2003. Denudational and thermal history along a transect across the Lambert Graben, northern Prince Charles Mountains, Antarctica derived from apatite fission track thermochronology. *Tectonics*, **22**, 1055; doi:10.1029/2002TC001477.
- LOGAN, P. & DUDDY, I. R. 1998. An investigation of the thermal history of the Ahnet–Reggane basins, central Algeria, the consequences for hydrocarbon generation accumulation. In: MACGREGOR, D. S., MOODY, R. T. J. & CLARK-LOWES, D. D. (eds) *Petroleum Geology of North Africa*. Geological Society, London, Special Publications, **132**, 131–155.
- LONERGAN, L. & JOHNSON, C. 1998. Reconstructing orogenic exhumation histories using synorogenic detrital zircons and apatites from the Betic Cordillera, SE Spain. *Basin Research*, **10**, 353–364.

- LORENCAK, M., KOHN, B. P., OSADETZ, K. G. & GLEADOW, A. J. W. 2004. Combined apatite fission track and (U–Th)/He thermochronometry in a slowly cooled terrane: results from a 3440-m-deep drill hole in the southern Canadian Shield. *Earth and Planetary Science Letters*, **227**, 87–104.
- MALUSA, M. G., ZATTIN, M., ANDÒ, S., GARZANTI, E. & VEZZOLI, G. 2009. Focused erosion in the Alps constrained by fission-track ages on detrital apatites. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 141–152.
- MANCKTELOW, N. S. & GRASEMANN, B. 1997. Time-dependent effects of heat advection and topography on cooling histories during erosion. *Tectonophysics*, **270**, 167–195.
- MATHIESEN, A., BIDSTRUP, T. & CHRISTIANSEN, F. G. 2000. Denudation and uplift history of the Jameson Land basin, East Greenland – constrained from maturity and apatite fission track data. *Global and Planetary Change*, **24**, 275–301.
- MCINNES, B. I. A., EVANS, N. J., FU, F. Q. & GARWIN, S. 2005. Application of thermochronology to hydrothermal ore deposits. *Reviews in Mineralogy and Geochemistry*, **58**, 467–498.
- MEESTERS, A. G. C. A. & DUNAI, T. J. 2002a. Solving the production-diffusion equation for finite diffusion domains of various shapes: Part I. Implications for low-temperature (U–Th)/He thermochronology. *Chemical Geology*, **186**, 333–344.
- MEESTERS, A. G. C. A. & DUNAI, T. J. 2002b. Solving the production-diffusion equation for finite diffusion domains of various shapes: Part II. Application to cases with [alpha]-ejection and nonhomogeneous distribution of the source. *Chemical Geology*, **186**, 347–363.
- MILLER, D. S. & WAGNER, G. A. 1979. Age and intensity of thermal events by fission track analysis; the Ries impact crater. *Earth and Planetary Science Letters*, **43**, 351–358.
- MITCHELL, M. M. 1997. Elevated mid-Cretaceous paleotemperatures in the western Otway Basin: consequences for hydrocarbon generation models. *The APPEA Journal*, **37**, 505–523.
- MOORE, M. E., GLEADOW, A. J. W. & LOVERING, J. F. 1986. Thermal evolution of rifted continental margins: new evidence from fission tracks in basement apatites from southeastern Australia. *Earth and Planetary Science Letters*, **78**, 255–270.
- MORWOOD, M. J., O'SULLIVAN, P. B., AZIZ, F. & RAZA, A. 1998. Fission-track ages of stone tools and fossils on the east Indonesian island of Flores. *Nature*, **392**, 173–176.
- MURRELL, G. R., SOBEL, E. R., CARRAPA, B. & ANDRIESEN, P. 2009. Calibration and comparison of etching techniques for apatite fission-track thermochronology. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 73–85.
- NAESER, C. W. 1967. The use of apatite and sphene for fission track age determinations. *Geological Society of America Bulletin*, **78**, 1523–1526.
- NAESER, C. W. 1979. Fission-track dating and geological annealing of fission tracks. In: JÄGER, E. & HUNZIKER, J. C. (eds) *Lectures in Isotope Geology*. Springer, New York, 154–169.
- NOBLE, W. D., FOSTER, D. A. & GLEADOW, A. J. W. 1997. The post-Pan-African thermal and extensional history of crystalline basement rocks in eastern Tanzania. *Tectonophysics*, **275**, 331–350.
- OLSEN, K. H. 1995. *Continental Rifts: Evolution, Structure, Tectonics*. Elsevier, New York.
- OMAR, G. I., STECKLER, M. S., BUCK, W. R. & KOHN, B. P. 1989. Fission track analysis of basement apatites at the western margin of the Gulf of Suez rift, Egypt: evidence for synchronicity of uplift and subsidence. *Earth and Planetary Science Letters*, **94**, 316–328.
- OSADETZ, K. G., KOHN, B. P., FEINSTEIN, S. & O'SULLIVAN, P. B. 2002. Thermal history of Canadian Williston basin from apatite fission-track thermochronology – implications for petroleum systems and geodynamic history. *Tectonophysics*, **349**, 221–249.
- O'SULLIVAN, P. B. & CURRIE, L. D. 1996. Thermotectonic history of Mt Logan, Yukon territory, Canada: implications of multiple episodes of middle to late Cenozoic denudation. *Earth and Planetary Science Letters*, **144**, 251–261.
- O'SULLIVAN, P. B. & PARRISH, R. R. 1995. The importance of apatite composition and single-grain ages when interpreting fission track data from plutonic rocks: a case study from the Coast Ranges, British Columbia. *Earth and Planetary Science Letters*, **132**, 213–224.
- O'SULLIVAN, P. B., KOHN, B. P., FOSTER, D. A. & GLEADOW, A. J. W. 1995. Fission track data from the Bathurst Batholith; evidence for rapid Mid-Cretaceous uplift and erosion within the eastern highlands of Australia. *Australian Journal of Earth Sciences*, **42**, 597–607.
- O'SULLIVAN, P. B., KOHN, B. P. & MITCHELL, M. M. 1998. Phanerozoic reactivation along a fundamental Proterozoic crustal fault, the Darling River Lineament, Australia; constraints from apatite fission track thermochronology. *Earth and Planetary Science Letters*, **164**, 451–465.
- O'SULLIVAN, P. B., MITCHELL, M. M., O'SULLIVAN, A. J., KOHN, B. P. & GLEADOW, A. J. W. 2000. Thermotectonic history of the Bassian Rise, Australia; implications for the breakup of eastern Gondwana along Australia's southeastern margins. *Earth and Planetary Science Letters*, **182**, 31–47.
- PAGEL, M., BRAUN, J. J., DISNAR, J. R., MARTINEZ, L., RENAC, C. & VASSEUR, G. 1997. Thermal history constraints from studies of organic matter, clay minerals, fluid inclusions, and apatite fission tracks at the Ardeche paleo-margin (BA1 drill hole, GPF Program), France. *Journal of Sedimentary Research, Section A: Sedimentary Petrology and Processes*, **67**, 235–245.
- PARRISH, R. R. 1983. Cenozoic thermal evolution and tectonics of the Coast Mountains of British Columbia. 1. Fission-track dating, apparent uplift rates, and patterns of uplift. *Tectonics*, **2**, 601–631.
- PAUL, T. A. & FITZGERALD, P. G. 1992. Transmission electron-microscopic investigation of fission tracks in fluorapatite. *American Mineralogist*, **77**, 336–344.

- PERSANO, C., STUART, F. M., BISHOP, P. & BARFOD, D. N. 2002. Apatite (U–Th)/He age constraints on the development of the Great Escarpment on the southeastern Australian passive margin. *Earth and Planetary Science Letters*, **200**, 79–90.
- PEVEAR, D. R. 1999. Illite and hydrocarbon exploration. *Proceedings of the National Academy of Science, USA*, **96**, 3440–3446.
- PRICE, P. B. & WALKER, R. M. 1963. Fossil tracks of charged particles in mica and the age of minerals. *Journal of Geophysical Research*, **68**, 4847–4862.
- RAAB, M. J., BROWN, R. W., GALLAGHER, K., CARTER, A. & WEBER, K. 2002. Late Cretaceous reactivation of major crustal shear zones in northern Namibia: constraints from apatite fission track analysis. *Tectonophysics*, **349**, 75–92.
- RAAB, M. J., BROWN, R. W., GALLAGHER, K., WEBER, K. & GLEADOW, A. J. W. 2005. Denudational and thermal history of the Early Cretaceous Brandberg and Okenyenya igneous complexes on Namibia's Atlantic passive margin. *Tectonics*, **24**, TC3006; doi:10.1029/2004TC001688.
- RABONE, J. A. L., CARTER, A., HURFORD, A. J. & DE LEEUW, N. H. 2008. Modelling the formation of fission tracks in apatite minerals using molecular dynamics simulations. *Physics and Chemistry of Minerals*, **35**(10), 583–596; 10.1007/s00269-008-0250-6.
- RAHL, J. M., REINERS, P. W., CAMPBELL, I. H., NICOLESCU, S. & ALLEN, C. M. 2003. Combined single-grain (U–Th)/He and U/Pb dating of detrital zircons from the Navajo Sandstone, Utah. *Geology*, **31**, 761–764.
- RAHN, M. K., HURFORD, A. J. & FREY, M. 1997. Rotation and exhumation of a thrust plane: apatite fission-track data from the Glarus thrust, Switzerland. *Geology*, **25**, 599–602.
- REDFIELD, T. F., SCHOLL, D. W., FITZGERALD, P. G. & BECK, M. E., JR. 2007. Escape tectonics and the extrusion of Alaska: Past, present, and future. *Geology*, **35**, 1039–1042.
- REINERS, P. W. & BRANDON, M. T. 2006. Using thermochronology to understand orogenic erosion. *Annual Review of Earth and Planetary Sciences*, **34**, 419–466.
- REINERS, P. W. & EHLERS, T. A. (eds) 2005. *Low-temperature Thermochronology: Techniques, Interpretations, and Applications*. Reviews in Mineralogy and Geochemistry, **58**. Mineralogical Society of America, Geochemical Society.
- REINERS, P. W., EHLERS, T. A., GARVER, J. I., MITCHELL, S. G., MONTGOMERY, D. R., VANCE, J. A. & NICOLESCU, S. 2002. Late Miocene exhumation and uplift of the Washington Cascades. *Geology*, **30**, 767–770.
- REINERS, P. W., EHLERS, T. A. & ZEITLER, P. K. 2005. Past, present, and future of thermochronology. *Reviews in Mineralogy and Geochemistry*, **58**, 1–18.
- REINERS, P. W., SPELL, T. L., NICOLESCU, S. & ZANETTI, K. A. 2004. Zircon (U–Th)/He thermochronometry: He diffusion and comparisons with  $^{40}\text{Ar}/^{39}\text{Ar}$  dating. *Geochimica et Cosmochimica Acta*, **68**, 1857–1887.
- RING, U., BRANDON, M. T., WILLETT, S. D. & LISTER, G. S. 1999. Exhumation processes. In: RING, U., BRANDON, M. T., WILLETT, S. D. & LISTER, G. S. (eds) *Exhumation Processes: Normal Faulting, Ductile Flow, and Erosion*. Geological Society, London, Special Publications, **154**, 1–27.
- ROHRMANN, M., VAN DER BEEK, P. & ANDRIESEN, P. 1994. Syn-rift thermal structure and post-rift evolution of the Oslo Rift (southeast Norway): New constraints from fission track thermochronology. *Earth and Planetary Science Letters*, **127**, 39–54.
- RUIZ, G. M. H., CARLOTTO, V., VAN HEININGEN, P. V. & ANDRIESEN, P. A. M. 2009. Steady-state exhumation pattern in the Central Andes – SE Peru. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 307–316.
- RUTHERFORD, E. 1905. Present problems in radioactivity. *Popular Science Monthly*, May, 1–34.
- SANDERS, C. A. E., ANDRIESEN, P. A. M. & CLOETINGH, S. A. P. L. 1999. Life cycle of the East Carpathian orogen: erosion history of a doubly vergent critical wedge assessed by fission track thermochronology. *Journal of Geophysical Research*, **104**, 29,095–29,112.
- SCHAEER, J. P., REIMER, G. M. & WAGNER, G. A. 1975. Actual and ancient uplift rate in the Gotthard region, Swiss Alps: a comparison between precise levelling and apatite fission track age. *Tectonophysics*, **29**, 293–300.
- SEWARD, D. & TULLOCH, A. J. 1991. Fission-track analysis of Tertiary uplift history of granitic basement in the Victoria Range, West Coast, New Zealand. *New Zealand Journal of Geology and Geophysics*, **34**, 115–120.
- SEWARD, D., GRUJIC, D. & SCHREURS, G. 2004. An insight into the breakup of Gondwana: identifying events through low-temperature thermochronology from the basement rocks of Madagascar. *Tectonics*, **23**, TC3007; doi:10.1029/2003TC001556.
- SHUSTER, D. L. & FARLEY, K. A. 2009. The influence of artificial radiation damage and thermal annealing on helium diffusion kinetics in apatite. *Geochimica et Cosmochimica Acta*, **73**(1), 183–196; doi:10.1016/j.gca.2008.10.013.
- SHUSTER, D. L., FLOWERS, R. M. & FARLEY, K. A. 2006. The influence of natural radiation damage on helium diffusion kinetics in apatite. *Earth and Planetary Science Letters*, **249**, 148–161.
- SHUSTER, D. L., VASCONCELOS, P. M., HEIM, J. A. & FARLEY, K. A. 2005. Weathering geochronology by (U–Th)/He dating of goethite. *Geochimica et Cosmochimica Acta*, **69**, 659–673.
- SILK, E. C. H. & BARNES, R. S. 1959. Examination of fission fragment tracks with an electron microscope. *Philosophical Magazine*, **4**, 970–972.
- SODERLUND, P., JUEZ-LARRE, J., PAGE, L. M. & DUNAI, T. J. 2005. Extending the time range of apatite (U–Th)/He thermochronometry in slowly cooled terranes: palaeozoic to Cenozoic exhumation history of southeast Sweden. *Earth and Planetary Science Letters*, **239**, 266–275.



- SORKHABI, R. B., STUMP, E., FOLAND, K. A. & JAIN, A. K. 1996. Fission-track and  $^{40}\text{Ar}/^{39}\text{Ar}$  evidence for episodic denudation of the Gangotri granites in the Garhwal Higher Himalaya, India. *Tectonophysics*, **260**, 187–199.
- SPIEGEL, C., KOHN, B., BELTON, D. X., BERNER, Z. & GLEADOW, A. J. W. 2009. Apatite (U–Th–Sm)/He thermochronology of rapidly cooled samples: the effect of He implantation. *Earth and Planetary Science Letters*, doi: 10.1016/j.epsl.2009.05.045.
- SPIEGEL, C., KOHN, B., RAZA, A., RAINER, T. & GLEADOW, A. 2007. The effect of long-term low-temperature exposure on apatite fission track stability: a natural annealing experiment in the deep ocean. *Geochimica et Cosmochimica Acta*, **71**, 4512–4537.
- SPIEGEL, C., KUHLEMANN, J., DUNKL, I. N. & FRISCH, W. 2001. Paleogeography and catchment evolution in a mobile orogenic belt: the Central Alps in Oligo-Miocene times. *Tectonophysics*, **341**, 33–47.
- STEPHENSON, J., GALLAGHER, K. & HOLMES, C. C. 2006. Low temperature thermochronology and strategies for multiple samples: 2: Partition modelling for 2D/3D distributions with discontinuities. *Earth and Planetary Science Letters*, **241**, 557–570.
- STOCKLI, D. F. 2005. Application of low-temperature thermochronometry to extensional tectonic settings. *Reviews in Mineralogy and Geochemistry*, **58**, 411–448.
- STÜWE, K., WHITE, L. & BROWN, R. 1994. The influence of eroding topography on steady-state isotherms. Application to fission track analysis. *Earth and Planetary Science Letters*, **124**, 63–74.
- SUMMERFIELD, M. A. 1991. *Global Geomorphology*. Longman, London.
- SUMMERFIELD, M. A. & BROWN, R. W. 1998. Geomorphic factors in the interpretation of fission-track data. In: VAN DEN HAUTE, P. & DE CORTE, F. (eds) *Advances in Fission-track Geochronology*. Kluwer Academic, Dordrecht, 269–284.
- TAGAMI, T. 2005. Zircon fission-track thermochronology and applications to fault studies. *Reviews in Mineralogy and Geochemistry*, **58**, 95–122.
- THOMSON, S. N. 1998. Assessing the nature of tectonic contacts using fission-track thermochronology: an example from the Calabrian Arc, southern Italy. *Terra Nova*, **10**, 32–36.
- THOMSON, S. N. 2002. Late Cenozoic geomorphic and tectonic evolution of the Patagonian Andes between latitudes 42°S and 76°S: an appraisal based on fission-track results from the transpressional intra-arc Liquine–Ofqui fault zone. *Geological Society of America Bulletin*, **114**, 1159–1173.
- TIMAR-GENG, Z., HENK, A. & WETZEL, A. 2009. Convective heat transfer in a steeply dipping fault zone and its impact on the interpretation of fission-track data – a modelling study. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 87–98.
- TINGATE, P. R. & DUDDY, I. R. 2002. The thermal history of the eastern Officer Basin (South Australia); evidence from apatite fission track analysis and organic maturity data. *Tectonophysics*, **349**, 251–275.
- UNDERDOWN, R., REDFERN, J. & LISKER, F. 2007. Constraining the burial history of the Ghadames Basin, North Africa: an integrated analysis using sonic velocities, vitrinite reflectance data and apatite fission track ages. *Basin Research*, **19**(4), 557–578; doi: 10.1111/j.1365-2117.2007.00335.x.
- VAN DER BEEK, P. A., DELVAUX, D., ANDRIESEN, P. A. M. & LEVI, K. G. 1996. Early Cretaceous denudation related to convergent tectonics in the Baikal region, SE Siberia. *Journal of the Geological Society, London*, **153**, 515–523.
- VAN DER BEEK, P., MBEDE, E., ANDRIESEN, P. & DELVAUX, D. 1998. Denudation history of the Malawi and Rukwa Rift flanks (East African Rift System) from apatite fission track thermochronology. *Journal of African Earth Sciences*, **26**, 363–385.
- VAN DER BEEK, P., ROBERT, X., MUGNIER, J.-L., BERNET, M., HUYGHE, P. & LABRIN, E. 2006. Late Miocene–Recent exhumation of the central Himalaya and recycling in the foreland basin assessed by apatite fission-track thermochronology of Siwalik sediments, Nepal. *Basin Research*, **18**, 413–434.
- VENTURA, B., LISKER, F. & KOPP, J. 2009. Thermal and denudation history of the Lusatian Block (NE Bohemian Massif, Germany) as indicated by apatite fission-track data. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 181–192.
- VENTURA, B., PINI, G. A. & ZUFFA, G. G. 2001. Thermal history and exhumation of the Northern Apennines (Italy): evidence from combined apatite fission track and vitrinite reflectance data from foreland basin sediments. *Basin Research*, **13**, 435–448.
- VERMEESCH, P. 2004. How many grains are needed for a provenance study? *Earth and Planetary Science Letters*, **224**, 441–451.
- VINCENT, S. J., MORTON, A. C., CARTER, A., GIBBS, S. & BARABADZE, T. G. 2007. Oligocene uplift of the Western Greater Caucasus: an effect of initial Arabia–Eurasia collision. *Terra Nova*, **19**, 160–166.
- WAGNER, G. A. 1966. Altersbestimmung an Tektiten und anderen natürlichen Gläsern mittels Spuren der spontanen Kernspaltung des Uran-238 (“fission track”-Methode). *Zeitschrift für Naturforschung*, **21a**, 733–745.
- WAGNER, G. A. 1968. Fission track dating of apatites. *Earth and Planetary Science Letters*, **4**, 411–415.
- WAGNER, G. A. 1972. The geological interpretation of fission track ages. *Transactions of the American Nuclear Society*, **15**, 117.
- WAGNER, G. A. 1979. Correction and interpretation of fission track ages. In: JÄGER, E. & HUNZIKER, J. C. (eds) *Lectures in Isotope Geology*. Springer, New York, 170–177.
- WAGNER, G. A. 1990. Apatite fission-track dating of the crystalline basement of Middle Europe: concepts and results. *Nuclear Tracks Radiation Measurements*, **17**, 277–282.
- WAGNER, G. A. & REIMER, G. M. 1972. Fission track tectonics: the tectonic interpretation of fission track apatite ages. *Earth and Planetary Science Letters*, **14**, 263–268.

- WAGNER, G. A. & VAN DEN HAUTE, P. 1992. *Fission-track Dating*. Ferdinand Enke, Stuttgart.
- WAGNER, G. A., COYLE, D. A. *ET AL.* 1997. Post-Variscan thermal and tectonic evolution of the KTB site and its surroundings. *Journal of Geophysical Research*, **102**, B8, 18,221–18,232.
- WAGNER, G. A., MILLER, D. S. & JÄGER, E. 1979. Fission track ages on apatite of Bergell rocks from central Alps and Bergell boulders in Oligocene sediments. *Earth and Planetary Science Letters*, **45**, 355–360.
- WANG, W. & ZHOU, Z. 2009. Reconstruction of paleotopography from low-temperature thermochronological data. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 99–110.
- WILLETT, S. D. 1992. Modelling thermal annealing of fission tracks in apatite. In: ZENTILLI, M. & REYNOLDS, P. H. (eds) *Short Course Handbook on Low Temperature Thermochronology*. Mineralogical Society of Canada, Toronto, 43–72.
- WILLETT, S. D. 1997. Inverse modeling of annealing of fission tracks in apatite. 1: A controlled random search method. *American Journal of Science*, **297**, 939–969.
- WILLETT, S. D. 1999. Orogeny and orography: the effects of erosion on the structure of mountain belts. *Journal of Geophysical Research*, **104**, 28,957–28,981.
- WILLETT, S. D., FISHER, D., FULLER, C., EN-CHAO, Y. & CHIA-YU, L. 2003. Erosion rates and orogenic-wedge kinematics in Taiwan inferred from fission-track thermochronometry. *Geology*, **31**, 945–948.
- WOLF, R. A., FARLEY, K. A. & KASS, D. M. 1998. Modeling of the temperature sensitivity of the apatite (U–Th)/He thermochronometer. *Chemical Geology*, **148**, 105–114.
- WOLF, R. A., FARLEY, K. A. & SILVER, L. T. 1996. Helium diffusion and low-temperature thermochronometry of apatite. *Geochimica et Cosmochimica Acta*, **60**, 4231–4240.
- XU, C., MANSY, J. L., VAN DEN HAUTE, P., GUILLOT, F., ZHOU, Z., CHEN, J. & DE GRAVE, J. 2009. Late- and post-Variscan evolution of the Ardennes in France and Belgium: constraints from apatite fission-track data. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 167–179.
- YAMADA, R., ONGIRAD, H., MATSUDA, T., OMURA, K., TAKEUCHI, A. & IWANO, H. 2009. Fission-track analysis of the Atotsugawa Fault (Hida Metamorphic Belt, central Japan): fault-related thermal anomaly and activation history. In: LISKER, F., VENTURA, B. & GLASMACHER, U. A. (eds) *Thermochronological Methods: From Palaeotemperature Constraints to Landscape Evolution Models*. Geological Society, London, Special Publications, **324**, 331–337.
- YAMADA, R., TAGAMI, T., NISHIMURA, S. & ITO, H. 1995. Annealing kinetics of fission tracks in zircon: an experimental study. *Chemical Geology (Isotope Geoscience Section)*, **122**, 249–258.
- ZAUN, P. E. & WAGNER, G. A. 1985. Fission track stability in zircons under geological conditions. *Nuclear Tracks and Radiation Measurements*, **10**, 303–307.
- ZEITLER, P. K., HERCZIG, A. L., MCDUGALL, I. & HONDA, M. 1987. U–Th–He dating of apatite: a potential thermochronometer. *Geochimica et Cosmochimica Acta*, **51**, 2865–2868.
- ZIEGLER, P. A. & CLOETHING, S. 2004. Dynamic processes controlling evolution of rifted basins. *Earth-Science Reviews*, **64**, 1–50.
- ZWINGMANN, H. & MANCKTELOW, N. 2004. Timing of Alpine fault gouges. *Earth and Planetary Science Letters*, **223**, 415–425.