

Contents

1	Spin vs Kramers restricted MP2	1
1.1	Spin-unrestricted non-relativistic MP2 energy	1
1.2	Spin-restricted non-relativistic MP2 energy	3
1.3	Kramers restricted relativistic MP2 energy	3
1.4	Reduction of the 16-block expansion	3
1.5	Complex (Re/Im) split: real-arithmetic form for closed-shell 1eX2C	4

Chapter 1

Spin vs Kramers restricted MP2

1.1 Spin-unrestricted non-relativistic MP2 energy

The non-relativistic MP2 energy in spin-orbitals, assuming Einstein summation, is

$$E_{\text{MP2}} = \frac{1}{4} \bar{t}_{ab}^{ij} \bar{g}_{ij}^{ab}. \quad (1.1)$$

Here, $\bar{g}_{ij}^{ab}$ denotes the antisymmetrized electron repulsion integrals,

$$\bar{g}_{ij}^{ab} = g_{ij}^{ab} - g_{ji}^{ab} = \langle ij|ab \rangle - \langle ji|ab \rangle = \langle ij||ab \rangle, \quad (1.2)$$

$$g_{ij}^{ab} = \langle ij|ab \rangle = \iint \phi_i^*(\mathbf{r1}) \phi_j^*(\mathbf{r2}) \frac{1}{|\mathbf{r1} - \mathbf{r2}|} \phi_a(\mathbf{r1}) \phi_b(\mathbf{r2}) d\mathbf{r1} d\mathbf{r2} \quad (1.3)$$

and

$$\bar{t}_{ab}^{ij} = \frac{\bar{g}_{ab}^{ij}}{\epsilon_i + \epsilon_j - \epsilon_a - \epsilon_b} = \frac{\bar{g}_{ab}^{ij}}{\Delta_{ab}^{ij}} \quad (1.4)$$

are the antisymmetrized MP2 amplitudes, where ϵ_p are the orbital energies. Indices i and j denote occupied orbitals, while indices a and b denote virtual orbitals. Note:

$$g_{ij}^{ab} = g_{ji}^{ba} = g_{ab}^{*ji} = g_{ba}^{*ji} \quad (1.5)$$

Before we look at spin-separated E_{MP2} , we need to realize that not all spin-combinations are non-zero. This can be understood by working out the spin-block separation of $\bar{g}_{ij}^{ab}$:

$$\begin{aligned} \bar{g}_{ij}^{ab} &= \bar{g}_{i\alpha j\alpha}^{a\alpha b\alpha} + \bar{g}_{i\beta j\beta}^{a\beta b\beta} + \bar{g}_{i\alpha j\beta}^{a\alpha b\beta} + \bar{g}_{i\beta j\alpha}^{a\beta b\alpha} + \bar{g}_{i\alpha j\beta}^{a\alpha b\beta} + \bar{g}_{i\alpha j\beta}^{a\beta b\alpha} \\ \bar{g}_{i\alpha j\alpha}^{a\alpha b\alpha} &= g_{i\alpha j\alpha}^{a\alpha b\alpha} - g_{j\alpha i\alpha}^{a\alpha b\alpha} \end{aligned} \quad (1.6)$$

$$\text{and } \bar{g}_{i\alpha j\beta}^{a\alpha b\beta} = g_{i\alpha j\beta}^{a\alpha b\beta} - 0 \quad (1.7)$$

$$(g_{j\beta i\alpha}^{a\alpha b\beta} \text{ is zero due to violation of column spin symmetry}) \quad (1.8)$$

Decomposing the energy expression into its individual spin components,

$$E_{\text{MP2}} = E_{\alpha\alpha}^{\alpha\alpha} + E_{\beta\beta}^{\beta\beta} + E_{\alpha\beta}^{\alpha\beta} + E_{\beta\alpha}^{\beta\alpha} + E_{\beta\alpha}^{\alpha\beta} + E_{\alpha\beta}^{\beta\alpha} \quad (1.9)$$

$$= \frac{1}{4} \left[\bar{t}_{a\alpha b\alpha}^{i\alpha j\alpha} \bar{g}_{i\alpha j\alpha}^{a\alpha b\alpha} + \bar{t}_{a\beta b\beta}^{i\beta j\beta} \bar{g}_{i\beta j\beta}^{a\beta b\beta} + \bar{t}_{a\alpha b\beta}^{i\alpha j\beta} \bar{g}_{i\alpha j\beta}^{a\alpha b\beta} + \bar{t}_{a\beta b\alpha}^{i\beta j\alpha} \bar{g}_{i\beta j\alpha}^{a\beta b\alpha} + \bar{t}_{a\beta b\alpha}^{i\alpha j\beta} \bar{g}_{i\alpha j\beta}^{a\beta b\alpha} + \bar{t}_{a\alpha b\beta}^{i\beta j\alpha} \bar{g}_{i\beta j\alpha}^{a\alpha b\beta} \right] \quad (1.10)$$

Note that $E_{\alpha\beta}^{\alpha\beta} = E_{\beta\alpha}^{\beta\alpha}$ in Eq. (1.35). This can be seen by considering the fourth term in Eq. (1.35), which, upon swapping the columns of both $\bar{t}$ and $\bar{g}$ and then swapping the dummy indices, becomes identical to the third term in Eq. (1.35). Similarly, it can also be seen that $E_{\beta\alpha}^{\alpha\beta} = E_{\alpha\beta}^{\beta\alpha}$. Consequently, Eq. (1.35) can be rewritten as

$$E_{\text{MP2}} = \frac{1}{4}\bar{t}_{a_\alpha b_\alpha}^{i_\alpha j_\alpha} \bar{g}_{i_\alpha j_\alpha}^{a_\alpha b_\alpha} + \frac{1}{4}\bar{t}_{a_\beta b_\beta}^{i_\beta j_\beta} \bar{g}_{i_\beta j_\beta}^{a_\beta b_\beta} + \frac{1}{2}\bar{t}_{a_\alpha b_\beta}^{i_\alpha j_\beta} \bar{g}_{i_\alpha j_\beta}^{a_\alpha b_\beta} + \frac{1}{2}\bar{t}_{a_\beta b_\alpha}^{i_\beta j_\alpha} \bar{g}_{i_\beta j_\alpha}^{a_\beta b_\alpha}. \quad (1.11)$$

Now consider the tensor network for $E_{\alpha\alpha}^{\alpha\alpha}$ only:

$$E_{\alpha\alpha}^{\alpha\alpha} = \frac{1}{4}\bar{t}_{a_\alpha b_\alpha}^{i_\alpha j_\alpha} \bar{g}_{i_\alpha j_\alpha}^{a_\alpha b_\alpha} \quad (1.12)$$

$$= \frac{1}{4}(t_{a_\alpha b_\alpha}^{i_\alpha j_\alpha} - t_{b_\alpha a_\alpha}^{i_\alpha j_\alpha})(g_{i_\alpha j_\alpha}^{a_\alpha b_\alpha} - g_{j_\alpha i_\alpha}^{a_\alpha b_\alpha}) \quad (1.13)$$

$$= \frac{1}{4}[t_{a_\alpha b_\alpha}^{i_\alpha j_\alpha} g_{i_\alpha j_\alpha}^{a_\alpha b_\alpha} - t_{a_\alpha b_\alpha}^{i_\alpha j_\alpha} g_{j_\alpha i_\alpha}^{a_\alpha b_\alpha} - t_{b_\alpha a_\alpha}^{i_\alpha j_\alpha} g_{i_\alpha j_\alpha}^{a_\alpha b_\alpha} + t_{b_\alpha a_\alpha}^{i_\alpha j_\alpha} g_{j_\alpha i_\alpha}^{a_\alpha b_\alpha}]. \quad (1.14)$$

The last term in Eq. (1.14) is equal to the first term, which can be seen by swapping the column indices of g and then exchanging the dummy indices a_α and b_α . An analogous relation holds between the second and third terms in Eq. (1.14). Therefore, the expression for $E_{\alpha\alpha}$ can be written in a compact form as

$$E_{\alpha\alpha}^{\alpha\alpha} = \frac{1}{2}[t_{a_\alpha b_\alpha}^{i_\alpha j_\alpha} g_{i_\alpha j_\alpha}^{a_\alpha b_\alpha} - t_{a_\alpha b_\alpha}^{i_\alpha j_\alpha} g_{j_\alpha i_\alpha}^{a_\alpha b_\alpha}] \quad (1.15)$$

$$= \frac{1}{2}t_{a_\alpha b_\alpha}^{i_\alpha j_\alpha} (g_{i_\alpha j_\alpha}^{a_\alpha b_\alpha} - g_{j_\alpha i_\alpha}^{a_\alpha b_\alpha}). \quad (1.16)$$

Similarly, the expression for $E_{\beta\beta}^{\beta\beta}$ can be written in compact form as

$$E_{\beta\beta}^{\beta\beta} = \frac{1}{2}t_{a_\beta b_\beta}^{i_\beta j_\beta} (g_{i_\beta j_\beta}^{a_\beta b_\beta} - g_{j_\beta i_\beta}^{a_\beta b_\beta}). \quad (1.17)$$

Now consider $2E_{\alpha\beta}^{\alpha\beta}$:

$$2E_{\alpha\beta}^{\alpha\beta} = \frac{1}{2}\bar{t}_{a_\alpha b_\beta}^{i_\alpha j_\beta} \bar{g}_{i_\alpha j_\beta}^{a_\alpha b_\beta} \quad (1.18)$$

$$= \frac{1}{2}(t_{a_\alpha b_\beta}^{i_\alpha j_\beta} - t_{b_\beta a_\alpha}^{i_\alpha j_\beta})(g_{i_\alpha j_\beta}^{a_\alpha b_\beta} - g_{j_\beta i_\alpha}^{a_\alpha b_\beta}). \quad (1.19)$$

The second term in each bracket vanishes because it violates column spin equivalence, and hence $2E_{\alpha\beta}^{\alpha\beta}$ reduces to

$$2E_{\alpha\beta}^{\alpha\beta} = \frac{1}{2}t_{a_\alpha b_\beta}^{i_\alpha j_\beta} g_{i_\alpha j_\beta}^{a_\alpha b_\beta}. \quad (1.20)$$

Similarly, for $2E_{\beta\alpha}^{\alpha\beta}$, the first terms in the antisymmetrization brackets vanish, and we are left with

$$2E_{\beta\alpha}^{\alpha\beta} = \frac{1}{2}t_{a_\alpha b_\beta}^{i_\alpha j_\beta} g_{i_\alpha j_\beta}^{a_\alpha b_\beta}. \quad (1.21)$$

Therefore, substituting the results of Eqs. (1.16), (1.17), (1.20) and (1.21) into Eq. (1.11), we get the expression for the total unrestricted MP2 energy as:

$$E_{\text{UMP2}} = \frac{1}{2}t_{a_\alpha b_\alpha}^{i_\alpha j_\alpha} (g_{i_\alpha j_\alpha}^{a_\alpha b_\alpha} - g_{j_\alpha i_\alpha}^{a_\alpha b_\alpha}) + \frac{1}{2}t_{a_\beta b_\beta}^{i_\beta j_\beta} (g_{i_\beta j_\beta}^{a_\beta b_\beta} - g_{j_\beta i_\beta}^{a_\beta b_\beta}) + t_{a_\alpha b_\beta}^{i_\alpha j_\beta} g_{i_\alpha j_\beta}^{a_\alpha b_\beta} \quad (1.22)$$

1.2 Spin-restricted non-relativistic MP2 energy

The spin-restricted expression for MP2 energy can be obtained from Eq. (1.22) by simply dropping the spin indices as

$$E_{\text{RMP2}} = \frac{1}{2} t_{ab}^{ij} (g_{ij}^{ab} - g_{ji}^{ab}) + \frac{1}{2} t_{ab}^{ij} (g_{ij}^{ab} - g_{ji}^{ab}) + t_{ab}^{ij} g_{ij}^{ab} \quad (1.23)$$

$$= t_{ab}^{ij} (2g_{ij}^{ab} - g_{ji}^{ab}) \quad (1.24)$$

1.3 Kramers restricted relativistic MP2 energy

If P denotes a spinor (Kramers up), then let $\bar{P}$ denote¹ the time reversed version (Kramers down) of P , i.e.,

$$|P\rangle = \mathcal{T}|\bar{P}\rangle; \quad \{|P\rangle\} \cup \{|\bar{P}\rangle\} = \{|p\rangle\} \quad (1.25)$$

where $\mathcal{T}$ is the time reversal operator:

$$\mathcal{T} = \begin{cases} i\sigma_y \hat{K}_0, & \text{2-component} \\ i\Sigma_y \hat{K}_0, & \text{4-component} \end{cases} \quad (1.26)$$

and $\hat{K}_0$ is the complex conjugation operator.

Due to the symmetry of the matrix elements under time reversal, the following relationships arise for electron repulsion integrals for closed shell systems:

$$g_{PQ}^{RS} = g_{\bar{P}\bar{Q}}^{*\bar{R}\bar{S}} = g_{\bar{P}\bar{S}}^{*\bar{R}Q} = g_{RQ}^{*P\bar{S}} \quad (1.27)$$

$$-g_{\bar{Q}\bar{R}}^{*SP} = -g_{RQ}^{*PS} = g_{QP}^{\bar{S}\bar{R}} = g_{PQ}^{R\bar{S}} = -g_{\bar{S}\bar{R}}^{*\bar{Q}\bar{P}} = -g_{\bar{R}\bar{S}}^{*\bar{P}\bar{Q}} = g_{\bar{S}\bar{P}}^{*Q\bar{R}} = g_{\bar{P}\bar{S}}^{*\bar{R}Q} \quad (1.28)$$

$$g_{\bar{R}\bar{S}}^{*PQ} = g_{PQ}^{\bar{R}\bar{S}} \quad (1.29)$$

$$g_{\bar{P}\bar{Q}}^{*R\bar{S}} = g_{PQ}^{\bar{R}\bar{S}} \quad (1.30)$$

Let's rewrite the MP2 energy expression, separating it into each Kramers pair permuted blocks,

$$E_{\text{MP2}} = \frac{1}{4} \left[\bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{AB} \right] \quad (1.31)$$

$$+ \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{\bar{A}\bar{B}} + \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{A\bar{B}} + \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{\bar{A}B} \quad (1.32)$$

$$+ \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{\bar{A}\bar{B}} + \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{A\bar{B}} + \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{\bar{A}B} + \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{A\bar{B}} + \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{\bar{A}\bar{B}} + \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{\bar{A}B} \quad (1.33)$$

$$+ \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{\bar{A}\bar{B}} + \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{A\bar{B}} + \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{\bar{A}B} + \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{\bar{A}\bar{B}} \quad (1.34)$$

$$+ \bar{t}_{AB}^{IJ} \bar{g}_{IJ}^{\bar{A}\bar{B}} \quad (1.35)$$

1.4 Reduction of the 16-block expansion

The 16-block expansion of Eq. (1.35) reduces to a much smaller, structurally compact form by exploiting two properties:

¹Note: $\bar{}$ over an index denotes time reversal and $\bar{}$ over an operator denotes antisymmetrization

1. **Time-reversal symmetry (TRS) folding.** For closed-shell Kramers-restricted references, summands related by full bar-flip $X \leftrightarrow \bar{X}$ on every index are complex conjugates of one another. Each pair collapses via $A + A^* = 2 \Re A$ (or $A - A^* = 2i \Im A$), so each conjugate-pair becomes a single $\Re$ - or $\Im$ -marked summand.
2. **Antisymmetrizer recombination.** Direct/exchange pairs of non-symmetric tensors produced by the trace merge back into single antisymmetric tensors $\bar{g}, \bar{t}$ over the Kramers-restricted index space.

The resulting six-term Kramers-restricted MP2 energy expression (I, J, A, B are Kramers-up dummy indices, $\bar{I}, \bar{A}, \dots$ are their Kramers-down partners; $\bar{g}, \bar{t}$ denote antisymmetric tensors; $\Re[\cdot]$ marks an evaluation-time real-part of a closed contraction) is:

$$\begin{aligned}
 E_{\text{MP2}}^{\text{KR}} = & - \Re \left[g_{I_1 I_2}^{A_1 A_2} \bar{t}_{\bar{A}_1 \bar{A}_2}^{I_2 I_1} \right] \\
 & + \Re \left[\bar{g}_{I_1 \bar{I}_1}^{A_1 \bar{A}_1} \bar{t}_{\bar{A}_1 \bar{A}_1}^{I_1 \bar{I}_1} \right] \\
 & + \Re \left[g_{I_1 I_2}^{\bar{A}_1 \bar{A}_2} \bar{t}_{\bar{A}_1 \bar{A}_2}^{I_1 I_2} \right] \\
 & + 2 \Re \left[g_{I_1 I_2}^{A_1 \bar{A}_1} t_{\bar{A}_1 \bar{A}_1}^{I_1 I_2} \right] \\
 & - 2 \Re \left[g_{\bar{I}_1 \bar{I}_2}^{A_1 \bar{A}_1} t_{\bar{A}_1 \bar{A}_1}^{I_2 \bar{I}_1} \right] \\
 & - 2 \Re \left[g_{I_1 \bar{I}_1}^{A_1 A_2} \bar{t}_{\bar{A}_2 \bar{A}_1}^{I_1 \bar{I}_1} \right].
 \end{aligned} \tag{1.36}$$

The structure makes the cost breakdown explicit: one doubly-antisymmetrized $\bar{g} \cdot \bar{t}$ term (line 2), three singly-antisymmetrized $g \cdot \bar{t}$ terms (lines 1, 3, 6), and two non-antisymmetrized $g \cdot t$ terms (lines 4, 5).

1.5 Complex (Re/Im) split: real-arithmetic form for closed-shell 1eX2C

For a closed-shell 1eX2C reference the spin-free Coulomb operator spin-traces each per-electron transition density to its *complex scalar* Pauli component, so every tensor in Eq. (1.36) is intrinsically complex (2-component); there is no quaternion structure to exploit and a 4-component (quaternion) decomposition would over-decompose the expression for zero benefit.

The *minimal* real-arithmetic form is therefore the 2-component *complex split*: rewrite each complex tensor as two real tensors

$$g = \Re g + i \Im g, \quad g^* = \Re g - i \Im g, \tag{1.37}$$

$$t = \Re t + i \Im t, \quad t^* = \Re t - i \Im t, \tag{1.38}$$

and propagate complex arithmetic symbolically. Each $\Re[g \cdot t]$ summand of Eq. (1.36) expands as

$$\Re[g \cdot t] = \Re g \cdot \Re t - \Im g \cdot \Im t, \tag{1.39}$$

which is a difference of two products of *real* tensors. The full six-term Kramers-restricted MP2

energy expression in the complex-split form (with $\Re g$, $\Im g$, $\Re t$, $\Im t$ all real) is:

$$\begin{aligned}
E_{\text{MP2}}^{\text{KR,c-split}} = & - \Re \left[\Re g_{I_1 I_2}^{A_1 A_2} \Re t_{A_1 A_2}^{I_2 I_1} - \Im g_{I_1 I_2}^{A_1 A_2} \Im t_{A_1 A_2}^{I_2 I_1} \right] \\
& + \Re \left[\Re g_{I_1 \bar{I}_1}^{A_1 \bar{A}_1} \Re t_{A_1 \bar{A}_1}^{I_1 \bar{I}_1} - \Im g_{I_1 \bar{I}_1}^{A_1 \bar{A}_1} \Im t_{A_1 \bar{A}_1}^{I_1 \bar{I}_1} \right] \\
& + \Re \left[\Re g_{I_1 I_2}^{\bar{A}_1 \bar{A}_2} \Re t_{\bar{A}_1 \bar{A}_2}^{I_1 I_2} - \Im g_{I_1 I_2}^{\bar{A}_1 \bar{A}_2} \Im t_{\bar{A}_1 \bar{A}_2}^{I_1 I_2} \right] \\
& + 2 \Re \left[\Re g_{I_1 I_2}^{A_1 \bar{A}_1} \Re t_{A_1 \bar{A}_1}^{I_1 I_2} - \Im g_{I_1 I_2}^{A_1 \bar{A}_1} \Im t_{A_1 \bar{A}_1}^{I_1 I_2} \right] \\
& - 2 \Re \left[\Re g_{\bar{I}_1 \bar{I}_2}^{A_1 \bar{A}_1} \Re t_{A_1 \bar{A}_1}^{\bar{I}_2 \bar{I}_1} - \Im g_{\bar{I}_1 \bar{I}_2}^{A_1 \bar{A}_1} \Im t_{A_1 \bar{A}_1}^{\bar{I}_2 \bar{I}_1} \right] \\
& - 2 \Re \left[\Re g_{I_1 \bar{I}_1}^{A_1 A_2} \Re t_{A_2 A_1}^{I_1 \bar{I}_1} - \Im g_{I_1 \bar{I}_1}^{A_1 A_2} \Im t_{A_2 A_1}^{I_1 \bar{I}_1} \right]. \tag{1.40}
\end{aligned}$$

The outer $\Re[\cdot]$ wrappers are now structurally redundant — every inner product is a product of real tensors and therefore real — but they are retained as evaluation-dispatch markers so the inner contractions can be performed as real (DGEMM) rather than complex (ZGEMM) products. This is the cost win of the complex split versus Eq. (1.36).