

Fundamentals

26 April 2026 22:45

[Fundamentals > Procedures > Liver Biopsy] Medium - VERIFIED

Q7. After a liver biopsy, what is the safest and most appropriate position to place the client in?

- A) High Fowler's position
- B) Prone position
- C) Supine position
- D) Right lateral position

Answer: D) Right lateral position

Explanation:

Correct - D: Liver on right side. Right lateral position applies pressure to biopsy site → prevents hemorrhage + bile leak. Keep 2-4 hours.

Incorrect - A/B/C: No pressure on liver, risk of bleeding increases.

Trick: Think "Press the side where organ is". Kidney biopsy = affected side down.

Source: NORCET-2023 | Potter Perry Fundamentals

Tags: #PYQ #HighYield #Fundamentals #Procedure #Priority

[Fundamentals > IM Injection > Pediatric] Easy - VERIFIED

Q8. Which of the following sites is the preferred location for administering an intramuscular (IM) injection in infants and young children?

- A) Vastus lateralis muscle (mid-thigh)
- B) Dorsogluteal muscle
- C) Deltoid muscle
- D) Ventrogluteal muscle

Answer: A) Vastus lateralis muscle (mid-thigh)

Explanation:

Correct - A: Vastus lateralis is largest muscle in infants, no major nerves/vessels. Safest site till child is walking.

Incorrect - B: Dorsogluteal contraindicated <3 years - sciatic nerve injury risk.

Incorrect - C: Deltoid underdeveloped, small volume only after 3+ years.

Incorrect - D: Ventrogluteal safe but used after child walks.

Trick: "Infant = Lateral thigh". "Adult = Ventrogluteal/Deltoid".

Source: NORCET-2023 | Wong's Pediatric

Tags: #PYQ #HighYield #Fundamentals #Pediatric #Procedure #Safety

[Fundamentals > Drug Administration > Transdermal] Medium - VERIFIED

Q11. Which of the following is TRUE regarding the optimal site for the application of a nitroglycerin patch?

- A) Apply over a bony prominence
 - B) Apply on a clean, hairless area near the heart
 - C) Apply directly over a moving joint
 - D) Apply on the distal lower extremities
- Answer: B) Apply on a clean, hairless area near the heart

Explanation:

Correct - B: Nitroglycerin patch needs flush skin contact. Upper chest/arm is hairless, vascular, close to heart = best absorption. Rotate sites daily to prevent skin irritation.

Incorrect - A/C: Bony areas/joints = poor adhesion + poor blood flow. Patch falls off.

Incorrect - D: Distal extremities have vasoconstriction, poor absorption.

Trick: "Patch on Chest" for cardiac drugs. Hair cut mat karo, shave nahi karna - irritation badhti.

Source: NORCET-2023 | Potter Perry Fundamentals

Tags: #PYQ #HighYield #Fundamentals #Pharma #DrugAdministration #Priority

[Fundamentals > Anatomy > Cardiovascular] High - VERIFIED

Q12. What is the correct sequence of venous blood flow from the external jugular vein back to the heart?

- A) External jugular vein -> Subclavian vein -> Brachiocephalic vein -> Superior Vena Cava -> Heart
- B) External jugular vein -> Internal jugular vein -> Heart
- C) External jugular vein -> Brachiocephalic vein -> Subclavian vein -> Heart
- D) External jugular vein -> Internal jugular vein -> Subclavian vein -> Heart

Answer: A) External jugular vein -> Subclavian vein -> Brachiocephalic vein -> Superior Vena Cava -> Heart

Explanation:

Correct - A: Ext Jugular → Subclavian. Subclavian + Int Jugular = Brachiocephalic. Right + Left Brachiocephalic = SVC → RA.

Incorrect - B/C/D: Ext Jugular never drains to Int Jugular. Order galat hai.

Trick: Mnemonic "External Subscribes to Bravo Company" = Ext

Jugular→Subclavian→Brachiocephalic→SVC.

Source: NORCET-2023 | Gray's Anatomy

Tags: #PYQ #HighYield #Fundamentals #Anatomy #NCLEX #Conceptual

[Fundamentals > CPR > Infant BLS] Medium - VERIFIED

Q13. During infant CPR, where exactly should chest compressions be applied?

- A) Upper third of the sternum
- B) Middle of the sternum
- C) Lower half of the sternum (just below the nipple line)
- D) Directly on the xiphoid process

Answer: C) Lower half of the sternum (just below the nipple line)

Explanation:

Correct - C: AHA 2020: Infant = 2 fingers or 2 thumb-encircling on lower 1/2 sternum, just below nipple line. Depth 1.5 inches/4cm. Rate 100-120/min.

Incorrect - A/B: Upper/middle = ineffective, heart lower me hota hai infant me.

Incorrect - D: Xiphoid = liver laceration + gastric rupture risk. NEVER compress xiphoid.

Trick: Adult = lower 1/2, Child = lower 1/2, Infant = lower 1/2. Sab same bas technique alag.

Source: NORCET-2023 | AHA BLS 2020

Tags: #PYQ #HighYield #Fundamentals #CPR #Pediatric #Emergency

[Fundamentals > Patient Safety > WHO Checklist] Easy - VERIFIED

Q14. Which of the following phases is NOT a part of the WHO Surgical Safety Checklist?

- A) Sign in
- B) Time out
- C) Time in
- D) Sign out

Answer: C) Time in

Explanation:

Correct - C: WHO has 3 phases only: 1) Sign In - before anesthesia, 2) Time Out - before incision, 3) Sign Out - before leaving OR. "Time in" fake term hai.

Incorrect - A/B/D: Ye teen hi official phases hain.

Trick: "In-Out-In-Out" - Sign In, Time Out, Sign Out. Beech me "Time In" nahi aata.

Source: NORCET-2023 | WHO Guidelines 2009

Tags: #PYQ #HighYield #Fundamentals #PatientSafety #OT #GK

[Fundamentals > Hygiene > Perineal Care] Easy - VERIFIED

Q16. What is the standard and correct technique for cleaning the female perineal area to prevent infection?

- A) Back to front
- B) Front to back

C) Inner to outer in circles

D) Side to side

Answer: B) Front to back

Explanation:

Correct - B: Front to back = urethra to anus. Prevents E.coli from bowel to contaminate urethra/vagina. UTI prevention ka #1 rule.

Incorrect - A: Back to front = 100% UTI guarantee. Fecal matter urethra me jayega.

Incorrect - C: Circles for surgical site, not perineal.

Trick: "Front door Clean, Back door Dirty" - Clean se dirty ki taraf jao.

Source: NORCET-2023 | Potter Perry Fundamentals

Tags: #PYQ #HighYield #Fundamentals #Hygiene #InfectionControl #Priority

[Fundamentals > Infection Control > Biohazard] Medium - VERIFIED

Q17. Which solution concentration is standardly recommended for cleaning and disinfecting a small biological blood spill (e.g., 5 mL) on a hospital floor?

A) 5% Sodium hypochlorite

B) 1% Sodium hypochlorite

C) 2% Glutaraldehyde

D) 70% Isopropyl alcohol

Answer: B) 1% Sodium hypochlorite

Explanation:

Correct - B: CDC/WHO: Small blood spill <10ml = 1% sodium hypochlorite for 10 min contact. Large spill >10ml = 10% i.e. 1:10 dilution. 1% = 10,000 ppm.

Incorrect - A: 5% too strong, corrosive to floor, toxic fumes.

Incorrect - C: Glutaraldehyde for instruments, not floor.

Incorrect - D: Alcohol evaporates fast, not for blood. Blood inactivates alcohol.

Trick: Small spill = 1%, Large spill = 10%. "1 for small, 10 for big".

Source: NORCET-2023 | CDC Biohazard Guidelines

Tags: #PYQ #HighYield #Fundamentals #InfectionControl #Biohazard #Safety

[Fundamentals > Assessment > Hearing Test] Easy - VERIFIED

Q20. What frequency of tuning fork is standardly used by healthcare providers for auditory testing (such as the Rinne and Weber tests)?

A) 128 Hz

B) 256 Hz

C) 512 Hz

D) 1024 Hz

Answer: C) 512 Hz

Explanation:

Correct - C: 512 Hz = best for hearing tests. Optimal tone + vibration decay. Rinne/Weber dono me use hota.

Incorrect - A: 128 Hz = vibration sense testing, not hearing. For neuropathy check.

Incorrect - B/D: 256/1024 not standard. 512 Hz international standard.

Trick: "512 for Hearing, 128 for Feeling". 5-Hearing, 1-Feeling.

Source: NORCET-2023 | Bates Physical Exam

Tags: #PYQ #HighYield #Fundamentals #Assessment #ENT #Procedure

[Fundamentals > Infection Control > Hand Hygiene] Easy - VERIFIED

Q22. Which of the following is NOT one of the "5 Moments for Hand Hygiene" as defined by the WHO?

A) Before touching a patient

B) Before touching a patient's surroundings

C) After a procedure or body fluid exposure risk

D) After touching a patient

Answer: B) Before touching a patient's surroundings

Explanation:

Correct - B: WHO 5 Moments: 1) Before touching patient, 2) Before clean/aseptic procedure, 3) After body fluid exposure, 4) After touching patient, 5) AFTER touching patient surroundings. "Before surroundings" nahi hai.

Incorrect - A/C/D: Ye teeno official moments hain.

Trick: Surroundings ko chhune se PEHLE nahi, BAAD me hand wash. Environment dirty mana jata hai.

Source: NORCET-2023 | WHO Guidelines

Tags: #PYQ #HighYield #Fundamentals #InfectionControl #WHO #PatientSafety

[Fundamentals > Poisoning > Hydrocarbon] High - CORRECTED

Q24. Gastric lavage is considered strictly contraindicated in which of the following types of poisoning cases?

A) Organophosphate poisoning

B) Kerosene (hydrocarbon) poisoning

C) Paracetamol poisoning

D) Barbiturate overdose

Answer: B) Kerosene (hydrocarbon) poisoning

Explanation:

Correct - B: Hydrocarbons like kerosene have low viscosity + high volatility. Lavage/induced vomiting → aspiration → fatal chemical pneumonitis. Absolute contraindication.

Incorrect - A/C/D: Organophosphate, Paracetamol, Barbiturate me lavage kar sakte if <1 hour of ingestion + airway protected. Atropine + PAM for OP.

Trick: "Kerosene me Kucch nahi karna" - No lavage, No charcoal, No vomiting. Only supportive NPO + O2.

Source: NORCET-2023 | CMC Vellore Toxicology

Tags: #PYQ #HighYield #Fundamentals #Poisoning #Emergency #Contraindication #TrickQ

[Fundamentals > IV Fluids > Osmolarity] Medium - VERIFIED

Q25. Which type of intravenous solution is utilized to draw fluid out of the intracellular and interstitial tissue spaces and shift it into the intravascular blood stream?

A) Hypotonic solution

B) Isotonic solution

C) Hypertonic solution

D) Colloid-free solution

Answer: C) Hypertonic solution

Explanation:

Correct - C: Hypertonic = >375 mOsm/L. Ex: 3% NS, D10W, D5NS. High osmolarity pulls water from cells to vessels via osmosis. Used in cerebral edema, hyponatremia.

Incorrect - A: Hypotonic = <250 mOsm/L. Pushes fluid INTO cells. Used in dehydration. Ex: 0.45% NS.

Incorrect - B: Isotonic = 275-295 mOsm/L. No fluid shift. Ex: 0.9% NS, RL. For volume replacement.

Trick: "Hyper = High = Pulls water OUT of cells". "Hypo = Low = Pushes water IN".

Source: NORCET-2023 | Potter Perry Fundamentals

Tags: #PYQ #HighYield #Fundamentals #IVFluids #Pharma #Conceptual #NCLEX

[Fundamentals > Procedures > Lumbar Puncture] Medium - VERIFIED

Q31. Which position is recommended for a patient immediately following a lumbar puncture procedure?

A) High Fowler's position

B) Prone position

C) Supine position

D) Right lateral position

Answer: C) Supine position

Explanation:

Correct - C: Flat supine 4-12 hours. Prevents CSF leak from dural puncture → prevents post-LP headache. Headache mechanism: ↓CSF pressure → brain sagging → meningeal traction.
Incorrect - A: Fowler's = ↑hydrostatic pressure → more CSF leak → severe headache.
Incorrect - B/D: Prone/Right lateral doesn't prevent leak.
Trick: "LP ke baad Lie Flat" - L for Lumbar, L for Lie. Also increase oral fluids + caffeine.
Source: NORCET-2023 | Brunner & Suddarth
Tags: #PYQ #HighYield #Fundamentals #Procedure #Neuro #PatientSafety

[Fundamentals > CPR > Two Rescuer] Medium - VERIFIED

Q36. In two-rescuer CPR, after how many cycles of compressions and ventilations should the rescuers switch roles to prevent fatigue?

- A) 2 cycles
- B) 3 cycles
- C) 5 cycles
- D) 10 cycles

Answer: C) 5 cycles

Explanation:

Correct - C: AHA 2020: Switch compressor every 5 cycles OR 2 minutes, whichever comes first. Prevents fatigue → maintains depth 2-2.4 inches + rate 100-120.

Incorrect - A/B: Too frequent switching wastes time.

Incorrect - D: 10 cycles = 4+ min = fatigue → shallow compressions.

Trick: "5 cycles = 2 minutes = Switch". 30:2 ratio x 5 = 150 compressions then switch.

Source: NORCET-2023 | AHA BLS 2020

Tags: #PYQ #HighYield #Fundamentals #CPR #Emergency #AHA

[Fundamentals > Hemodynamic > CVP] High - VERIFIED

Q45. A patient is receiving a continuous intravenous Normal Saline infusion. The nurse assesses a Central Venous Pressure (CVP) of 18 mmHg. What is the priority nursing action?

- A) Increase the infusion rate
- B) Administer a continuous diuretic drip
- C) Stop the IV infusion immediately
- D) Lower the head of the bed

Answer: C) Stop the IV infusion immediately

Explanation:

Correct - C: Normal CVP = 2-8 mmHg. 18 mmHg = Severe fluid overload/CHF/RV failure. Priority: STOP IV to prevent pulmonary edema. Then notify doctor, sit patient up, O2, diuretics.

Incorrect - A: Increasing fluid = kill patient. Fluid overload worse.

Incorrect - B: Diuretic after stopping IV, not while infusing.

Incorrect - D: Lowering HOB = worse dyspnea in fluid overload.

Trick: "High CVP = Hold IV". Low CVP <2 = give fluid. CVP reflects RA pressure.

Source: NORCET-2023 | Lewis Med-Surg

Tags: #PYQ #HighYield #Fundamentals #Hemodynamic #Emergency #Priority #NCLEX

[Fundamentals > Dosage Calculation > IV Flow Rate] Medium - VERIFIED

Q50. Calculate the IV flow rate required to infuse 500 mL of 5% Dextrose over 5 hours using a microdrip set (drop factor = 60 drops/mL).

- A) 25 drops/min
- B) 60 drops/min
- C) 100 drops/min
- D) 125 drops/min

Answer: C) 100 drops/min

Explanation:

Correct - C: Formula: (Volume x Drop Factor) / Time in min = (500 x 60) / 300 = 100 gtt/min.

Microdrip Trick: mL/hr = gtt/min always. 500mL/5hr = 100mL/hr = 100 gtt/min.

Incorrect - A/B/D: Calculation error.

Trick: "Microdrip Magic: $\text{mL/hr} = \text{gtt/min}$ ". Macrodrip 15gtt: $(\text{mL/hr})/4$. Macrodrip 20gtt: $(\text{mL/hr})/3$.

Source: NORCET-2023 | Drug Calculation Book

Tags: #PYQ #HighYield #Fundamentals #Calculation #IVFluids #Formula #Math

[Fundamentals > Bandaging > Techniques] Easy - VERIFIED

Q2. Which bandage technique is most commonly used to wrap joints like the knee, ankle, or elbow?

- A) Spiral
- B) Circular
- C) Capeline
- D) Figure of 8

Answer: D) Figure of 8

Explanation:

Correct - D: Figure of 8 = crosses over joint, allows flexion/extension + provides compression. For knee, ankle, elbow, wrist.

Incorrect - A: Spiral = for uniform cylindrical parts like arm, thigh. Slips on joints.

Incorrect - B: Circular = for anchoring, not joints.

Incorrect - C: Capeline = for head/stump amputation.

Trick: "Joint = 8". 8 ka shape joint pe fit baithta hai.

Source: NORCET AIIMS 2021 | Potter Perry

Tags: #PYQ #HighYield #Fundamentals #Bandaging #Procedure

[Fundamentals > IM Injection > Sites] High - VERIFIED

Q3. Which of the following sites is strictly avoided for intramuscular (IM) injections in adults due to a high risk of nerve injury?

- A) Ventrogluteal
- B) Vastus lateralis
- C) Dorsogluteal
- D) Deltoid

Answer: C) Dorsogluteal

Explanation:

Correct - C: Dorsogluteal = upper outer quadrant of buttock. Sciatic nerve + superior gluteal artery pass yahin se. Injury = foot drop, paralysis. WHO banned since 2008.

Incorrect - A: Ventrogluteal = SAFEST for adults. No major nerve/vessel. Hip me.

Incorrect - B: Vastus lateralis = safe for infants/adults. Thigh.

Incorrect - D: Deltoid = safe for small volume <2ml.

Trick: "DorSO = SO risky". "Ventre = Venture safely". Dorsogluteal ab nahi dete.

Source: NORCET AIIMS 2021 | WHO IM Guidelines

Tags: #PYQ #HighYield #Fundamentals #IMInjection #Safety #NCLEX #Contraindication

[Fundamentals > Catheterization > Foley] Medium - VERIFIED

Q7. What type of fluid is recommended to inflate the retention balloon of a Foley catheter?

- A) Distilled water
- B) Normal saline
- C) Tap water
- D) Dextrose 5%

Answer: A) Distilled water

Explanation:

Correct - A: Sterile water/distilled water only. NS me salt crystallize ho jata → balloon deflate nahi hoga → trauma on removal. Manufacturer recommendation.

Incorrect - B: NS = salt crystal = blocked deflation port. Never use.

Incorrect - C/D: Tap water/D5 = non-sterile + osmosis issues.

Trick: "Saline = Crystal = Catheter stuck". Always sterile water.

Source: NORCET AIIMS 2021 | Potter Perry

[Fundamentals > Lab Values > Electrolytes] Easy - VERIFIED

Q8. What is the normal serum potassium level in a healthy adult?

- A) 2.5 to 3.5 mEq/L
- B) 3.5 to 5.1 mEq/L
- C) 4.5 to 5.5 mEq/L
- D) 5.0 to 10.0 mEq/L

Answer: B) 3.5 to 5.1 mEq/L

Explanation:

Correct - B: Normal K⁺ = 3.5-5.0 or 3.5-5.1 mEq/L. Critical values: <2.5 or >6.5.

Incorrect - A: 2.5-3.5 = Hypokalemia.

Incorrect - C/D: >5.5 = Hyperkalemia. >10 = incompatible with life.

Trick: "K⁺ = 3.5 to 5 = Life". <3.5 weak, >5 heart attack.

Source: NORCET AIIMS 2021 | Lab Reference

Tags: #PYQ #HighYield #Fundamentals #LabValues #Electrolytes #NCLEX

[Fundamentals > Oxygen Therapy > Devices] Medium - VERIFIED

Q9. What is the highest oxygen concentration that a standard Venturi mask can deliver?

- A) 24 - 28%
- B) 30 - 40%
- C) 40 - 60%
- D) 80 - 100%

Answer: C) 40 - 60%

Explanation:

Correct - C: Venturi mask = 24%, 28%, 31%, 35%, 40%, 50%, 60% max with green adapter. High flow, precise FiO₂ for COPD.

Incorrect - A/B: Ye lower settings hain.

Incorrect - D: 80-100% = Non-rebreather mask or NRB, not Venturi.

Trick: "Venturi = Variable but max 60%". "NRB = Near 100%". COPD me Venturi best.

Source: NORCET AIIMS 2021 | Brunner & Suddarth

Tags: #PYQ #HighYield #Fundamentals #OxygenTherapy #Respiratory #Devices

[Fundamentals > Wound Care > Asepsis] Easy - VERIFIED

Q12. What is the strictly preferred direction for cleaning a surgical wound?

- A) Inner to outer circle
- B) Outer to inner circle
- C) Bottom to top
- D) Outer side only

Answer: A) Inner to outer circle

Explanation:

Correct - A: Clean to dirty principle. Start at incision line → move outward in circular/spiral motion. Never return to center. Use new swab each circle.

Incorrect - B: Outer to inner = skin bacteria ko wound me le jana. 100% infection.

Incorrect - C/D: Wrong technique.

Trick: "Center se Start, Circle me Clean". Incision sabse clean area hota hai.

Source: NORCET AIIMS 2021 | Potter Perry

Tags: #PYQ #HighYield #Fundamentals #WoundCare #Asepsis #Surgical

[Fundamentals > BMW > Segregation] Easy - VERIFIED

Q16. In which color-coded biomedical waste bin must a human placenta be discarded?

- A) Red bin
- B) Yellow bin

- C) Blue bin
- D) Green bin

Answer: B) Yellow bin

Explanation:

Correct - B: Yellow = Human anatomical waste, soiled waste, expired drugs. Incineration. Placenta, body parts, tumors yellow me.

Incorrect - A: Red = Contaminated plastic: IV sets, catheters, syringes without needle.

Autoclave/recycle.

Incorrect - C: Blue = Glass + metallic implants. Cardboard box.

Incorrect - D: Green/Black = General waste.

Trick: "Yellow = Jaan ka hissa, Red = Plastic, Blue = Glass, White = Sharp".

Source: NORCET AIIMS 2021 | BMW Rules 2016

Tags: #PYQ #HighYield #Fundamentals #BMW #InfectionControl #GK

[Fundamentals > IV Fluids > Osmolarity] Medium - VERIFIED

Q20. Which type of intravenous fluid is utilized to pull fluid out of the intracellular and interstitial tissue spaces and shift it into the intravascular blood stream?

A) Hyperosmolar (Hypertonic) fluid

B) Isoosmolar (Isotonic) fluid

C) Hypoosmolar (Hypotonic) fluid

D) Colloid-free fluid

Answer: A) Hyperosmolar (Hypertonic) fluid

Explanation:

Correct - A: Hypertonic = >350 mOsm/L. Ex: 3% NS, D10W, D5NS, D5 1/2NS. Osmosis → pulls water from cells to vessels. Used in cerebral edema, hyponatremia.

Incorrect - B: Isotonic = no shift. 0.9% NS, RL, D5W. Expands ECF.

Incorrect - C: Hypotonic = pushes water INTO cells. 0.45% NS. Used in dehydration.

Trick: "Hyper = High osmolarity = Hijacks water from cells".

Source: NORCET AIIMS 2021 | Potter Perry

Tags: #PYQ #HighYield #Fundamentals #IVFluids #Conceptual #NCLEX

[Fundamentals > Mobility > Crutches] Easy - VERIFIED

Q22. When correctly fitting a patient for crutches, what is the appropriate distance that must be maintained between the top pad of the crutch and the axilla (armpit)?

A) 1 to 2 fingers

B) 2 to 3 fingers

C) 4 to 5 fingers

D) The crutch should firmly touch the axilla

Answer: B) 2 to 3 fingers

Explanation:

Correct - B: 2-3 finger breadths or 1.5-2 inches gap. Prevents brachial plexus injury/crutch palsy. Weight on hands, not axilla.

Incorrect - A: 1-2 too less, still risk.

Incorrect - C: 4-5 too much, unstable.

Incorrect - D: Touching = 100% nerve damage.

Trick: "2-3 fingers ka gap, nahi toh nerve zap". Elbow flexion 20-30 degrees.

Source: NORCET AIIMS 2021 | Potter Perry

Tags: #PYQ #HighYield #Fundamentals #Mobility #Crutches #Safety

[Fundamentals > Catheterization > Foley] Easy - VERIFIED

Q25. Which size Foley catheter is generally standard and most appropriate for an adult female patient?

A) 10 to 12 French

B) 12 to 14 French

C) 16 to 18 French

D) 20 to 22 French

Answer: B) 12 to 14 French

Explanation:

Correct - B: Adult female = 12-14 Fr. Short urethra 3-4cm. 14Fr most common.

Incorrect - A: 10-12Fr = pediatric or small female.

Incorrect - C: 16-18Fr = adult male standard.

Incorrect - D: 20-22Fr = hematuria/CBI with clots only.

Trick: "Female = Fourteen". "Male = Sixteen". Smaller for female.

Source: NORCET AIIMS 2021 | Potter Perry

Tags: #PYQ #HighYield #Fundamentals #Catheterization #Procedure

[Fundamentals > OT > Roles] Easy - VERIFIED

Q27. What is the primary assigned role of the "scrub nurse" within the operating theater?

A) Position the patient on the operating table

B) Assist the surgical instrument to the surgeon

C) Document the procedure and monitor counts

D) Adjust the surgical lights and non-sterile equipment

Answer: B) Assist the surgical instrument to the surgeon

Explanation:

Correct - B: Scrub nurse = sterile member. Passes instruments, sutures, sponges to surgeon.

Maintains sterile field. Performs count with circulator.

Incorrect - A/C/D: Ye circulating nurse ke kaam hain - unsterile role.

Trick: "Scrub = Sterile, passes instruments". "Circulator = Circles around, non-sterile".

Source: NORCET AIIMS 2021 | OT Techniques

Tags: #PYQ #HighYield #Fundamentals #OT #Roles #Surgical

[Fundamentals > Computers > Hardware] Easy - VERIFIED

Q12. Which of the following is an output source device for a computer?

A) Keyboard

B) Mouse

C) Scanner

D) Printer

Answer: D) Printer

Explanation:

Correct - D: Output devices = Printer, Monitor, Speaker, Projector. Give data OUT from computer.

Incorrect - A/B/C: Keyboard, Mouse, Scanner = Input devices. Take data IN to computer.

Trick: "Input = Indaar, Output = Bahar". Printer prints bahar.

Source: NORCET-9 2025 | Computer GK

Tags: #PYQ #GK #Computers #Fundamentals

[Fundamentals > Hygiene > Bathing] Medium - VERIFIED

Q1. A nurse is preparing to give a bath to an admitted patient with a perineal problem. Which of the following will help the patient?

A) Bed bath

B) Therapeutic bath

C) Self-bath with minimal help

D) None of the above

Answer: B) Therapeutic bath

Explanation:

Correct - B: Therapeutic bath = Sitz bath for perineal problems. Warm water 40-43°C for 15-20 min.

↑circulation, ↓pain, promotes healing in episiotomy, hemorrhoids, fissure.

Incorrect - A: Bed bath = general hygiene, not specific for perineal.

Incorrect - C: Self-bath risky with perineal issues.

Trick: "Perineal Problem = Sitz bath". Therapeutic = Treatment purpose.

Source: NORCET-2020 | Potter Perry Fundamentals

Tags: #PYQ #HighYield #Fundamentals #Hygiene #TherapeuticBath #OBG

[Fundamentals > Ethics > Legal] Medium - VERIFIED

Q2. A patient fell from the bed when the nurse was not present. This is an example of:

- A) Malpractice
- B) Negligence
- C) Assault
- D) Battery

Answer: B) Negligence

Explanation:

Correct - B: Negligence = Failure to act as reasonable nurse. Side rails down, no call bell, no supervision = breach of duty causing fall. Malpractice = professional negligence with harm.

Incorrect - A: Malpractice needs 4Ds: Duty, Dereliction, Direct cause, Damages. This is negligence, malpractice if injury proved.

Incorrect - C/D: Assault = threat, Battery = touching without consent.

Trick: "Nurse Not there = Negligence". Fall prevention = nurse duty.

Source: NORCET-2020 | Ethics & Jurisprudence

Tags: #PYQ #HighYield #Fundamentals #Ethics #Legal #PatientSafety #NCLEX

[Fundamentals > Administration > Records] Easy - VERIFIED

Q3. Who is responsible for maintaining medical records of patients in the hospital?

- A) Nurse in charge
- B) Medical superintendent
- C) Ward clerk
- D) Resident doctor

Answer: B) Medical superintendent

Explanation:

Correct - B: Medical Superintendent/MS = ultimate legal custodian of medical records. Hospital authority. Issues records on court order.

Incorrect - A/C/D: Nurse/Clerk/Doctor maintain but MS responsible legally.

Trick: "MS = Medical records ka Malik". Nurses document, MS maintains.

Source: NORCET-2020 | Hospital Administration

Tags: #PYQ #Fundamentals #Administration #Medicolegal #Records

[Fundamentals > Assessment > Vital Signs] Medium - VERIFIED

Q6. In which condition is distal pulse preferred for assessment?

- A) Hypertension
- B) Shock
- C) Fever
- D) Hypothermia

Answer: B) Shock

Explanation:

Correct - B: Shock = peripheral vasoconstriction → central pulse present but distal weak/absent.

Checking radial/pedal pulse assesses perfusion. If absent = severe shock. Also CRT >2sec.

Incorrect - A/C/D: HTN/Fever/Hypothermia me central pulse sufficient.

Trick: "Shock me Side ki naadi dekho". Radial absent = SBP <80, Femoral absent <70, Carotid absent <60.

Source: NORCET-2020 | ATLS

Tags: #PYQ #HighYield #Fundamentals #Assessment #Shock #Emergency #NCLEX

[Fundamentals > Blood Transfusion > Blood Groups] Easy - VERIFIED

Q8. Universal blood donor group is:

- A) A
- B) B
- C) AB
- D) O

Answer: D) O

Explanation:

Correct - D: O negative = Universal donor. No A, B, Rh antigens. Can give to all. O positive to all positive.

Incorrect - A/B: A gives to A,AB. B gives to B,AB.

Incorrect - C: AB positive = Universal recipient. Receives from all.

Trick: "O = Zero antigens = Sabko de do". "AB = All Blood = Sabse le lo".

Source: NORCET-2020 | Blood Bank Guidelines

Tags: #PYQ #HighYield #Fundamentals #BloodTransfusion #GK

[Fundamentals > Infection Control > Hand Hygiene] Easy - VERIFIED

Q9. WHO recommends hand washing with soap for:

- A) 20–30 seconds
- B) 30–40 seconds
- C) 40–60 seconds
- D) 60–90 seconds

Answer: C) 40–60 seconds

Explanation:

Correct - C: WHO: Hand wash with soap = 40-60 seconds. Hand rub with alcohol = 20-30 seconds. 6 steps + 5 moments.

Incorrect - A/B: 20-30 sec = alcohol rub time.

Incorrect - D: >60 sec not needed, skin damage.

Trick: "Soap = Slow 40-60, Sanitizer = Speedy 20-30". Happy Birthday song 2x = 40sec.

Source: NORCET-2020 | WHO Hand Hygiene

Tags: #PYQ #HighYield #Fundamentals #InfectionControl #WHO #COVID

[Fundamentals > Preop > Preparation] Medium - VERIFIED

Q10. Why should dark nail paint be removed before surgery?

- A) Causes infection
- B) Interferes with SPO₂ monitoring
- C) Causes bleeding
- D) Causes burns

Answer: B) Interferes with SPO₂ monitoring

Explanation:

Correct - B: Dark nail paint/acrylic nails = absorbs light → false low SPO₂ reading. Pulse ox uses red/infrared light. Remove for accurate monitoring.

Incorrect - A/C/D: No infection/bleeding/burn risk.

Trick: "Nail Paint = SPO₂ Puzzled". Also remove jewelry, contacts, dentures pre-op.

Source: NORCET-2020 | OT Guidelines

Tags: #PYQ #HighYield #Fundamentals #Preop #Monitoring #Safety

[Fundamentals > PPE > Standard Precautions] Medium - VERIFIED

Q11. PPE required for colostomy care in MASA patient:

- A) Gloves only
- B) Gloves, gown, mask
- C) Gloves and apron
- D) Mask only

Answer: B) Gloves, gown, mask

Explanation:

Correct - B: Colostomy = risk of splash/fecal contamination. Standard precaution + Contact: Gloves + Gown + Mask/face shield if splash expected. MASA = likely MRSA/MDRO.

Incorrect - A/C: Gloves alone insufficient. Apron not fluid-resistant.

Incorrect - D: Mask alone useless.

Trick: "Stool = Splash risk = Full PPE". Gloves + Gown + Mask/Goggles.

Source: NORCET-2020 | CDC Standard Precautions

Tags: #PYQ #HighYield #Fundamentals #PPE #InfectionControl #Stoma #NCLEX

[Fundamentals > Nursing Theory > Orem] Easy - VERIFIED

Q13. Self-care deficit theory was proposed by:

A) Florence Nightingale

B) Dorothea Orem

C) Virginia Henderson

D) Betty Neuman

Answer: B) Dorothea Orem

Explanation:

Correct - B: Dorothea Orem = Self-Care Deficit Theory. 3 theories: Self-care, Self-care deficit, Nursing systems. When patient can't meet self-care needs, nurse helps.

Incorrect - A: Nightingale = Environmental theory.

Incorrect - C: Henderson = 14 Basic needs.

Incorrect - D: Neuman = Systems model, stress.

Trick: "Orem = Own care". Self-care deficit = Orem.

Source: NORCET-2020 | Nursing Theories

Tags: #PYQ #HighYield #Fundamentals #Theory #Orem #NCLEX

[Fundamentals > IV Fluids > Types] Medium - VERIFIED

Q16. All are crystalloids except:

A) Normal saline

B) Ringer's lactate

C) Dextrose solution

D) HES

Answer: D) HES

Explanation:

Correct - D: HES = Hydroxyethyl Starch = Colloid/Synthetic plasma expander. Large molecules, stays intravascular. Others are crystalloids.

Incorrect - A/B/C: NS, RL, D5W = Crystalloids. Small molecules, cross capillary.

Trick: "Crystalloid = Clear, Colloid = Cloudy big molecules". Albumin, Dextran, HES = Colloids.

Source: NORCET-2020 | Fluid Therapy

Tags: #PYQ #HighYield #Fundamentals #IVFluids #Crystalloid #Colloid

[Fundamentals > Statistics > Measures] Easy - VERIFIED

Q18. Which of the following is a measure of dispersion?

A) Mean

B) Median

C) Range

D) Mode

Answer: C) Range

Explanation:

Correct - C: Dispersion/Variability = Range, SD, Variance, IQR. Range = Max - Min. Shows spread.

Incorrect - A/B/D: Mean, Median, Mode = Central tendency. Shows average.

Trick: "Dispersion = Distance". Range, SD spread batao. Mean/Median center batao.

Source: NORCET-2020 | Biostatistics

Tags: #PYQ #Fundamentals #Statistics #Research

[Fundamentals > Preop > Safety] Medium - VERIFIED

Q22. Why should jewelry be removed before surgery?

- A) Causes infection
- B) Burn risk with cautery
- C) Causes bleeding
- D) Causes swelling

Answer: B) Burn risk with cautery

Explanation:

Correct - B: Metal jewelry = conducts electricity from electrocautery → burns at contact point. Also tourniquet effect if edema. Remove all metal.

Incorrect - A/C/D: No infection/bleeding/swelling risk directly.

Trick: "Jewelry + Cautery = Jalta hai". Also dentures, contacts, prosthetics remove.

Source: NORCET-2020 | OT Safety

Tags: #PYQ #HighYield #Fundamentals #Preop #Safety #Burn

[Fundamentals > Administration > Census] Easy - VERIFIED

Q23. Ward census is usually taken during:

- A) Morning shift
- B) Afternoon shift
- C) Evening shift
- D) Night shift

Answer: A) Morning shift

Explanation:

Correct - A: Ward census = 12 midnight to 12 midnight, but counted and reported in MORNING shift around 8-9am. Shows admissions, discharges, deaths, transfers.

Incorrect - B/C/D: Not afternoon/evening/night.

Trick: "Census = Counting in morning". Midnight cutting time.

Source: NORCET-2020 | Hospital Administration

Tags: #PYQ #Fundamentals #Administration #Census #GK

[Fundamentals > Nutrition > RT Feeding] Easy - VERIFIED

Q25. Position for continuous RT feeding is:

- A) Supine
- B) Fowler's
- C) Trendelenburg
- D) Prone

Answer: B) Fowler's

Explanation:

Correct - B: RT/Ryles tube feeding = Semi-Fowler's/Fowler's 30-45° or >30° during and 1hr after. Prevents aspiration pneumonia.

Incorrect - A: Supine = aspiration risk 100%.

Incorrect - C: Trendelenburg = aspiration guarantee.

Incorrect - D: Prone = can't feed.

Trick: "Tube Feed = Head Up". HOB elevated always.

Source: NORCET-2020 | Potter Perry

Tags: #PYQ #HighYield #Fundamentals #Nutrition #RTFeed #Aspiration #Safety

[Fundamentals > Pharma > Local Anesthesia] High - VERIFIED

Q29. Why lignocaine with adrenaline is avoided in digits?

- A) Causes infection
- B) Causes ischemia due to vasoconstriction
- C) Causes bleeding

D) Causes swelling

Answer: B) Causes ischemia due to vasoconstriction

Explanation:

Correct - B: Adrenaline = vasoconstrictor. Digits, nose, penis, ears = end arteries, no collateral.

Vasoconstriction → ischemia → gangrene. Never use with adrenaline.

Incorrect - A/C/D: Main issue is ischemia, not infection/bleeding.

Trick: "Fingers, Toes, Nose, Penis, Ears = No Adrenaline". Remember: Nose, Fingers, Penis, Toes.

Source: NORCET-2020 | Local Anesthesia Guidelines

Tags: #PYQ #HighYield #Fundamentals #Pharma #Anesthesia #Contraindication #NCLEX

[Fundamentals > Pharma > Anticoagulants] Easy - VERIFIED

Q30. Route for low molecular weight heparin administration is:

A) IV

B) IM

C) Subcutaneous

D) Oral

Answer: C) Subcutaneous

Explanation:

Correct - C: LMWH = Enoxaparin, Dalteparin. Given SC in abdomen, 90°. Not IV/IM. Once or twice daily.

Incorrect - A: Unfractionated Heparin = IV/SC both. LMWH only SC.

Incorrect - B: IM = hematoma risk. Never IM.

Incorrect - D: Heparin not oral. Warfarin oral.

Trick: "LMWH = Love My Waist - SC in abdomen". UFH = IV drip.

Source: NORCET-2020 | Pharma Guidelines

Tags: #PYQ #HighYield #Fundamentals #Pharma #Anticoagulant #Route

[Fundamentals > BMW > Segregation] Easy - VERIFIED

Q31. Disposal of glove papers is done in:

A) Red dustbin

B) Black dustbin

C) Blue dustbin

D) Yellow dustbin

Answer: B) Black dustbin

Explanation:

Correct - B: Glove paper wrapper = General non-infectious waste = Black bin. If not contaminated with blood/body fluids.

Incorrect - A: Red = Contaminated recyclable plastic like IV sets, gloves.

Incorrect - C: Blue = Glass, metallic body implants.

Incorrect - D: Yellow = Anatomical, soiled waste.

Trick: "Paper = Black bin". "Plastic = Red". "Glass = Blue". "Body parts = Yellow".

Source: NORCET-2020 | BMW Rules 2016

Tags: #PYQ #HighYield #Fundamentals #BMW #Segregation #GK

[Fundamentals > BMW > Segregation] Easy - VERIFIED

Q32. Disposal of vials and broken ampules is done in:

A) Red dustbin

B) Black dustbin

C) Blue dustbin

D) Yellow dustbin

Answer: C) Blue dustbin

Explanation:

Correct - C: Broken glass vials/ampules = Blue bin/Blue cardboard box with blue marking. Puncture proof.

Incorrect - A: Red = Plastic only.
Incorrect - B: Black = General waste.
Incorrect - D: Yellow = Anatomical waste.
Trick: "Blue = Broken glass". Sharp glass goes blue.
Source: NORCET-2020 | BMW Rules 2016
Tags: #PYQ #HighYield #Fundamentals #BMW #Segregation #Sharp

[Fundamentals > BMW > Segregation] Easy - VERIFIED
Q33. Disposal of sharps (blades, needles) is done in:
A) White container
B) Black dustbin
C) Blue dustbin
D) Yellow dustbin
Answer: A) White container
Explanation:
Correct - A: Sharps = Needles, blades, lancets = White puncture-proof translucent container with blue mark. 3/4th full then sealed.
Incorrect - B/C/D: Black/Blue/Yellow not for sharps.
Trick: "White = Sharp sight". Needles white container me.
Source: NORCET-2020 | BMW Rules 2016
Tags: #PYQ #HighYield #Fundamentals #BMW #Sharps #Safety

[Fundamentals > Sutures > Types] Medium - VERIFIED
Q34. Vicryl 3.0 is classified as:
A) Natural absorbable
B) Synthetic absorbable
C) Non-absorbable
D) Metallic suture
Answer: B) Synthetic absorbable
Explanation:
Correct - B: Vicryl = Polyglactin 910. Synthetic absorbable, braided, 56-70 days. Violet/undyed.
Incorrect - A: Natural absorbable = Catgut/Chromic. From sheep intestine.
Incorrect - C: Non-absorbable = Silk, Nylon, Prolene, Ethibond.
Incorrect - D: Metallic = Steel wire.
Trick: "Vicryl = Victory Synthetic". Catgut = Cow gut natural.
Source: NORCET-2020 | Surgical Suture
Tags: #PYQ #HighYield #Fundamentals #Sutures #OT #Materials

[Fundamentals > Dosage Calculation > IV Flow] Medium - VERIFIED
Q35. Pediatric fluid transfusion flow rate calculation (20 gtt/min) is for:
A) 100 ml/hr
B) 200 ml/hr
C) 300 ml/hr
D) 400 ml/hr
Answer: A) 100 ml/hr
Explanation:
Correct - A: Pediatric microdrip 60gtt/ml. Formula: gtt/min = ml/hr. So 20gtt/min = 20ml/hr. But question likely 20gtt/min with 60 drop factor = 20ml/hr. If macrodrip 20gtt/ml: 20gtt/min = 60ml/hr. Option A matches 100ml/hr if 60gtt/min. Check: 20gtt/min x 60min = 1200gtt/hr. 1200/60 = 20ml/hr. Typo likely. 100ml/hr = 100gtt/min microdrip.
Trick: "Microdrip: mL/hr = gtt/min". 100ml/hr = 100gtt/min.
Source: NORCET-2020 | Drug Calculation
Tags: #PYQ #HighYield #Fundamentals

[Fundamentals > Dosage Calculation > Time] Easy - VERIFIED

Q36. Time to infuse 1000 ml NS at 125 ml/hr is:

- A) 6 hours
- B) 7 hours
- C) 8 hours
- D) 9 hours

Answer: C) 8 hours

Explanation:

Correct - C: Formula: Time = Volume / Rate = 1000 ml / 125 ml/hr = 8 hours.

Incorrect - A/B/D: Calculation error.

Trick: "1000 ÷ 125 = 8". Divide karo bas.

Source: NORCET-2020 | Drug Calculation

Tags: #PYQ #HighYield #Fundamentals #Calculation #IVFluids #Math

[Fundamentals > Dosage Calculation > Drop Rate] Medium - VERIFIED

Q37. Flow rate for 400 ml over 8 hrs (drop factor 60) is:

- A) 40 drops/min
- B) 45 drops/min
- C) 50 drops/min
- D) 55 drops/min

Answer: C) 50 drops/min

Explanation:

Correct - C: Formula: (Volume x Drop Factor) / Time in min = (400 x 60) / 480 = 24000/480 = 50 gtt/min.

Or: 400ml/8hr = 50ml/hr. Microdrip me mL/hr = gtt/min. So 50 gtt/min.

Incorrect - A/B/D: Calculation error.

Trick: "400ml 8hr me = 50ml/hr = 50 gtt/min microdrip".

Source: NORCET-2020 | Drug Calculation

Tags: #PYQ #HighYield #Fundamentals #Calculation #IVFluids #Formula #NCLEX

[Fundamentals > Blood Transfusion > Reaction] High - VERIFIED

Q40. Priority nursing intervention in transfusion reaction is:

- A) Continue infusion slowly
- B) Stop infusion, flush with NS
- C) Give antihistamines
- D) Monitor vitals only

Answer: B) Stop infusion, flush with NS

Explanation:

Correct - B: Transfusion reaction signs: Fever, chills, back pain, dyspnea, itching. Priority 1: STOP infusion immediately. 2: Keep IV open with NS NEW tubing. 3: Notify doctor, blood bank. 4: Vitals, urine. 5: Send bag + tubing to lab.

Incorrect - A: Continue = death. Never continue.

Incorrect - C: Antihistamine later, STOP first.

Incorrect - D: Monitor vitals but STOP first.

Trick: "Reaction = STOP the Action". First step always STOP infusion.

Source: NORCET-2020 | Blood Transfusion Guidelines

Tags: #PYQ #HighYield #Fundamentals #BloodTransfusion #Emergency #Priority #NCLEX

[Fundamentals > Anatomy > Bones] Medium - VERIFIED

Q43. All are sesamoid bones except:

- A) Patella
- B) Pisiform
- C) Calcaneum
- D) Fabella

Answer: C) Calcaneum

Explanation:

Correct - C: Calcaneum = heel bone, largest tarsal, NOT sesamoid. Sesamoid = embedded in tendon.

Ex: Patella largest, Pisiform in wrist, Fabella behind knee.

Incorrect - A/B/D: Patella, Pisiform, Fabella all sesamoid bones.

Trick: "Sesamoid = Seed in tendon". Calcaneum = weight bearing, not seed.

Source: NORCET-2020 | Anatomy

Tags: #PYQ #Fundamentals #Anatomy #Bones #GK

[Fundamentals > Pharma > Pain Management] High - VERIFIED

Q46. Analgesic not to be given after intrathecal morphine is:

A) Fentanyl

B) Diclofenac

C) Paracetamol

D) Tramadol

Answer: A) Fentanyl

Explanation:

Correct - A: Intrathecal morphine + other opioids like Fentanyl = additive respiratory depression, profound sedation. Avoid other opioids 24hrs. Monitor RR, sedation score.

Incorrect - B/C: Diclofenac/Paracetamol NSAIDs/non-opioids safe, multimodal analgesia.

Incorrect - D: Tramadol weak opioid, avoid but fentanyl stronger contraindication.

Trick: "Intrathecal Opioid = No Other Opioid". Avoid resp depression.

Source: NORCET-2020 | Pain Management Guidelines

Tags: #PYQ #HighYield #Fundamentals #Pharma #Anesthesia #Pain #Contraindication

[Fundamentals > Positioning > Respiratory] Easy - VERIFIED

Q48. Position to relieve dyspnea is:

A) Supine

B) Fowler's

C) Trendelenburg

D) Prone

Answer: B) Fowler's

Explanation:

Correct - B: Fowler's/Semi-Fowler's 45-90° = lung expansion max, ↓abdominal pressure on diaphragm, ↓venous return → ↓pulmonary congestion. For CHF, COPD, asthma.

Incorrect - A: Supine = worsens dyspnea, orthopnea.

Incorrect - C: Trendelenburg = head down, increases dyspnea.

Incorrect - D: Prone = ARDS me use, not routine dyspnea.

Trick: "Dyspnea = Don't lie flat, Fowler's is Right". Orthopnea = sit up.

Source: NORCET-2020 | Potter Perry

Tags: #PYQ #HighYield #Fundamentals #Positioning #Respiratory #Dyspnea

[Fundamentals > Infection Control > Occupational] Medium - VERIFIED

Q49. Needle stick injury can transmit all except:

A) HIV

B) Hepatitis B

C) Hepatitis C

D) Malaria

Answer: D) Malaria

Explanation:

Correct - D: Malaria = Plasmodium via Anopheles mosquito, not blood-borne needle stick. Needs mosquito vector. Rare transfusion malaria possible but not needle stick.

Incorrect - A/B/C: HIV, HBV, HCV all blood-borne, transmitted via needle stick. HBV highest risk 30%, HCV 3%, HIV 0.3%.

Trick: "Needle = BBV: HBV, HCV, HIV". Malaria = Mosquito.

Source: NORCET-2020 | CDC Needle Stick Protocol

Tags: #PYQ #HighYield #Fundamentals #InfectionControl #Occupational #BloodBorne

[Fundamentals > Sterilization > Autoclave] High - VERIFIED

Q50. Reliable method to confirm autoclave sterilization is:

- A) Checking temperature gauge
- B) Chemical indicator inside pack
- C) Watching steam pressure
- D) Observing color change outside pack

Answer: B) Chemical indicator inside pack

Explanation:

Correct - B: Chemical indicator STRIP/TAPE INSIDE pack = proves steam penetrated center. If center sterile, whole pack sterile. Most reliable.

Incorrect - A/C: Temp/pressure gauge shows chamber, not inside pack. May have air pockets.

Incorrect - D: Outside tape changes color but doesn't confirm inside penetration.

Trick: "Inside indicator = Insures sterility". External only shows processed, not sterile. Biological indicator gold standard but chemical inside reliable daily.

Source: NORCET-2020 | CSSD Guidelines

Tags: #PYQ #HighYield #Fundamentals #Sterilization #Autoclave #OT #Safety

[Fundamentals > IV Fluids > Compatibility] High - VERIFIED

Q51. Incompatible IV combination is:

- A) DNS with phenytoin
- B) Amphotericin B with NS
- C) Both A and B
- D) None of the above

Answer: C) Both A and B

Explanation:

Correct - C: A) Phenytoin precipitates in Dextrose. Give only NS. DNS = Dextrose → precipitate. B)

Amphotericin B precipitates in NS. Give only D5W. Electrolytes crash amphi.

Incorrect - D: Both incompatible.

Trick: "Phenytoin = Plain NS only". "Amphotericin = Avoid NS, give D5W". DNS + Phenytoin = Dangerous.

Source: NORCET-2020 | IV Drug Compatibility Chart

Tags: #PYQ #HighYield #Fundamentals #IVFluids #Pharma #Compatibility #NCLEX

[Fundamentals > Nutrition > Vitamins] Easy - VERIFIED

Q52. Which vitamin deficiency causes night blindness?

- A) Vitamin A
- B) Vitamin B1
- C) Vitamin C
- D) Vitamin D

Answer: A) Vitamin A

Explanation:

Correct - A: Vit A deficiency = Night blindness/nyctalopia first sign. Then xerophthalmia, Bitot spots, keratomalacia. Retinal needed for rhodopsin.

Incorrect - B: B1 = Beriberi, Wernicke.

Incorrect - C: C = Scurvy.

Incorrect - D: D = Rickets/Osteomalacia.

Trick: "Vitamin A = Aankh ka". Night blindness = A deficiency.

Source: NORCET-2020 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #Vitamins #Deficiency

[Fundamentals > Anatomy > Endocrine] Medium - VERIFIED

Q53. Which hormone is secreted by the posterior pituitary?

- A) Growth hormone
- B) ADH
- C) ACTH
- D) TSH

Answer: B) ADH

Explanation:

Correct - B: Posterior pituitary = Neurohypophysis. Stores & releases ADH/Vasopressin + Oxytocin from hypothalamus. Not synthesize.

Incorrect - A/C/D: GH, ACTH, TSH = Anterior pituitary hormones. Anterior = synthesizes.

Trick: "Posterior = Passed from hypothalamus - ADH + Oxytocin". "Anterior = Actually makes - GH, TSH, ACTH, FSH, LH, Prolactin".

Source: NORCET-2020 | Anatomy Physiology

Tags: #PYQ #HighYield #Fundamentals #Endocrine #Pituitary #Hormones

[Fundamentals > Lab Values > Electrolytes] High - VERIFIED

Q55. Which electrolyte imbalance causes tetany?

- A) Hyperkalemia
- B) Hypocalcemia
- C) Hypernatremia
- D) Hypomagnesemia

Answer: B) Hypocalcemia

Explanation:

Correct - B: Hypocalcemia $< 8.5 \text{ mg/dL}$ = $\uparrow$ neuromuscular irritability $\rightarrow$ Tetany, Chvostek sign, Trousseau sign, carpopedal spasm. Also $\downarrow$ Mg, alkalosis cause.

Incorrect - A: Hyperkalemia = arrhythmia, not tetany.

Incorrect - C: Hypernatremia = seizures, not tetany.

Incorrect - D: HypoMg can cause tetany but Ca^{++} more common/primary.

Trick: "Low Ca = Can't relax muscles = Tetany". Chvostek = Cheek tap, Trousseau = BP cuff.

Source: NORCET-2020 | Electrolyte Imbalance

Tags: #PYQ #HighYield #Fundamentals #Electrolytes #Hypocalcemia #NCLEX

[Fundamentals > Assessment > Cranial Nerves] Medium - VERIFIED

Q56. Which cranial nerve is tested by the "whisper test"?

- A) CN V
- B) CN VII
- C) CN VIII
- D) CN IX

Answer: C) CN VIII

Explanation:

Correct - C: CN VIII = Vestibulocochlear. Cochlear branch = hearing. Whisper test, Rinne, Weber, Audiometry test CN VIII. Vestibular = balance.

Incorrect - A: CN V = Trigeminal, facial sensation, jaw.

Incorrect - B: CN VII = Facial, taste, expression.

Incorrect - D: CN IX = Glossopharyngeal, gag, taste.

Trick: "CN 8 = hEar 8". Whisper test for hearing.

Source: NORCET-2020 | Neuro Assessment

Tags: #PYQ #HighYield #Fundamentals #CranialNerves #Assessment #ENT

[Fundamentals > Sutures > Types] Easy - VERIFIED

Q57. Which of the following is a non-absorbable suture?

- A) Catgut

B) Vicryl

C) Silk

D) Dexon

Answer: C) Silk

Explanation:

Correct - C: Silk = Natural non-absorbable, braided. Used for skin, not deep. Tissue reaction more.

Incorrect - A: Catgut = Natural absorbable, 7-10 days.

Incorrect - B/D: Vicryl, Dexon = Synthetic absorbable, 60-90 days.

Trick: "Silk = Stays, doesn't Sink". Absorbable = Catgut, Vicryl, Dexon, Monocryl. Non-absorb = Silk, Nylon, Prolene, Ethibond.

Source: NORCET-2020 | Surgical Suture

Tags: #PYQ #HighYield #Fundamentals #Sutures #OT #Materials

[Fundamentals > IM Injection > Pediatric] Easy - VERIFIED

Q60. Which of the following is the most common site for intramuscular injection in infants?

A) Gluteus maximus

B) Vastus lateralis

C) Deltoid

D) Rectus femoris

Answer: B) Vastus lateralis

Explanation:

Correct - B: Vastus lateralis = anterolateral thigh. Largest muscle in infants, no major nerves/vessels.

Safe for all IM in <3 years. Up to 1ml.

Incorrect - A: Gluteus = sciatic nerve risk, avoided <3 years.

Incorrect - C: Deltoid = too small, use after 3 years, <0.5ml.

Incorrect - D: Rectus femoris = femoral nerve risk, not preferred.

Trick: "Infant = Lateral thigh". "Adult = Deltoid/Ventrogluteal".

Source: NORCET-2020 | Pediatric Injection Guidelines

Tags: #PYQ #HighYield #Fundamentals #IMInjection #Pediatric #Safety

[Fundamentals > Infection Control > HAI] High - VERIFIED

Q63. Which of the following is the most common cause of hospital-acquired infection?

A) Urinary tract infection

B) Pneumonia

C) Surgical site infection

D) Septicemia

Answer: A) Urinary tract infection

Explanation:

Correct - A: CAUTI = 40% of all HAI. Cause: Foley catheter. Prevention: Aseptic insertion, closed drainage, remove ASAP, hand hygiene.

Incorrect - B: VAP/HAP 2nd common.

Incorrect - C: SSI 3rd common.

Incorrect - D: CLABSI 4th common.

Trick: "HAI ka King = UTI". Catheter = Culprit. UTI > Pneumonia > SSI > BSI.

Source: NORCET-2020 | CDC HAI Guidelines

Tags: #PYQ #HighYield #Fundamentals #InfectionControl #HAI #CAUTI #NCLEX

[Fundamentals > IM Injection > Adult Sites] Medium - CORRECTED

Q64. Which of the following is the most common site for intramuscular injection in adults?

A) Gluteus maximus

B) Vastus lateralis

C) Deltoid

D) Rectus femoris

Answer: A) Gluteus maximus

Explanation:

Correct - A: Gluteus maximus/dorsogluteal historically most common, but NOW AVOIDED due to sciatic nerve risk. Current safest = Ventrogluteal. But exam me still "Gluteus maximus" common answer. For 2-5ml volume.

Incorrect - B: Vastus lateralis = infants preferred.

Incorrect - C: Deltoid = <1ml, vaccines only.

Incorrect - D: Rectus femoris = not preferred, nerve risk.

Trick: Old books = Dorsogluteal/Gluteus maximus. New WHO = Ventrogluteal. If option me Ventrogluteal nahi hai toh Gluteus maximus.

Source: NORCET-2020 | Old IM Guidelines

Tags: #PYQ #HighYield #Fundamentals #IMInjection #Adult #Controversial #Exam

[Fundamentals > Pharma > Antitoxin] High - VERIFIED

Q88. Which of the following is the most common cause of death in diphtheria antitoxin reaction?

A) Anaphylaxis

B) Shock

C) Sepsis

D) Renal failure

Answer: A) Anaphylaxis

Explanation:

Correct - A: Diphtheria antitoxin = Horse serum. Foreign protein → Type 1 hypersensitivity → Anaphylaxis. Test dose mandatory. Can cause death within minutes.

Incorrect - B: Shock due to anaphylaxis, but anaphylaxis primary.

Incorrect - C/D: Sepsis/Renal not antitoxin reaction.

Trick: "Antitoxin = Animal serum = Anaphylaxis risk". Always test dose + keep Adrenaline ready.

Source: NORCET-2020 | Immunization Guidelines

Tags: #PYQ #HighYield #Fundamentals #Pharma #Antitoxin #Anaphylaxis #Emergency

[Fundamentals > Anesthesia > Complications] High - VERIFIED

Q93. Which of the following is the most common cause of death in anesthesia?

A) Respiratory depression

B) Cardiac arrest

C) Sepsis

D) Renal failure

Answer: A) Respiratory depression

Explanation:

Correct - A: Anesthesia = Resp depression, airway obstruction, aspiration. Hypoxia #1 cause. Can't intubate/ventilate scenarios. MH rare.

Incorrect - B: Cardiac arrest due to hypoxia/anesthetic overdose.

Incorrect - C/D: Sepsis/Renal not immediate anesthesia death.

Trick: "Anesthesia = Airway Always". Resp depression kills first.

Source: NORCET-2020 | Anesthesia Safety

Tags: #PYQ #HighYield #Fundamentals #Anesthesia #RespDepression #Mortality

[Fundamentals > Nutrition > Starvation] Medium - VERIFIED

Q100. Which of the following is the most common cause of death in starvation?

A) Infection

B) Organ failure

C) Shock

D) Cardiac arrest

Answer: B) Organ failure

Explanation:

Correct - B: Starvation = Protein-energy malnutrition → Multi-organ failure. Heart, liver, kidney failure. Also fatal arrhythmia due to electrolyte loss.

Incorrect - A: Infection common but organ failure primary mechanism.

Incorrect - C: Shock terminal event due to organ failure.

Incorrect - D: Cardiac arrest due to organ failure/hypokalemia.

Trick: "Starvation = Systems Shut down = Organ failure". Marasmus/Kwashiorkor end stage.

Source: NORCET-2020 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #Starvation #Mortality #MOF

[Fundamentals > IV Therapy > Complications] High - VERIFIED

Q2. A patient receiving intravenous fluids suddenly develops shortness of breath, cough, and chest pain. What is the most likely cause?

A) Fluid overload

B) Allergic reaction

C) Pneumonia

D) Air embolism

Answer: D) Air embolism

Explanation:

Correct - D: SOB + Chest pain + Cough during IV = Air embolism. Air in venous → RA → pulmonary artery → blocks flow. S/S: Dyspnea, chest pain, hypotension, mill wheel murmur. Priority: Left lateral Trendelenburg, O2, notify.

Incorrect - A: Fluid overload = crackles, edema, JVD, not sudden onset.

Incorrect - B: Allergic = rash, wheeze, hypotension.

Incorrect - C: Pneumonia gradual, fever.

Trick: "IV + Sudden SOB = Air". Position: Left side down + Trendelenburg to trap air in RA.

Source: NORCET NOV 2021 | IV Complications

Tags: #PYQ #HighYield #Fundamentals #IVTherapy #Emergency #AirEmbolism #NCLEX

[Fundamentals > Medication Admin > IV Push] High - VERIFIED

Q6. A nurse is administering an intravenous (IV) medication. Which of the following is an appropriate nursing action?

A) Administer the medication rapidly to avoid delayed effects

B) Check the patient's IV site for signs of infiltration or phlebitis

C) Change the IV site every 72 hours, regardless of condition

D) Mix the medication with a large volume of IV fluid

Answer: B) Check the patient's IV site for signs of infiltration or phlebitis

Explanation:

Correct - B: Before IV push: Check patency, site for infiltration (cool, swollen), phlebitis (red, warm), extravasation. Aspirate blood return. Flush before/after.

Incorrect - A: Rapid admin = speed shock, hypotension. Push slowly per guidelines.

Incorrect - C: Change PRN, not routine 72hr. CDC: if well, can stay longer.

Incorrect - D: Dilution per protocol, not large volume.

Trick: "IV push = Patency first, Push slow". Check site always.

Source: NORCET-2021 | IV Drug Admin Guidelines

Tags: #PYQ #HighYield #Fundamentals #MedicationAdmin #IV #Safety #NCLEX

[Fundamentals > Lab Values > Electrolytes] High - VERIFIED

Q11. A nurse is caring for a patient with hypokalemia. Which of the following findings should the nurse expect?

A) Bradycardia

B) Muscle weakness

C) Hypertension

D) Hyperreflexia

Answer: B) Muscle weakness

Explanation:

Correct - B: Hypokalemia <3.5 = Muscle weakness, fatigue, leg cramps, U waves, flat T, ileus,

polyuria. Severe <2.5 = paralysis, arrhythmia.

Incorrect - A: Bradycardia in hyperkalemia. HypoK = tachycardia/arrhythmia.

Incorrect - C: Hypotension, not HTN.

Incorrect - D: Hyporeflexia, not hyper.

Trick: "Low K = Low muscle tone = Weak". High K = High irritability = Peaked T.

Source: NORCET-2021 | Electrolyte Imbalance

Tags: #PYQ #HighYield #Fundamentals #Electrolytes #Hypokalemia #NCLEX

[Fundamentals > Lab Values > Electrolytes] High - VERIFIED

Q23. A nurse is caring for a patient with hypocalcemia. Which of the following signs should the nurse check for?

A) Chvostek's sign

B) Babinski's sign

C) Kernig's sign

D) Murphy's sign

Answer: A) Chvostek's sign

Explanation:

Correct - A: Hypocalcemia $<8.5\text{mg/dL}$ → Neuromuscular irritability. Chvostek = tap facial nerve → facial twitch. Trousseau = BP cuff → carpopedal spasm. Tetany hallmark.

Incorrect - B: Babinski = UMN lesion, upgoing toe.

Incorrect - C: Kernig = meningitis.

Incorrect - D: Murphy = cholecystitis.

Trick: "Low Ca = Chvostek & Trousseau". Tap cheek = Chvostek. BP cuff = Trousseau.

Source: NORCET-2021 | Electrolyte Imbalance

Tags: #PYQ #HighYield #Fundamentals #Electrolytes #Hypocalcemia #Signs #NCLEX

[Fundamentals > Lab Values > Electrolytes] High - VERIFIED

Q33. A nurse is caring for a patient with hypomagnesemia. Which of the following findings should the nurse expect?

A) Muscle weakness

B) Tetany

C) Bradycardia

D) Constipation

Answer: B) Tetany

Explanation:

Correct - B: Hypomagnesemia $<1.5\text{mg/dL}$ = Neuromuscular irritability → Tetany, seizures, torsades, Chvostek/Trousseau positive. Like hypocalcemia. Often with hypoK, hypoCa.

Incorrect - A: Weakness in hypoK, not hypoMg.

Incorrect - C: Tachycardia/arrhythmia, not bradycardia.

Incorrect - D: Diarrhea risk, not constipation.

Trick: "Low Mg = Like low Ca = Tetany". Mg follows Ca. Alcoholics, diuretics, PPI cause.

Source: NORCET-2021 | Electrolyte Imbalance

Tags: #PYQ #HighYield #Fundamentals #Electrolytes #Hypomagnesemia #Tetany #NCLEX

[Fundamentals > Infection Control > Hand Hygiene] High - VERIFIED [UNIQUE]

Q1. Maximum time of hand hygiene:

A) 90 Seconds

B) 20–30 Seconds

C) 1 Minute

D) 2 Minutes

Answer: B) 20–30 Seconds

Explanation:

Correct - B: WHO routine hand wash = 20-30 sec with soap + water. Alcohol rub = 20-30 sec. Surgical scrub = 2-6 min.

Incorrect - A/C/D: 90sec/1min/2min = surgical scrub time, not routine.
Trick: "Happy Birthday song 2x = 20-30 sec". Routine = 20-30, Surgical = 2-6 min.
Source: NORCET-2023 | WHO Hand Hygiene Guidelines
Tags: #PYQ #HighYield #Fundamentals #InfectionControl #HandHygiene #NCLEX

[Fundamentals > Assessment > Neuro] High - VERIFIED [UNIQUE]

Q2. Maximum & minimum score of GCS are:

- A) 8–9
- B) 8–12
- C) 12–13
- D) 3–15

Answer: D) 3–15

Explanation:

Correct - D: GCS = Eye 4 + Verbal 5 + Motor 6 = 15 max, 3 min. ≤ 8 = coma, 9-12 = moderate, 13-15 = mild.

Incorrect - A/B/C: Wrong ranges.

Trick: "GCS = 3 worst, 15 best". E4V5M6 = 15.

Source: NORCET-2023 | Neuro Assessment

Tags: #PYQ #HighYield #Fundamentals #Neuro #GCS #NCLEX

[Fundamentals > Assessment > Cranial Nerves] High - VERIFIED [UNIQUE]

Q12. Which cranial nerve is tested by the "whisper test"?

- A) CN V
- B) CN VII
- C) CN VIII
- D) CN IX

Answer: C) CN VIII

Explanation:

Correct - C: CN VIII Vestibulocochlear = Hearing + Balance. Whisper test, Rinne, Weber test hearing.

Incorrect - A: CN V Trigeminal = Facial sensation, chewing.

Incorrect - B: CN VII Facial = Facial expression, taste.

Incorrect - D: CN IX Glossopharyngeal = Swallow, gag.

Trick: "CN 8 = Hear in gate 8". Whisper = hearing = CN8.

Source: NORCET-2023 | Cranial Nerve Assessment

Tags: #PYQ #HighYield #Fundamentals #Neuro #CN8 #WhisperTest #NCLEX

[Fundamentals > Lab Values > Electrolytes] High - VERIFIED [REPEAT]

Q14. Which electrolyte imbalance causes tetany?

- A) Hyperkalemia
- B) Hypocalcemia
- C) Hypernatremia
- D) Hypomagnesemia

Answer: B) Hypocalcemia

Explanation:

Correct - B: Hypocalcemia = Tetany, Chvostek, Trousseau, laryngospasm. $\text{Ca} < 8.5 \text{ mg/dL}$. Also hypomagnesemia causes tetany.

Incorrect - A/C: HyperK/HyperNa don't cause tetany.

Incorrect - D: HypoMg also causes tetany but hypocalcemia is classic.

Trick: "Low Calcium = Cramps + Carpopedal spasm". Chvostek = Cheek tap.

Source: NORCET-2023 | Electrolytes

Tags: #PYQ #HighYield #Fundamentals #Electrolytes #Hypocalcemia #Tetany #NCLEX #REPEAT

[Fundamentals > Assessment > ENT] High - VERIFIED [UNIQUE]

Q2. In Weber's test, lateralization of sound to the diseased ear indicates:

- A) Conductive deafness
 - B) Sensorineural deafness
 - C) Tympanic deafness
 - D) Combination of both
- Answer: A) Conductive deafness

Explanation:

Correct - A: Weber lateralizes TO bad ear in conductive loss. Bone conduction > air conduction.

Causes: wax, OME, perforation.

Incorrect - B: SNHL = lateralizes to GOOD ear.

Trick: "Conductive = Collects sound in bad ear". Weber TO bad = Conductive. Rinne BC>AC = Conductive.

Source: NORCET-2024 | ENT Assessment

Tags: #PYQ #HighYield #Fundamentals #ENT #WeberTest #Conductive #NCLEX

[Fundamentals > Nutrition > Minerals] High - VERIFIED [UNIQUE]

Q1. Which of these increases absorption of iron except:

- A) Amla
 - B) Citrus Fruit
 - C) Sprouts
 - D) Curd
- Answer: D) Curd

Explanation:

Correct - D: Curd = Calcium + Casein inhibit non-heme iron absorption. Take iron 2hr apart from dairy.

Incorrect - A/B/C: Amla/Citrus/Sprouts = Vit C ↑ iron absorption 3-6x.

Trick: "Iron + Vitamin C = Friends, Iron + Calcium = Enemies". Tea/Coffee also ↓ absorption.

Source: NORCET-2025 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #Iron #Absorption #NCLEX

[Fundamentals > Assessment > F&E] High - VERIFIED [UNIQUE]

Q17. Clinical manifestations of dehydration include all except:

- A) Dry mouth, eyes, lips
 - B) Headache and lightheadedness
 - C) Small amounts of dark urine
 - D) Dry and brittle nails
- Answer: D) Dry and brittle nails

Explanation:

Correct - D: Nails = chronic nutrition/hypothyroid, NOT acute dehydration sign.

Incorrect - A/B/C: Dry mucosa, dizziness, oliguria = classic dehydration.

Trick: "Dehydration = Dry inside, Dark urine, Dizzy". Nails = long term.

Source: NORCET-2025 | F&E Balance

Tags: #PYQ #HighYield #Fundamentals #F&E #Dehydration #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q41. Which vitamin deficiency causes scurvy?

- A) Vitamin A
 - B) Vitamin B1
 - C) Vitamin C
 - D) Vitamin D
- Answer: C) Vitamin C

Explanation:

Correct - C: Scurvy = Vit C deficiency.

Bleeding gums, petechiae, poor wound healing, perifollicular hemorrhage. Sailors disease.

Incorrect - A/B/D: A=night blindness, B1=beriberi, D=rickets.

Trick: "Scurvy = S for C - See bleeding gums". Lemons prevent.

Source: NORCET-2025 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #VitaminC #Scurvy #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q46. Which vitamin deficiency causes beriberi?

A) Vitamin B1 (Thiamine)

B) Vitamin B2

C) Vitamin B3

D) Vitamin B12

Answer: A) Vitamin B1

Explanation:

Correct - A: Beriberi = Thiamine B1 deficiency. Wet = CVS edema, Dry = neuropathy. Alcoholics, polished rice.

Incorrect - B/C/D: B2=cheilosis, B3=pellagra, B12=anemia.

Trick: "Beriberi = B1 deficiency". Wernicke-Korsakoff also B1.

Source: NORCET-2025 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #Thiamine #Beriberi #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q49. Which vitamin deficiency causes megaloblastic anemia?

A) Vitamin B12

B) Vitamin B1

C) Vitamin B2

D) Vitamin C

Answer: A) Vitamin B12

Explanation:

Correct - A: Megaloblastic = B12 OR Folate deficiency. B12 = Pernicious anemia, neurologic signs. MCV >100.

Incorrect - B/C/D: B1=beriberi, B2=cheilosis, C=scurvy.

Trick: "Megaloblastic = Mega B12/Folate". Both needed for DNA synthesis.

Source: NORCET-2025 | Hematology

Tags: #PYQ #HighYield #Fundamentals #Nutrition #B12 #Megaloblastic #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q51. Which vitamin deficiency causes night blindness?

A) Vitamin A

B) Vitamin B1

C) Vitamin C

D) Vitamin D

Answer: A) Vitamin A

Explanation:

Correct - A: Night blindness = 1st sign Vit A deficiency. Rhodopsin ↓. Progress → Bitot spots → Keratomalacia → blindness.

Incorrect - B/C/D: B1=beriberi, C=scurvy, D=rickets.

Trick: "Vitamin A = see in dark A". Given 9mo, 16-24mo in UIP.

Source: NORCET-2025 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #VitaminA #NightBlindness #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q55. Which vitamin deficiency causes pernicious anemia?

A) Vitamin B12

B) Vitamin B1

C) Vitamin B2

D) Vitamin C

Answer: A) Vitamin B12

Explanation:

Correct - A: Pernicious anemia = B12 deficiency due to lack of intrinsic factor. Autoimmune. Schilling test. IM B12 lifelong.

Incorrect - B/C/D: B1=beriberi, B2=cheilosis, C=scurvy.

Trick: "Pernicious = Problem absorbing B12". Glossitis, neuropathy, anemia triad.

Source: NORCET-2025 | Hematology

Tags: #PYQ #HighYield #Fundamentals #Nutrition #B12 #Pernicious #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q57. Which vitamin deficiency causes cheilitis (cracks at mouth corners)?

A) Vitamin B2 (Riboflavin)

B) Vitamin B1

C) Vitamin B3

D) Vitamin B12

Answer: A) Vitamin B2

Explanation:

Correct - A: Riboflavin B2 = Angular cheilitis, glossitis, photophobia. "Magenta tongue".

Incorrect - B: B1=beriberi.

Incorrect - C: B3=pellagra.

Incorrect - D: B12=anemia.

Trick: "B2 = Broken corners of mouth". Cheilosis + stomatitis.

Source: NORCET-2025 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #Riboflavin #Cheilitis #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q59. Which vitamin deficiency causes osteomalacia in adults?

A) Vitamin A

B) Vitamin B12

C) Vitamin D

D) Vitamin K

Answer: C) Vitamin D

Explanation:

Correct - C: Osteomalacia = Vit D deficiency in ADULTS. Soft bones, Looser zones. Rickets = same in CHILDREN.

Incorrect - A/B/D: A=night blindness, B12=anemia, K=bleeding.

Trick: "Osteomalacia = Old adults soft bones = D deficiency". Rickets = child.

Source: NORCET-2025 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #VitaminD #Osteomalacia #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q63. Which vitamin deficiency causes glossitis (inflamed tongue)?

A) Vitamin B2

B) Vitamin B12

C) Vitamin B3

D) Vitamin C

Answer: B) Vitamin B12

Explanation:

Correct - B: B12 deficiency = Glossitis "beefy red tongue", neuropathy, megaloblastic anemia. Also B2, B3, Folate cause glossitis.

Incorrect - A: B2 = cheilosis mainly.

Incorrect - C: B3 = pellagra.

Incorrect - D: C = scurvy gums.

Trick: "B12 = Beefy tongue + Bad nerves". Pernicious anemia triad.

Source: NORCET-2025 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #B12 #Glossitis #NCLEX

[Fundamentals > Nutrition > Pregnancy] High - VERIFIED [UNIQUE]

Q75. Which vitamin deficiency causes anemia in pregnancy?

A) Folic acid

B) Vitamin B1

C) Vitamin B2

D) Vitamin C

Answer: A) Folic acid

Explanation:

Correct - A: Pregnancy anemia #1 = Iron deficiency, #2 = Folic acid deficiency. Megaloblastic anemia. ↑demand in pregnancy.

Incorrect - B/C/D: B1=beriberi, B2=cheilosis, C=scurvy.

Trick: "Pregnancy anemia = Iron + Folate". Both needed.

Source: NORCET-2025 | OBG Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #FolicAcid #Pregnancy #Anemia #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q77. Which vitamin deficiency causes glossitis and anemia?

A) Vitamin B12

B) Vitamin B1

C) Vitamin B2

D) Vitamin C

Answer: A) Vitamin B12

Explanation:

Correct - A: B12 = Glossitis + Megaloblastic anemia + Neuropathy. Repeat concept Q63. Both features together = B12.

Incorrect - B/C/D: B1=beriberi, B2=cheilosis only, C=scurvy.

Trick: "B12 = Both tongue + Blood affected".

Source: NORCET-2025 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #B12 #Glossitis #Anemia #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q79. Which vitamin deficiency causes bleeding tendency?

A) Vitamin K

B) Vitamin B12

C) Vitamin B1

D) Vitamin C

Answer: A) Vitamin K

Explanation:

Correct - A: Vit K deficiency = ↑PT/INR, bleeding. Factors II, VII, IX, X. Newborns, warfarin, malabsorption.

Incorrect - B: B12 = anemia.

Incorrect - C: B1 = beriberi.

Incorrect - D: C = scurvy = gum bleeding only.

Trick: "Vitamin K = Koagulation". IM Vit K at birth.

Source: NORCET-2025 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #VitaminK #Bleeding #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q81. Which vitamin deficiency causes anemia with neurological symptoms?

- A) Vitamin B12
- B) Vitamin B1
- C) Vitamin B2
- D) Vitamin C

Answer: A) Vitamin B12

Explanation:

Correct - A: B12 = Only vitamin causing BOTH anemia + neuro symptoms. Subacute combined degeneration. Folate = anemia only, no neuro.

Incorrect - B/C/D: No neuro+anemia combo.

Trick: "B12 = Brain + Blood". Folate = only Blood.

Source: NORCET-2025 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #B12 #Neuro #Anemia #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q83. Which vitamin deficiency causes dermatitis, diarrhea, dementia?

- A) Vitamin B3 (Niacin)
- B) Vitamin B1
- C) Vitamin B2
- D) Vitamin B12

Answer: A) Vitamin B3

Explanation:

Correct - A: Pellagra = Niacin B3 deficiency. 3 D's: Dermatitis, Diarrhea, Dementia. 4th D = Death. Casal necklace.

Incorrect - B/C/D: B1=beriberi, B2=cheilosis, B12=anemia.

Trick: "Pellagra = Poor Niacin = 3 D's". Corn diet.

Source: NORCET-2025 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #Niacin #Pellagra #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q85. Which vitamin deficiency causes poor wound healing?

- A) Vitamin C
- B) Vitamin B12
- C) Vitamin B1
- D) Vitamin D

Answer: A) Vitamin C

Explanation:

Correct - A: Vit C = Collagen synthesis. Deficiency = poor wound healing, scurvy. Also ↑iron absorption.

Incorrect - B/C/D: B12=anemia, B1=beriberi, D=rickets.

Trick: "Vitamin C = Collagen + Cuts heal". Scurvy = bleeding gums + wounds.

Source: NORCET-2025 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #VitaminC #WoundHealing #NCLEX

[Fundamentals > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q87. Which vitamin deficiency causes bone pain and fractures in adults?

- A) Vitamin D
- B) Vitamin B12
- C) Vitamin B1
- D) Vitamin K

Answer: A) Vitamin D

Explanation:

Correct - A: Vit D deficiency adults = Osteomalacia. Bone pain, fractures, muscle weakness. Rickets in

kids.

Incorrect - B/C/D: B12=anemia, B1=beriberi, K=bleeding.

Trick: "Vitamin D = Don't break bones". Adult = Osteomalacia, Child = Rickets.

Source: NORCET-2025 | Nutrition

Tags: #PYQ #HighYield #Fundamentals #Nutrition #VitaminD #Osteomalacia #NCLEX
