

Pediatric

26 April 2026 21:30

[Pediatric > GI > Diarrhea Management] Medium - VERIFIED

Q1. A mother of a 4-month-old infant with diarrhea is being taught about preventing dehydration. Which of the following is the most appropriate feeding choice?

- A) Fruit juice
- B) Diluted carbonated drink
- C) Soy-based lactose-free formula
- D) Regular formula mixed with electrolyte solution

Answer: C) Soy-based lactose-free formula

Explanation:

Correct - C: Diarrhea causes secondary lactase deficiency. Lactose-free soy formula provides nutrition without worsening diarrhea. WHO/IMNCI recommends continued breastfeeding + ORS, but if formula-fed, switch to lactose-free temporarily.

Incorrect - A/B: High osmolarity pulls water into gut → osmotic diarrhea worsens.

Incorrect - D: Don't mix ORS with formula. Give separately.

Trick: Students choose D thinking "electrolyte" is good, but mixing dilutes both.

Source: NORCET-2023 | IMNCI Guidelines

Tags: #PYQ #HighYield #Pediatric #IMNCI #Nutrition #TrickQ

[Pediatric > Neonatal > Hyperbilirubinemia] Medium - VERIFIED

Q34. Which nursing intervention is a top priority when caring for a newborn actively receiving continuous phototherapy for hyperbilirubinemia?

- A) Remove eye patches frequently
- B) Change the baby's position every 4 hours
- C) Encourage breastfeeding as much as possible
- D) Apply thick lotion to prevent skin breakdown

Answer: C) Encourage breastfeeding as much as possible

Explanation:

Correct - C: Phototherapy = ↑insensible water loss + loose stools. Risk: Dehydration. Priority: Hydration via frequent breastfeeding q2-3h. Also promotes stool excretion of bilirubin.

Incorrect - A: Eye patches MUST stay on 24/7. Remove only for feeds. Retinal damage risk.

Incorrect - B: Position change q2h, not q4h. Expose max skin surface.

Incorrect - D: NO lotions/oils - causes burns under lights. Skin kept dry.

Trick: "Photo = Fluid Loss = Feed More". No lotion, no diaper, yes eye patch, yes rotation.

Source: NORCET-2023 | Wong's Pediatric

Tags: #PYQ #HighYield #Pediatric #Neonatal #Phototherapy #Priority #NCLEX

[Pediatric > Neuro > Meningitis] High - VERIFIED

Q42. Which of the following is considered an early clinical manifestation of meningitis in a neonate?

- A) Neck stiffness (Nuchal rigidity)
- B) Positive Kernig's sign
- C) High-pitched cry
- D) Photophobia

Answer: C) High-pitched cry

Explanation:

Correct - C: Neonates <3 months: No nuchal rigidity/Kernig/Brudzinski. Open fontanelles decompress pressure. Early signs: Poor feeding, lethargy, irritability, HIGH-PITCHED CRY, seizures, bulging fontanelle.

Incorrect - A/B/D: These are ADULT signs. Photophobia in older kids, not neonates.

Trick: "Baby can't complain neck pain, so baby CRIES high-pitched". Neonate = Nonspecific signs.

Source: NORCET-2023 | Ghai Pediatrics

Tags: #PYQ #HighYield #Pediatric #Neuro #Infection #Emergency #NCLEX

[Pediatric > Growth > Anthropometry] Easy - VERIFIED

Q48. How is the physical growth parameter (height) measured in a healthy 2-year-old child?

- A) Standing height using a stadiometer
- B) Crown-rump length
- C) Supine length (lying down)
- D) Arm span

Answer: C) Supine length (lying down)

Explanation:

Correct - C: WHO/CDC: <24 months = Supine length on infantometer. ≥24 months = Standing height on stadiometer. 2-yr-old = 24 months = transition age, but standard is still supine if <2yr. Standing height ~0.7cm less than supine.

Incorrect - A: Standing after 2 years when child can stand straight.

Incorrect - B: Crown-rump = sitting height, for disproportionate growth.

Trick: "2 saal ke neeche Letao, 2 ke upar Khada karo". Length vs Height terminology.

Source: NORCET-2023 | WHO Growth Standards

Tags: #PYQ #HighYield #Pediatric #Growth #Assessment #WHO

[Pediatric > Growth > Milestones] Easy - VERIFIED

Q6. At what age does a healthy infant's birth weight typically triple?

- A) 6 months
- B) 9 months
- C) 1 year
- D) 2 years

Answer: C) 1 year

Explanation:

Correct - C: Birth weight: Double by 5-6 months, Triple by 12 months/1 year, Quadruple by 2 years.

Incorrect - A: 6 months = double only.

Incorrect - B/D: 9 months ya 2 years me triple nahi hota.

Trick: "Double 6, Triple 12, Quadruple 24 months". 6-12-24 pattern.

Source: NORCET AIIMS 2021 | Ghai Pediatrics

Tags: #PYQ #HighYield #Pediatric #Growth #Milestones #Formula

[Pediatric > Neonatal > KMC] Easy - VERIFIED

Q11. What is the primary clinical purpose of providing Kangaroo Mother Care (KMC) to a premature infant?

- A) To prevent infection
- B) To provide contraception for the mother
- C) To maintain the baby's body temperature
- D) To prevent hyperthermia

Answer: C) To maintain the baby's body temperature

Explanation:

Correct - C: KMC = Skin-to-skin contact, exclusive breastfeeding, early discharge. Primary goal: Prevent hypothermia in LBW/preterm. Maintains temp 36.5-37.5°C. Also ↑bonding, ↑weight gain, ↑O₂ sat.

Incorrect - A: Infection ↓ hoti hai but primary nahi.

Incorrect - B/D: No relation to contraception. Prevents hypo, not hyperthermia.

Trick: "Kangaroo = Keep warm". LBW <2500g ke liye mandatory.

Source: NORCET AIIMS 2021 | WHO KMC Guidelines

Tags: #PYQ #HighYield #Pediatric #Neonatal #KMC #WHO #LBW

[Pediatric > Procedures > Suctioning] Medium - VERIFIED

Q13. Which size suction catheter is typically appropriate for clearing the airway of a pediatric patient?

- A) 8 French
- B) 10 French
- C) 12 French
- D) 14 French

Answer: A) 8 French

Explanation:

Correct - A: Pediatrics: 6-10 Fr. Formula = (Age in years + 16) / 2. Or ETT size x 2. 8Fr safe for infants/children.

Incorrect - B: 10 Fr = older child/teen.

Incorrect - C/D: 12-14 Fr = adults. Pediatric airway me trauma karega.

Trick: "Small child = Small size". Adult = 12-16Fr. Never occlude >50% of ETT.

Source: NORCET AIIMS 2021 | PALS Guidelines

Tags: #PYQ #HighYield #Pediatric #Procedures #Suctioning #Airway

[Pediatric > Immunization > Routes] Medium - VERIFIED

Q15. All of the following vaccines are administered via the intramuscular (IM) route EXCEPT:

- A) Hepatitis B
- B) Measles
- C) DPT
- D) Tetanus Toxoid

Answer: B) Measles

Explanation:

Correct - B: Measles/MMR/MR = Subcutaneous route, right upper arm. Live attenuated.

Incorrect - A/C/D: Hep B, DPT, TT = IM, anterolateral thigh infants or deltoid adults. All contain adjuvant.

Trick: "Measles = Muscle me nahi, SubCut me". BCG = Intradermal. OPV/Rota = Oral. Rest mostly IM.

Source: NORCET AIIMS 2021 | UIP Schedule

Tags: #PYQ #HighYield #Pediatric #Immunization #Routes #UIP

[Pediatric > Neonatal > Infection] High - VERIFIED

Q8. A newborn whose mother is positive for Chlamydia trachomatis should be optimally treated with which of these to prevent ophthalmia neonatorum?

- A) Oral Erythromycin
- B) Erythromycin 0.5% ointment
- C) Tetracycline 1% ointment
- D) Silver nitrate 1% drops

Answer: A) Oral Erythromycin

Explanation:

Correct - A: Chlamydia trachomatis = systemic infection. Oral Erythromycin 50mg/kg/day x 14 days needed. Prevents pneumonia also. Topical not enough.

Incorrect - B/C: Erythromycin/Tetra ointment = for Gonococcal ophthalmia prophylaxis at birth, NOT treatment. Chlamydia needs systemic.

Incorrect - D: Silver nitrate = for Gonorrhea prophylaxis, banned now due to chemical conjunctivitis. Not for Chlamydia.

Trick: "Chlamydia = Chronic systemic = Oral needed". "Gonorrhea = Genital local = Topical ok".

Chlamydia causes pneumonia later if not treated.

Source: NORCET-9 2025 | Nelson Pediatrics

Tags: #PYQ #HighYield #Pediatric #Neonatal #Infection #STI #Pharma #NCLEX

[Pediatric > Neonatal > Thermoregulation] High - VERIFIED

Q7. Immediate nursing intervention for a neonate with hypothermia is:

- A) Cover with blanket
 - B) Place baby under radiant warmer
 - C) Give warm fluids
 - D) Apply hot water bag
- Answer: B) Place baby under radiant warmer

Explanation:

Correct - B: Neonatal hypothermia $<36.5^{\circ}\text{C}$. Priority: Radiant warmer or incubator. Skin-to-skin KMC if stable. Rewarm 0.5°C/hr . Monitor temp, sugar, O₂.

Incorrect - A: Blanket alone insufficient, blocks observation. Use with warmer.

Incorrect - C: No oral fluids in hypothermia, risk aspiration.

Incorrect - D: Hot water bag = burns. Never use.

Trick: "Hypothermia = Heater chahiye". Warm slowly, not rapidly.

Source: NORCET-2020 | WHO ENC Guidelines

Tags: #PYQ #HighYield #Pediatric #Neonatal #Hypothermia #Emergency #Priority

[Pediatric > NICU > Feeding] Medium - VERIFIED

Q26. Enteral feed flushing with air instead of water is usually done in:

- A) ICU
- B) NICU
- C) PICU
- D) Surgical ward

Answer: B) NICU

Explanation:

Correct - B: NICU = VLBW/preterm babies. Fluid restriction strict. Air 1-2ml used to flush OG/NG tube to clear, avoid fluid overload. Check placement.

Incorrect - A/C/D: ICU/PICU/Surgical = water 5-30ml used to flush. Fluid tolerance more.

Trick: "NICU = No extra water, use Air". Premature = fluid overload risk.

Source: NORCET-2020 | NICU Protocols

Tags: #PYQ #HighYield #Pediatric #NICU #Feeding #FluidBalance

[Pediatric > Immunization > Vaccine Types] Medium - VERIFIED

Q54. Which of the following is a live attenuated vaccine?

- A) Polio (Salk)
- B) BCG
- C) Hepatitis B
- D) Rabies (inactivated)

Answer: B) BCG

Explanation:

Correct - B: Live attenuated: BCG, MMR, OPV Sabin, Varicella, Yellow fever, Rotavirus. BCG = Mycobacterium bovis live attenuated.

Incorrect - A: Salk Polio = Killed/inactivated.

Incorrect - C: Hep B = Recombinant subunit.

Incorrect - D: Rabies = Inactivated.

Trick: "Live = BCG, Measles, Mumps, Rubella, OPV, Varicella". Killed = Salk, Hep A, Rabies, Influenza.

Source: NORCET-2020 | UIP Schedule

Tags: #PYQ #HighYield #Pediatric #Immunization #VaccineTypes #UIP

[Pediatric > GI > Diarrhea] High - VERIFIED

Q66. Which of the following is the most common cause of diarrhea in children?

- A) Rotavirus
- B) E. coli
- C) Salmonella
- D) Shigella

Answer: A) Rotavirus

Explanation:

Correct - A: Rotavirus = #1 cause of severe dehydrating diarrhea in <5 years globally. Winter peak.

Vaccine available. Watery, non-bloody.

Incorrect - B/C/D: E. coli, Salmonella, Shigella = bacterial, bloody diarrhea. Less common than rota.

Trick: "Rota = Real common in kids". Winter diarrhea = Rota. Summer = E.coli.

Source: NORCET-2020 | Ghai Pediatrics

Tags: #PYQ #HighYield #Pediatric #GI #Diarrhea #Viral #WHO

[Pediatric > Respiratory > Pneumonia] High - VERIFIED

Q67. Which of the following is the most common cause of pneumonia in children?

A) Streptococcus pneumoniae

B) Staphylococcus aureus

C) Klebsiella pneumoniae

D) Haemophilus influenzae

Answer: A) Streptococcus pneumoniae

Explanation:

Correct - A: S. pneumoniae = Most common bacterial pneumonia all ages. Lobar pneumonia. PCV vaccine given.

Incorrect - B: Staph aureus = post-influenza, abscess, pneumatoceles.

Incorrect - C: Klebsiella = alcoholics, DM, currant jelly sputum.

Incorrect - D: H. influenzae type B = 2nd common before Hib vaccine, now less.

Trick: "Pneumo = Pneumonia". PCV vaccine for Pneumo.

Source: NORCET-2020 | Ghai Pediatrics

Tags: #PYQ #HighYield #Pediatric #Respiratory #Pneumonia #Bacterial

[Pediatric > Neonatal > Meningitis] High - VERIFIED

Q68. Which of the following is the most common cause of meningitis in neonates?

A) Group B Streptococcus

B) E. coli

C) Listeria monocytogenes

D) All of the above

Answer: D) All of the above

Explanation:

Correct - D: Neonatal meningitis <28 days: GBS 50%, E. coli 20%, Listeria 5-10%. All 3 main causes. Vertical transmission.

Incorrect - A/B/C: Alone galat, all 3 common.

Trick: "Neonatal meningitis = GEL: GBS, E.coli, Listeria". After 1 month = S. pneumo, N. meningitidis, H. flu.

Source: NORCET-2020 | Nelson Pediatrics

Tags: #PYQ #HighYield #Pediatric #Neonatal #Meningitis #Infection #NCLEX

[Pediatric > Infectious > Measles] High - VERIFIED

Q83. Which of the following is the most common cause of death in measles?

A) Pneumonia

B) Encephalitis

C) Diarrhea

D) Malnutrition

Answer: A) Pneumonia

Explanation:

Correct - A: Measles = immune suppression → Secondary bacterial pneumonia > Giant cell pneumonia. #1 cause of death worldwide. Esp <5 years, malnourished.

Incorrect - B: Encephalitis 1/1000 cases, fatal but less common than pneumonia.

Incorrect - C/D: Diarrhea/Malnutrition contribute but pneumonia #1.

Trick: "Measles = Malnourished lung infection". Pneumonia kills most kids.

Source: NORCET-2020 | WHO Measles Guidelines

Tags: #PYQ #HighYield #Pediatric #Infectious #Measles #Pneumonia #Mortality

[Pediatric > Infectious > Pertussis] High - VERIFIED

Q85. Which of the following is the most common cause of death in pertussis?

- A) Pneumonia
- B) Encephalopathy
- C) Apnea
- D) Seizures

Answer: A) Pneumonia

Explanation:

Correct - A: Pertussis/Whooping cough = Secondary bacterial pneumonia #1 cause of death. Esp <6 months. Also apnea in infants.

Incorrect - B: Encephalopathy 1% cases, due to hypoxia.

Incorrect - C: Apnea common in infants but pneumonia kills more.

Incorrect - D: Seizures due to hypoxia/encephalopathy.

Trick: "Pertussis = Pneumonia kills". Whooping cough → lung infection.

Source: NORCET-2020 | Ghai Pediatrics

Tags: #PYQ #HighYield #Pediatric #Infectious #Pertussis #Pneumonia #Mortality

[Pediatrics > Growth > Milestones] High - VERIFIED [UNIQUE]

Q10. Birth weight quadruples by:

- A) 5 Months
- B) 1 Year
- C) 18 Months
- D) 2 Years

Answer: D) 2 Years

Explanation:

Correct - D: Birth weight doubles 5 months, triples 1 year, quadruples 2 years. Height doubles 4 years.

Incorrect - A: 5 months = doubles.

Incorrect - B: 1 year = triples.

Incorrect - C: 18 months = between 3x-4x.

Trick: "2-3-4 rule: 2x at 5mo, 3x at 1yr, 4x at 2yr". Height 2x at 4yr.

Source: NORCET-2023 | Growth Milestones

Tags: #PYQ #HighYield #Pediatrics #Growth #Weight #NCLEX

[Pediatrics > Nutrition > Vitamins] High - VERIFIED [UNIQUE]

Q11. Which vitamin deficiency causes night blindness?

- A) Vitamin A
- B) Vitamin B1
- C) Vitamin C
- D) Vitamin D

Answer: A) Vitamin A

Explanation:

Correct - A: Vit A deficiency = Night blindness, xerophthalmia, Bitot spots, keratomalacia. Rhodopsin formation needs Vit A.

Incorrect - B: B1 = Beriberi.

Incorrect - C: C = Scurvy.

Incorrect - D: D = Rickets.

Trick: "Vit A = A for eyes at night". Retinol = retinal function.

Source: NORCET-2023 | Nutritional Deficiencies

Tags: #PYQ #HighYield #Pediatrics #Nutrition #VitA #NightBlindness #NCLEX

[Pediatrics > Procedures > IM] High - VERIFIED [UNIQUE]

Q17. Which of the following is the most common site for intramuscular injection in infants?

- A) Gluteus maximus
- B) Vastus lateralis
- C) Deltoid
- D) Rectus femoris

Answer: B) Vastus lateralis

Explanation:

Correct - B: Infant <2yr = Vastus lateralis thigh. No major nerves/vessels, well-developed muscle. Avoid gluteal = sciatic risk.

Incorrect - A: Gluteus = after 3yr walking.

Incorrect - C: Deltoid = after 3yr, small volume.

Incorrect - D: Rectus femoris = not preferred.

Trick: "Baby Vastus = Very safe". 0-2yr = Vastus lateralis.

Source: NORCET-2023 | IM Injection Sites

Tags: #PYQ #HighYield #Pediatrics #Procedures #IM #VastusLateralis #NCLEX

[Pediatrics > Infectious > GI] High - VERIFIED [UNIQUE]

Q22. Which of the following is the most common cause of diarrhea in children?

- A) Rotavirus
- B) E. coli
- C) Salmonella
- D) Shigella

Answer: A) Rotavirus

Explanation:

Correct - A: Rotavirus = #1 cause severe dehydrating diarrhea <5yr. Vaccine available. Winter seasonality.

Incorrect - B: E.coli = Traveler's diarrhea.

Incorrect - C: Salmonella = Food poisoning.

Incorrect - D: Shigella = Bacillary dysentery, blood.

Trick: "Rota = Road to dehydration". Vaccine = Rotavac.

Source: NORCET-2023 | Pediatric Diarrhea

Tags: #PYQ #HighYield #Pediatrics #Infectious #Rotavirus #Diarrhea #NCLEX

[Pediatrics > Respiratory > Pneumonia] High - VERIFIED [UNIQUE]

Q23. Which of the following is the most common cause of pneumonia in children?

- A) Streptococcus pneumoniae
- B) Staphylococcus aureus
- C) Klebsiella pneumoniae
- D) Haemophilus influenzae

Answer: A) Streptococcus pneumoniae

Explanation:

Correct - A: S.pneumoniae = Pneumococcus = #1 bacterial CAP in children >3mo. PCV vaccine prevents.

Incorrect - B: Staph aureus = Post-viral, pneumatoceles.

Incorrect - C: Klebsiella = Aspiration, alcoholics.

Incorrect - D: H.influenzae = Hib vaccine reduced, still otitis/bronchitis.

Trick: "PneumoCoccus = Pneumonia Commonest". Vaccine = PCV.

Source: NORCET-2023 | Pediatric Pneumonia

Tags: #PYQ #HighYield #Pediatrics #Respiratory #Pneumonia #Pneumococcus #NCLEX

[Pediatrics > Neonatal > Meningitis] High - VERIFIED [UNIQUE]

Q24. Which of the following is the most common cause of meningitis in neonates?

- A) Group B Streptococcus
- B) E. coli
- C) Listeria monocytogenes
- D) All of the above

Answer: D) All of the above

Explanation:

Correct - D: Neonatal meningitis = GBS 50%, E.coli 20%, Listeria 5-10%. Early onset <7d = GBS/E.coli.
Late = Listeria.

Incorrect - A/B/C: Each alone incomplete, all are causes.

Trick: "Neonatal meningitis = GBS + E.coli + Listeria". Mom's vaginal flora.

Source: NORCET-2023 | Neonatal Sepsis

Tags: #PYQ #HighYield #Pediatrics #Neonatal #Meningitis #GBS #NCLEX

[Pediatrics > Infectious > Pertussis] High - VERIFIED [UNIQUE]

Q41. Which of the following is the most common cause of death in pertussis?

- A) Pneumonia
- B) Encephalopathy
- C) Apnea
- D) Seizures

Answer: A) Pneumonia

Explanation:

Correct - A: Secondary pneumonia = #1 death pertussis, especially <6mo. Whooping cough → exhaustion → infection.

Incorrect - B/C: Encephalopathy/Apnea = complications.

Trick: "Pertussis = Pneumonia kills". Whoop + cyanosis dangerous.

Source: NORCET-2023 | Pertussis

Tags: #PYQ #HighYield #Pediatrics #Infectious #Pertussis #Pneumonia #NCLEX

[Pediatrics > Neonatal > LBW] High - VERIFIED [UNIQUE]

Q6. The criteria for low birth weight baby:

- A) 2.5 kg
- B) 3 kg
- C) 1.5 kg
- D) 1 kg

Answer: A) 2.5 kg

Explanation:

Correct - A: LBW = <2500g at birth. VLBW <1500g, ELBW <1000g. WHO cutoff.

Incorrect - B: 3kg = normal.

Incorrect - C/D: 1.5kg = VLBW, 1kg = ELBW.

Trick: "LBW = Less than 2.5 Below Weight". <2.5kg needs NICU.

Source: NORCET-2025 | Neonatology WHO

Tags: #PYQ #HighYield #Pediatrics #Neonatal #LBW #NCLEX

[Pediatrics > Infectious > Meningitis] High - VERIFIED [UNIQUE]

Q45. Which of the following is the most common cause of meningitis in children?

- A) Streptococcus pneumoniae
- B) Haemophilus influenzae
- C) Neisseria meningitidis
- D) E. coli

Answer: B) Haemophilus influenzae

Explanation:

Correct - B: H.influenzae type b = #1 in unvaccinated <5yr. Hib vaccine reduced it. Now Strep pneumo commonest.

Incorrect - A: Strep pneumo = adults + post-vaccine era kids.

Incorrect - C: N.meningitidis = teens/young adults.

Incorrect - D: E.coli = neonates.

Trick: "Hib = Head infection in babies". Hib vaccine at 6,10,14wk.

Source: NORCET-2025 | Pediatrics

Tags: #PYQ #HighYield #Pediatrics #Infectious #Meningitis #Hib #NCLEX
