

Network level shifts in E/I balance during chronic epilepsy revealed by aperiodic EEG dynamics

Garima Chauhan, Kshitij Kumar, Deepti Chugh, Subramaniam Ganesh, Arjun Ramakrishnan

Indian Institute of Technology Kanpur, Uttar Pradesh, India

2025



Project page



Poster made with my Claude skill

Abstract

Chronic seizures progressively reshape brain-wide connectivity and excitation–inhibition (E/I) balance, but medication effects make these dynamics difficult to study in patients. Using a chronic epileptic Lafora disease (Epm2a^{−/−}) mouse model, we tracked how network-level E/I balance evolves across epileptogenesis and ictogenesis using the aperiodic exponent, a noninvasive EEG proxy for E/I ratio and beta-band oscillatory metrics. Laforin deficient (Epm2a^{−/−}) Lafora disease mice exhibited steeper baseline aperiodic slopes and elevated beta activity, indicating increased tonic inhibition. Isoflurane, a GABA-A receptor agonist, produced stronger spectral slope steepening in wild-type animals than in knockouts, reflecting altered GABAergic tone in Epm2a^{−/−} mice. During chemically induced seizures, aperiodic measures showed systematic variation across pre-ictal, ictal, and post-ictal periods, revealing phase-specific shifts between inhibitory and excitatory dominance. These results suggest that cortical E/I balance varies dynamically across seizure phases and severity. By resolving these transitions in real time, aperiodic EEG features offer a promising biomarker for seizure staging and vulnerability assessment. This work provides mechanistic insight into network-level adaptations in chronic epilepsy while avoiding pharmacological confounds.

Method

Using a chronic epileptic Lafora disease (Epm2a^{−/−}) mouse model, we tracked how network-level E/I balance evolves across epileptogenesis and ictogenesis using the aperiodic exponent, a noninvasive EEG proxy for E/I ratio and beta-band oscillatory metrics.

Results

Laforin deficient Lafora disease mice exhibited steeper baseline aperiodic slopes and elevated beta activity, indicating increased tonic inhibition. Isoflurane produced stronger spectral slope steepening in wild-type animals. Aperiodic measures showed systematic variation across pre-ictal, ictal, and post-ictal periods.

Introduction

Epilepsy is a dynamic network disorder in which recurrent seizures progressively reshape brain-wide structure, connectivity, and the balance between excitation and inhibition (E/I). Repeated seizures trigger long-lasting neuroplastic changes (Wang et al., 2025), destabilize local and global functional organization (Barrios-Martinez et al., 2025), and increase vulnerability to subsequent events (Liang et al., 2020). Disruptions in E/I balance are considered a central hallmark of epileptogenesis (Liu et al., 2023), reflected in alterations in the Hurst exponent in chronic temporal lobe epilepsy (Xie et al., 2024), dynamic E/I fluctuations across seizure phases (Žiburkus et al., 2013), and seizure-induced strengthening of excitatory synapses via BDNF signalling (Nasrallah et al., 2022). However, how E/I balance evolves across seizure stages and grades remains poorly understood, particularly in humans, where ethical constraints and anti-epileptic drugs (AEDs) obscure intrinsic circuit dy...

Conclusion

These results suggest that cortical E/I balance varies dynamically across seizure phases and severity. By resolving these transitions in real time, aperiodic EEG features offer a promising biomarker for seizure staging and vulnerability assessment.