

21-301 Assignment 05

Karan Sikka
ksikka@cmu.edu
November 1, 2013

1

We know that a maximal chain of $[n]$ contains $n + 1$ sets, and there is exactly one set of every size from 0 to n .

We know that the sizes of A_i strictly increase in the order they are given (from 1 to k). That is that A_1 is the smallest and A_k is the largest A_i .

We can construct all maximal chains of $[n]$ containing all A_i by the following method:

Let your chain contain all A_i . This is a chain given the subset relationship of all the A_i . Notice that this chain may not be maximal if there is not a set of every size from 0 to n . We know that we have sets of sizes $|A_1|, |A_2|, \dots, |A_k|$.

1. Now, add a maximal chain of 2^{A_1} .
2. Then find $|A_j| - |A_i|$ sets in $2^{[x]}$ of all sizes strictly between $(|A_i|, |A_j|)$ such that they form a chain, and add them to the chain.
3. Finally find $n - |A_k|$ sets in $2^{[x]}$ of all sizes strictly between $(|A_k|, n)$ such that they form a chain, and add them to the chain.

Note that by this method, you can form any maximal chain of $2^{[x]}$ which contains all the A_i . Now we count how many distinct outcomes there are of performing the above procedure.

1. There are $|A_1|!$ ways of doing step 1.
2. There are $\prod_{i,j|j-i=1} (|A_j| - |A_i|)!$ ways of doing step 2. This is because you can choose from $(|A_j| - |A_i|)$ elements for the first set you add, one fewer element for the next set, up til there is only one way to form the last set (then you have A_j).
3. There are $(n - |A_k|)!$ ways of doing step 3 by the same logic.

The product of those three quantities gives the answer.

2

We observe that the size of intersection of \mathcal{F} and any maximal chain of $2^{[n]}$, is at most 2, since if there were 3 sets in common with the chain, those 3 sets would form a chain which would break the constraint on \mathcal{F} given in the problem. With 2, you do not break the constraint.

The total number of sets in \mathcal{F} that also exist in some maximal chain is at most $2n!$ (two in a maximal chain, and there are $n!$).

We can also count this quantity by counting the number of maximal chains containing A for all $A \in \mathcal{F}$. This quantity is $\sum_{A \in \mathcal{F}} |A|!(n - |A|)!$, as shown in the proof of Sperner's Theorem in lecture.

Then, we do some algebra:

$$1 \geq \sum_{A \in \mathcal{F}} \frac{|A|!(n-|A|)!}{2n!} = \sum_{A \in \mathcal{F}} \frac{1}{2^{\binom{n}{\lfloor \frac{n}{2} \rfloor}}} = \frac{\mathcal{F}}{2^{\binom{n}{\lfloor \frac{n}{2} \rfloor}}}$$

Which implies that $\mathcal{F} \geq 2^{\binom{n}{\lfloor \frac{n}{2} \rfloor}}$ and since n is even, $\mathcal{F} \geq 2^{\binom{n}{\frac{n}{2}}}$.

3

4

5

6

7
