

02-512 Assignment 06

Karan Sikka

ksikka@cmu.edu

November 26, 2014

1

(a) The variables we are trying to find are the rate of infection, λ_1 and the rate of recovery, λ_2 .

We can run the CTMM as a simulation and compare it with real data. The output of the simulation will be sequence of states over time where each state will be (S_t, I_t, R_t) .

Let (Sr_t, Ir_t, Rr_t) be the real data. One possible objective function to minimize is

$$L(\lambda_1, \lambda_2) = \sum_{\text{real datapoints}} (S_t - Sr_t)^2 + (I_t - Ir_t)^2 + (R_t - Rr_t)^2$$

You can use steepest/gradient descent, Newton-Raphson's method, or a similar algorithm to find parameters yielding a local minimum. Rather than analytically computing the gradient, you'd have to approximate it using finite difference methods. Performance may be a concern, depending on how long the simulation has to run for.

(b) Let G be the growth rate, x_1 be the conc of nutrient 1 x_2 be the conc of nutrient 2.

$$G = \theta_1 x_1^2 + \theta_2 x_1 + \theta_3 x_2^2 + \theta_4 x_2 + \theta_5$$

Where $\vec{\theta}$ are the parameters we're trying to estimate.

Let $Gr(x_1, x_2)$ be the experimenally determined growth rate.

One possible objective function to minimize is

$$L(x_1, x_2) = \sum (Gr(x_1, x_2) - G(x_1, x_2))^2$$

You can use steepest descent once again, like in part a.

(c) Call parameters we are estimating, f_B and f_b , which are the frequencies of the B allele and b allele respectively.

Let B be the number of people observed with brown eyes, and b be the number of people observed with blue eyes.

The likelihood function is as follows:

$$Pr(B = B, b = b | p = p) =$$

2

(a) Let TTR be time to resistance.

TODO GAUSSIAN LIKELIHOOD

(b) TODO GAUSSIAN² LIKELIHOOD

(c) Do both and take the one with the smaller error?

(d) Metropolis

(e) Sampling vs solving

3

Given two points, the eqn of a line is derived as follows:

(Points given as $t_2 \ x_2 \ t_1 \ x_1$)

$$t = (x_2 - x_1)/(t_2 - t_1) x + b$$

$$t_1 - (x_2 - x_1)/(t_2 - t_1) x_1 = b$$

or

$$t_2 - (x_2 - x_1)/(t_2 - t_1) x_2 = b$$

$$t - t_2 = (x_2 - x_1)/(t_2 - t_1) * (x - x_2)$$

$$\text{if } 0 \leq t < 2, t - 2 = (5/2)(x - 5)$$

$$\text{if } 2 \leq t < 5, t - 5 = ((6-5)/(5-2))(x-6)$$

$$\text{if } 5 \leq t < 8, t - 8 = ((10-6)/(8-5))(x-10)$$

$$\text{if } 8 \leq t \leq 10, t - 10 = ((20-10)/(8-10))(x-20)$$

(b) TODO interp formula

(c) TODO interp formula + some deriv stuff.

4

(a)

If b_i is 0, there are no boojum on island i . Based on this observation, we note the following:

$$Pr(b_i = 0|f) = (1 - f)^{s_i}$$

$$Pr(b_i = 1|f) = 1 - Pr(b_i = 0|f) = 1 - (1 - f)^{s_i}$$

$$Pr(b|f) = \prod_{i=1}^n Pr(b_i = b_i)$$

(b)

$$f \approx \frac{1}{n} \sum_{i=1}^n \frac{y_i}{s_i}$$

(c)

expected number of people on island i

$$y_i = b_i * f * s_i$$