

Compatibility with Disk like Transfer Systems II

JMM 2025

David DeMark, Mike Hill, Yigal Kamel, Nelson Niu, **Kurt Stoeckl***,
Danika Van Niel and Guoqi Yan



Slides Part I



Slides Part II

Transfer Systems

Definition

Let \mathcal{O} be a binary relation on $Sub(G)$ refining \subset . Then, \mathcal{O} is said to be a G -transfer system if it is closed under

- conjugation,
- restriction, and
- composition.

Transfer Systems

Definition

Let \mathcal{O} be a binary relation on $Sub(G)$ refining \subset . Then, \mathcal{O} is said to be a G -transfer system if it is closed under

- conjugation,
- restriction, and
- composition.


Theorem (A.2 of [Rub21])

Let R be a binary relation on $Sub(G)$ refining \subset . Let $T(R)$ denote the closure of R under

- *conjugation, then*
- *restriction, and then*
- *composition.*

Then $T(R)$ is the smallest G -transfer system containing R .

Example



Theorem (A.2 of [Rub21])

Let R be a binary relation on $\text{Sub}(G)$ refining \subset . Let $T(R)$ denote the closure of R under

- conjugation, then
- restriction, and then
- composition.




Then $T(R)$ is the smallest G -transfer system containing R .

Disk like Transfer Systems

Definition

We say a transfer system \mathcal{O} is **disk like** if it admits the presentation $\mathcal{O} = T(D_G)$, where D_G is a set of relations/transfers with target G .


- Left, a non-disk like $C_{p^2,q}$ -transfer system.
- Mid, a non-disk like $C_{p^2,q}$ -transfer system.
- Right, a disk like $C_{p^2,q}$ -transfer system, its generators in solid green.






Saturated Transfer Systems

Definition

A transfer system \mathcal{O} is **saturated** if it satisfies the 2 out of 3 property.




- Left, a saturated $C_{p^2,q}$ -transfer system.
- Mid, a saturated $C_{p^2,q}$ -transfer system.
- Right, a non-saturated $C_{p^2,q}$ -transfer system.





Compatible Transfer Systems

Definition ([Cha24, Definition 4.6])

Let \mathcal{O}_a and \mathcal{O}_m be a pair of G -transfer systems such that $\mathcal{O}_m \subseteq \mathcal{O}_a$. We say $(\mathcal{O}_a, \mathcal{O}_m)$ are **compatible** if we can complete all squares of the form



with $e, r \in \mathcal{O}_m$ and $a \in \mathcal{O}_a$.



Maximal Compatible Transfer Systems


Proposition ([BH22])

If $(\mathcal{O}_a, \mathcal{O}_m)$ and $(\mathcal{O}_a, \mathcal{O}'_m)$ are both compatible, then $(\mathcal{O}_a, \mathcal{O}_m \vee \mathcal{O}'_m)$ is compatible.

Corollary

For a fixed transfer system \mathcal{O}_a ,

- there exists a maximal compatible transfer system \mathcal{O}_m , and*
- all other compatible transfers systems are sub-transfer systems of \mathcal{O}_m .*



Why?

Work including [BH15, GW18, BBR21, BP21, Rub21, BMO24, Cha24], provides the correspondences

N_∞ -operads

Additive Transfers

Multiplicative Norms

Bi-incomplete Transfers and Norms

Transfer Systems

Disk Like Transfer Systems

Saturated Transfer Systems

Compatible Transfer Systems

Why?

Work including [BH15, GW18, BBR21, BP21, Rub21, BMO24, Cha24], provides the correspondences

N_∞ -operads	Transfer Systems
Additive Transfers	Disk Like Transfer Systems
Multiplicative Norms	Saturated Transfer Systems
Bi-incomplete Transfers and Norms	Compatible Transfer Systems


Corollary

For a fixed transfer system \mathcal{O}_a ,

- there exists a maximal compatible transfer system \mathcal{O}_m , and
- all other compatible transfers systems are sub-transfer systems of \mathcal{O}_m .

Thus identifying the maximal compatible transfer system identifies all bi-incomplete/compatible multiplicative norms for a fixed additive transfer.

Maximal Compatible Pairs of Disk like Transfers of $C_{p,q}$



How?

In Part 1 with David, we saw that

Proposition (DHKNSVNY)

The maximal compatible transfer \mathcal{O}_m of \mathcal{O}_a is always saturated.

Is \mathcal{O}_m the 'maximal saturated sub-transfer system' of \mathcal{O}_a ?

How?

In Part 1 with David, we saw that

Proposition (DHKNSVNY)


The maximal compatible transfer \mathcal{O}_m of \mathcal{O}_a is always saturated.

Is \mathcal{O}_m the 'maximal saturated sub-transfer system' of \mathcal{O}_a ?




No!

- 1 There can exist multiple incomparable saturated transfer systems smaller than \mathcal{O}_a .
- 2 Saturated elements can exist in the open interval $(\mathcal{O}_m, \mathcal{O}_a)$.

There can be multiple incomparable saturated transfer systems smaller than \mathcal{O}_a




Saturated elements can exist in the open interval $(\mathcal{O}_m, \mathcal{O}_a)$



Computing \mathcal{O}_m

Worst case need to check A C B for all $e, r \in \mathcal{O}_m$ and $a \in \mathcal{O}_a$.



Computing \mathcal{O}_m

Worst case need to check $A \xrightarrow{e} C \xrightarrow{f} B$ for all $e, f \in \mathcal{O}_m$ and $a \in \mathcal{O}_a$.


Lemma (DHKNSVNY)

If $\mathcal{O}_a = \text{Comp}(\text{Rest}(\text{Conj}(B_a)))$ and $\mathcal{O}_m = \text{Comp}(\text{Rest}(\text{Conj}(B_m)))$.
Then, $(\mathcal{O}_a, \mathcal{O}_m)$ are compatible, if, and only if, $\mathcal{O}_m \subseteq \mathcal{O}_a$ and

$A \xrightarrow{e} C \xrightarrow{f} B$ for all $e, f \in \text{Rest}(\text{Conj}(B_m))$ and $a \in \text{Rest}(\text{Conj}(B_a))$.

Computing \mathcal{O}_m

Worst case need to check $A \xrightarrow{e} C \xrightarrow{f} B$ for all $e, f \in \mathcal{O}_m$ and $a \in \mathcal{O}_a$.



Lemma (DHKNSVNY)

If $\mathcal{O}_a = \text{Comp}(\text{Rest}(\text{Conj}(B_a)))$ and $\mathcal{O}_m = \text{Comp}(\text{Rest}(\text{Conj}(B_m)))$.
Then, $(\mathcal{O}_a, \mathcal{O}_m)$ are compatible, if, and only if, $\mathcal{O}_m \subseteq \mathcal{O}_a$ and

$A \xrightarrow{e} C \xrightarrow{f} B$ for all $e, f \in \text{Rest}(\text{Conj}(B_m))$ and $a \in \text{Rest}(\text{Conj}(B_a))$.




Also: It is possible to not conjugate one of the sets of generators!

Computing \mathcal{O}_m Via Its Complement

Proposition (DHKNSVNY)

The complement of the maximal compatible transfer system of \mathcal{O}_a satisfies

$$\mathcal{O}_m^c := \mathcal{O}_a \setminus \mathcal{O}_m = \{e \in \mathcal{O}_a : \exists r, r' \in \text{Res}(e), a \in \mathcal{O}_a, \nexists a' \in \mathcal{O}_a \text{ such that}$$



Idea:


- Delete the top left factors of all saturation failures in \mathcal{O}_a .
- Every occurrence of the pattern above deletes e, r and c .

Computing \mathcal{O}_m Via Its Complement

Proposition (DHKNSVNY)

The complement of the maximal compatible transfer system of \mathcal{O}_a satisfies

$$\mathcal{O}_m^c := \mathcal{O}_a \setminus \mathcal{O}_m = \{e \in \mathcal{O}_a : \exists r, r' \in \text{Res}(e), a \in \mathcal{O}_a, \nexists a' \in \mathcal{O}_a \text{ such that}$$




Proof sketch:

Computing \mathcal{O}_m Via Its Complement

Proposition (DHKNSVNY)

The complement of the maximal compatible transfer system of \mathcal{O}_a satisfies

$$\mathcal{O}_m^c := \mathcal{O}_a \setminus \mathcal{O}_m = \{e \in \mathcal{O}_a : \exists r, r' \in \text{Res}(e), a \in \mathcal{O}_a, \nexists a' \in \mathcal{O}_a \text{ such that}$$



Proof sketch:


- $e \in \mathcal{O}_m$ if, and only if, $(\mathcal{O}_a, T(e))$ is compatible.

Computing \mathcal{O}_m Via Its Complement

Proposition (DHKNSVNY)

The complement of the maximal compatible transfer system of \mathcal{O}_a satisfies

$$\mathcal{O}_m^c := \mathcal{O}_a \setminus \mathcal{O}_m = \{e \in \mathcal{O}_a : \exists r, r' \in \text{Res}(e), a \in \mathcal{O}_a, \nexists a' \in \mathcal{O}_a \text{ such that}$$



Proof sketch:


- $e \in \mathcal{O}_m$ if, and only if, $(\mathcal{O}_a, T(e))$ is compatible.
- Then check the compatibility of $(\mathcal{O}_a, T(e))$ with prior lemma.

Case: $\mathcal{O}_m^c = \emptyset$

Case: $\mathcal{O}_m^c = \emptyset$

Proposition (DHKNSVNY)

$\mathcal{O}_m^c := \mathcal{O}_a \setminus \mathcal{O}_m = \{e \in \mathcal{O}_a : \exists r, r' \in \text{Res}(e), a \in \mathcal{O}_a, \nexists a' \in \mathcal{O}_a \text{ such that}$



Proposition (DHKNSVNY)



A transfer system \mathcal{O}_a is self compatible, if, and only if, it is saturated.

Proof:

- If \mathcal{O}_a is saturated then $\mathcal{O}_m^c = \emptyset$.
- Every saturation failure of \mathcal{O}_a is in $\mathcal{O}_m^c = \emptyset$, thus \mathcal{O}_a is saturated.




Example: Computing \mathcal{O}_m in C_{p^2, q^2} using the complement

$$\mathcal{O}_m^c := \mathcal{O}_a \setminus \mathcal{O}_m = \{e \in \mathcal{O}_a : \exists r, r' \in \text{Res}(e), a \in \mathcal{O}_a, \nexists a' \in \mathcal{O}_a \text{ such that}$$






Example: Computing \mathcal{O}_m in C_{p^2, q^2} using the complement

$$\mathcal{O}_m^c := \mathcal{O}_a \setminus \mathcal{O}_m = \{e \in \mathcal{O}_a : \exists r, r' \in \text{Res}(e), a \in \mathcal{O}_a, \nexists a' \in \mathcal{O}_a \text{ such that}$$



Example: Computing \mathcal{O}_m in C_{p^2, q^2} using the complement

$$\mathcal{O}_m^c := \mathcal{O}_a \setminus \mathcal{O}_m = \{e \in \mathcal{O}_a : \exists r, r' \in \text{Res}(e), a \in \mathcal{O}_a, \nexists a' \in \mathcal{O}_a \text{ such that}$$



Some Questions and Future Directions

- How can we use the disk like assumption to aid in computing \mathcal{O}_m ?
- Can we use subset bounds on \mathcal{O}_m for faster computation?
- Can we compute all maximal compatible pairs for all disk like transfer systems of a fixed group G in a relatively efficient manner?
 - i.e. maybe it is 'hard' to compute \mathcal{O}_m for arbitrary \mathcal{O}_a ,
 - but we can induct to all $(\mathcal{O}_a, \mathcal{O}_m)$ from computing $(T(H \rightarrow G), \mathcal{O}_m)$?

Mentioned Sources I

- [BBR21] Scott Balchin, David Barnes, and Constanze Roitzheim. N_∞ -operads and associahedra. *Pacific J. Math.*, 315(2):285–304, 2021.
- [BH15] Andrew J. Blumberg and Michael A. Hill. Operadic multiplications in equivariant spectra, norms, and transfers. *Adv. Math.*, 285:658–708, 2015.
- [BH22] Andrew J. Blumberg and Michael A. Hill. Bi-incomplete Tambara functors. In *Equivariant topology and derived algebra*, volume 474 of *London Math. Soc. Lecture Note Ser.*, pages 276–313. Cambridge Univ. Press, Cambridge, 2022.
- [BMO24] Scott Balchin, Ethan MacBrough, and Kyle Ormsby. Composition closed premodel structures and the Kreweras lattice. *European J. Combin.*, 116:Paper No. 103879, 22, 2024.
- [BP21] Peter Bonventre and Luís A. Pereira. Genuine equivariant operads. *Adv. Math.*, 381:Paper No. 107502, 133, 2021.

Mentioned Sources II

- [Cha24] David Chan. Bi-incomplete Tambara functors as \mathcal{O} -commutative monoids. *Tunis. J. Math.*, 6(1):1–47, 2024.
- [GW18] Javier J. Gutiérrez and David White. Encoding equivariant commutativity via operads. *Algebr. Geom. Topol.*, 18(5):2919–2962, 2018.
- [Rub21] Jonathan Rubin. Detecting Steiner and linear isometries operads. *Glasg. Math. J.*, 63(2):307–342, 2021.