

Homotopy probs and other G -Operadic Structures¹

Kurt Stoeckl
supervised by
Marcy Robertson

University of Melbourne

AustMS 2023







¹Extending "Koszul Operads Governing Props and Wheeled Props" arXiv:2308.08718

G-Operadic

Idea: G -operadic structures model the composition of functions via graphs, where the edges of the graph are modelled by some group G .

| G | Operadic Family | | |
|-----------|-----------------|----------|------|
| | Operad | Properad | Prop |
| Non-Symm. | ... | ... | ... |
| Symmetric | ... | ... | ... |
| Braided | ... | ... | ... |
| Ribbon | ... | ... | ... |

Operadic Structures: Generalising the Graphs

| Example Operadic | Its Presentation | |
|------------------------------------|---|--|
| | Generators | Relations |
| The Operad Governing Ass. Algebras |  |  |
| The Properad Governing Bialgebras |  |  |
| The Prop Governing Hopf Algebras |  |  |

Trees \subset Connected Graphs \subset Graphs

G -Props: Varying G

Definition

- Symmetric props are symmetric monoidal categories,

in each case generated by a single object.

G-Props: Varying G

Definition

- Symmetric props are symmetric monoidal categories,
 - Non-symmetric props are non-symmetric monoidal categories,
 - Braided props (also called probs) are braided monoidal categories,
- in each case generated by a single object.


G-Props: Varying G

Definition

- Symmetric props are symmetric monoidal categories,
 - Non-symmetric props are non-symmetric monoidal categories,
 - Braided props (also called probs) are braided monoidal categories,
- in each case generated by a single object.

These are all instances of a more general definition called a G -prop.

$$G = (G_n, \gamma_n : G_n \rightarrow \mathbb{S}_n)_{n \in \mathbb{N}}$$




G-Props: Varying G

Definition

- Symmetric props are symmetric monoidal categories,
 - Non-symmetric props are non-symmetric monoidal categories,
 - Braided props (also called probs) are braided monoidal categories,
- in each case generated by a single object.

These are all instances of a more general definition called a G -prop.

$$G = (G_n, \gamma_n : G_n \rightarrow \mathbb{S}_n)_{n \in \mathbb{N}}$$



Claim: G -props can also be defined as algebras over operads!

Koszul Operads

Let \mathbb{P} be the operad, whose algebras are symmetric props in $\text{Vect}_{\mathbb{K}}$.

Theorem (Stoeckl [2023])

The operad \mathbb{P} is Koszul.

Koszul Operads

Let \mathbb{P} be the operad, whose algebras are symmetric props in $\text{Vect}_{\mathbb{K}}$.

Theorem (Stoeckl [2023])

The operad \mathbb{P} is Koszul.

A algebraic operad P is Koszul if, and only if, it has a quadratic model P_{∞} .

- model: $P_{\infty} \xrightarrow{f} P$, f is an epimorphism and quasi-isomorphism.
- quadratic: $P_{\infty} = (F(E), d)$ and $d(E) \rightarrow F(E)^2$
- quadratic \implies minimal

Why might we care?

Corollary

Algebras over \mathbb{P}_{∞} are homotopy-props.

Koszul Operads Governing Operadic Structures

Theorem

The operads governing all mainstream symmetric (and non-symmetric) operadic structures are all Koszul.

This is the culmination of many years of work by many authors.

- Non-symmetric Operads: [Van der Laan, 2003]
- Modular operads: [Ward, 2022]
- Operadic structures living on connected graphs:
 - [Kaufmann and Ward, 2023] with cubical Feynman categories.
 - [Batanin and Markl, 2023] with partial operads in operadic categories
- All operadic structures: [Stoeckl, 2023]

The Key Ideas in Extending the Theory

- ① Non-symmetric Operads.
- ② Symmetric connected operadic structures.
- ③ Disconnected operadic structures.

$1 \rightarrow 2$: Hide symmetric action in action of a groupoid (or equiv.).

$2 \rightarrow 3$: Use Gröbner bases methods instead of polytopes.

The Key Ideas in Extending the Theory

- ① Non-symmetric Operads.
- ② Symmetric connected operadic structures.
- ③ Disconnected operadic structures.

$1 \rightarrow 2$: Hide symmetric action in action of a groupoid (or equiv.).

$2 \rightarrow 3$: Use Gröbner bases methods instead of polytopes.

Remainder of this talk:

- Describe these two ideas.
- Outline why they also work for G -operadic structures.

→ Understand the operad governing props.


The Operad Governing Props

Generalised Graphs \approx Vertices + Flags/Half-Edges

Proposition (Specialisation of Lemma 14.2 Yau and Johnson [2015])

Let Gr^\uparrow be the set of strict isomorphism classes of directed, vertex labelled generalised graphs, with no directed cycles, and partition it by profiles.

$$Gr^\uparrow\left(\binom{n}{m}, \dots, \binom{n_k}{m_k}\right) := \{\gamma \in Gr^\uparrow : \gamma \text{ profile } \binom{n}{m}, v_i \text{ profile } \binom{n_i}{m_i}\}$$



The Operad Governing Props


Generalised Graphs \approx Vertices + Flags/Half-Edges

Proposition (Specialisation of Lemma 14.2 Yau and Johnson [2015])

Let Gr^\uparrow be the set of strict isomorphism classes of directed, vertex labelled generalised graphs, with no directed cycles, and partition it by profiles.

$$Gr^\uparrow\left(\begin{pmatrix} n \\ m \end{pmatrix}, \dots, \begin{pmatrix} n_k \\ m_k \end{pmatrix}\right) := \{\gamma \in Gr^\uparrow : \gamma \text{ profile } \begin{pmatrix} n \\ m \end{pmatrix}, v_i \text{ profile } \begin{pmatrix} n_i \\ m_i \end{pmatrix}\}$$

Then Gr^\uparrow with \circ_i given by graph substitution is a coloured operad in Set , whose algebras are symmetric props in Set .



The Operad Governing Props

Proposition (Specialisation of Lemma 14.4 Yau and Johnson [2015])

The operad Gr^\uparrow governing symmetric props in Set , can be enriched into the operad \mathbb{P} governing symmetric props in $Vect_{\mathbb{K}}$.

$$\mathbb{P}\left(\begin{smallmatrix} n \\ m_1, \dots, m_k \end{smallmatrix}\right) \cong \mathbb{K}\langle Gr^\uparrow\left(\begin{smallmatrix} n \\ m_1, \dots, m_k \end{smallmatrix}\right) \rangle,$$


$$k_1 \begin{array}{c} \textcircled{v_2} \\ \textcircled{v_1} \\ | \end{array} + k_2 \begin{array}{c} \textcircled{v_2} \\ \textcircled{v_1} \\ | \end{array} \in \mathbb{P}\left(\begin{smallmatrix} 0 \\ 2, 2 \end{smallmatrix}\right)$$

Towards a Quadratic Presentation

- We want to provide a quadratic presentation $\mathbb{P} \cong F(E)/\langle R \rangle$.


Towards a Quadratic Presentation

- We want to provide a quadratic presentation $\mathbb{P} \cong F(E)/\langle R \rangle$.
- Need equalities like **r** to hold formally,




Towards a Quadratic Presentation

- We want to provide a quadratic presentation $\mathbb{P} \cong F(E)/\langle R \rangle$.
- Need equalities like **r** to hold formally, **not** by def of graph subst.



Towards a Quadratic Presentation

- We want to provide a quadratic presentation $\mathbb{P} \cong F(E)/\langle R \rangle$.
- Need equalities like **r** to hold formally, **not** by def of graph subst.



Key Idea: Hide symmetric structure of graph in action of a groupoid

- Reconsider \mathbb{P} as a groupoid coloured operad.


The Free Full Nesting Operad

Proposition

Let E be the set of graphs in Gr^\uparrow with two vertices, partitioned as follows.

$$E = \left\{ \begin{array}{c} \text{graph 1} \\ \text{graph 2} \\ \text{graph 3} \end{array} \right\}.$$

The free operad $F(E)$ is equivalent to 'fully nested graphs' with \circ_i given by nested graph substitution.




A Quadratic Presentation of the Operad Governing Props

Proposition (Reformulation of Stoeckl [2023])

The groupoid coloured operad \mathbb{P} admits a quadratic presentation $\mathbb{P} \cong F(E)/\langle R \rangle$.

The quadratic relations R are,



Why does this work for all G -props?

Why does this work for all G -props?

- All G -operadic structures only differ in their equivariance axioms.

Why does this work for all G -props?


- All G -operadic structures only differ in their equivariance axioms.
 - Symmetric equivariance axioms hidden in action of a groupoid.
 - G -equivariance axioms in action of a groupoid.

Why does this work for all G -props?

- All G -operadic structures only differ in their equivariance axioms.
 - Symmetric equivariance axioms hidden in action of a groupoid.
 - G -equivariance axioms in action of a groupoid.
- Varying G , the operads governing G -props have 'same' presentation,
 - with different groupoid colouring!

An Informal Illustration for Braided Props

Governing operad \mathbb{P}_B can be described in terms of **braided** graphs.



$$\mathbb{P}_B \cong F(E_B)/\langle R_B \rangle$$

where

- E_B is the set of **braided** graphs with two vertices.
- $F(E_B)$ is the set of fully nested **braided** graphs.
- R_B is R re-interpreted with nested **braided** graphs.

Koszulness of Operads Governing G -Operadic Structures

Koszulness of Operads Governing G -Operadic Structures

Theorem (Stoeckl [2023])

Let $\mathcal{O} = F_{\Sigma}(\mathcal{X})/\langle \mathcal{G} \rangle$ be a \mathbb{V} -coloured operad, where $\text{Aut}(v)$ is finite for all $v \in \text{ob}(\mathbb{V})$. If $(\mathcal{O}^f)_$ admits a quadratic Groebner basis then \mathcal{O} is Koszul.*

Proposition

If G has a solvable word problem, then the operad \mathcal{O} governing each G -operadic structure satisfies: $(\mathcal{O}^f)_$ has a quadratic Groebner basis.*

Koszulness of Operads Governing G -Operadic Structures

Theorem (Stoeckl [2023])

Let $\mathcal{O} = F_{\Sigma}(\mathcal{X})/\langle \mathcal{G} \rangle$ be a \mathbb{V} -coloured operad, *where $\text{Aut}(v)$ is finite for all $v \in \text{ob}(\mathbb{V})$* . If $(\mathcal{O}^f)_*$ admits a quadratic Groebner basis then \mathcal{O} is Koszul.

Proposition

If G has a solvable word problem, then the operad \mathcal{O} governing each G -operadic structure satisfies: $(\mathcal{O}^f)_*$ has a quadratic Groebner basis.

Recall $G = (G_n, \gamma_n : G_n \rightarrow \mathbb{S}_n)_{n \in \mathbb{N}}$.

Corollary

The groupoid coloured operads governing all G -operadic structures are Koszul, *if G_n is a finite group for all $n \in \mathbb{N}$* .

What I'm Thinking On

- The operads governing all G -operadic structures have quadratic Groebner bases.
- Using this result, what is the nicest possible way to produce associated Koszul operads?
 - Can seek to extend groupoid coloured theory.
 - Can use coloured operads with quadratic-unary presentations.
 - Mirroring Dehling and Vallette [2021].


Later,

- Currently working with Marcy Robertson and Phil Hackney on homotopy coherent nerves between various models of symmetric ∞ -(wheeled)-prop(erad)s.
- These nerves could also extend nicely in the G -case.

Mentioned Sources

- M. Batanin and M. Markl. Koszul duality for operadic categories. *Compositionality*, 5(4):56, 2023. ISSN 2631-4444. arXiv:2105.05198.
- M. Dehling and B. Vallette. Symmetric homotopy theory for operads. *Algebr. Geom. Topol.*, 21(4):1595–1660, 2021. ISSN 1472-2747. doi: 10.2140/agt.2021.21.1595. URL <https://doi.org/10.2140/agt.2021.21.1595>. arXiv:1503.02701.
- R. M. Kaufmann and B. C. Ward. Koszul Feynman categories. *Proc. Amer. Math. Soc.*, 151(8):3253–3267, 2023. ISSN 0002-9939. doi: 10.1090/proc/16372. URL <https://doi.org/10.1090/proc/16372>. arXiv:2108.09251.
- K. Stoeckl. Koszul operads governing props and wheeled props. *arXiv:2308.08718*, 2023.
- P. Van der Laan. Coloured koszul duality and strongly homotopy operads. *arXiv preprint math/0312147*, 2003.
- B. C. Ward. Massey products for graph homology. *IMRN: International Mathematics Research Notices*, 2022(11), 2022.
- D. Yau and M. W. Johnson. *A foundation for PROPs, algebras, and modules*, volume 203. American Mathematical Soc., 2015.

What does it mean to have a quadratic Groebner basis?



- Every graph has a unique minimal shuffle tree monomial forming it.
- Every non-minimal tree monomial can be rewritten to the minimal tree via the relations of the operad.