Correspondence Chengxiang Fang cxfang@whu.edu.cn

Lysobacter ximonensis sp. nov., isolated from soil

Yang Wang, Jun Dai, Lei Zhang, Xuesong Luo, Yaowei Li, Gong Chen, Yali Tang, Ya Meng and Chengxiang Fang

China Center for Type Culture Collection (CCTCC), College of Life Sciences, Wuhan University, Wuhan 430072, PR China

A bacterial isolate, strain XM415^T, obtained from soil of Tibet in China was characterized using a polyphasic taxonomic approach. Strain XM415^T was aerobic, Gram-negative, gliding, rod-shaped, oxidase-negative and catalase-positive. Strain XM415^T showed the highest 16S rRNA gene sequence similarity with *Lysobacter niastensis* GH41-7^T (96.0 %). Ubiquinone Q-8 and branched fatty acids, such as iso- $C_{16:0}$ (24.0 %), iso- $C_{15:0}$ (22.6 %), iso- $C_{17:1}\omega 9c$ (6.7 %), iso- $C_{14:0}$ (6.1 %) and iso- $C_{11:0}$ 3-OH (5.2 %), were predominant in strain XM415^T as well as in all type strains of recognized *Lysobacter* species. The DNA G+C content of XM415^T was 63.5 mol%. The genotypic and phenotypic data show that strain XM415^T represents a novel species of the genus *Lysobacter*, for which the name *Lysobacter ximonensis* sp. nov. is proposed. The type strain is XM415^T (=CCTCC AB 207091^T =NRRL B-51263^T).

The genus Lysobacter was first proposed by Christensen & Cook (1978), and the type species is Lysobacter enzymogenes. Although the description of an antibiotic-producing species, 'Lysobacter lactamgenus', has been published (Ono et al., 1984; Kimura et al., 1996), its taxonomic position has not vet been confirmed and its name has not been validly published. The species of the genus Lysobacter are related to members of the genera Xanthomonas, Pseudoxanthomonas, Stenotrophomonas, Thermomonas and Xylella, containing ubiquinone Q-8 as the major respiratory quinone and having fatty acid profiles with a predominance of isobranched fatty acids. At the time of writing, the genus Lysobacter encompasses 13 species with validly published names: Lysobacter antibioticus, L. brunescens, L. enzymogenes, L. gummosus (Christensen & Cook, 1978), L. concretionis (Bae et al., 2005), L. daejeonensis, L. vangpyeongensis (Weon et al., 2006), L. koreensis (Lee et al., 2006), L. niabensis, L. niastensis (Weon et al., 2007), L. defluvii (Yassin et al., 2007), L. capsici (Park et al., 2008) and L. spongiicola (Romanenko et al., 2008). Members of the genus are strongly proteolytic and characteristically lyse a variety of micro-organisms (both Gram-negative and Grampositive bacteria), as well as nematodes (Christensen, 2005; Yassin et al., 2007), suggesting that they have a particular biological function in microbial ecosystems.

In this study, strain XM415 $^{\rm T}$ was isolated from a soil sample presented by Chaohua Liao (a Chinese geologist) from Ximo, at an altitude of 750 m in Tibet (most areas of Tibet are above 2000 m). The soil sample was suspended in sterilized water and diluted solutions were spread on 0.1 \times

The GenBank/EMBL/DDBJ accession number for the 16S rRNA gene sequence of strain XM415^T is EU237492.

tryptic soy broth (TSB; Difco) agar. The plates were incubated at 30 °C for 4 days, and strain XM415 T was subsequently isolated. Cellular morphology was determined by using light microscopy with 2-day-old cells. The gliding motility test was performed as described by Bowman (2000). Oxidase activity was evaluated via the oxidation of 1 % p-aminodimethylaniline oxalate. Catalase activity was determined by observation of bubble production after the application of 3 % (v/v) hydrogen peroxide solution. Tests were also made for hydrolysis of starch, CM-cellulose (0.1 %, w/v), chitin from crab shells (1 %, w/v), casein (5 %, w/v) and tyrosine (0.5 %, w/v) (Smibert & Krieg, 1994). The temperature range (5, 10, 15, 20, 28, 30 and 37 °C) and pH range (pH 4-11 at intervals of 1 pH unit) and the requirement for 0, 0.5, 1, 2, 3 and 5% NaCl (w/v) for growth were determined by using $0.1 \times TSB$ medium.

The API ZYM, API 20E and API 20NE systems (bioMérieux) were used to determine enzyme activities, fermentation tests and biochemical properties of strain XM415^T following the manufacturer's instructions. Utilization of some carbohydrates and organic acids by strain XM415^T was tested by using the Biolog Microbial Identification System according to the protocol provided by the manufacturer.

Genomic DNA was extracted with a Bacteria Genomic DNA Isolation kit (Shanghai Chaoshi bio Technologies Co. Ltd). The 16S rRNA gene was amplified by PCR with bacterial universal primers 27F (5'-GAGTTTGAT-CCTGGCTCAG-3') and 1527R (5'-AGAAAGGAGGTG-ATCCAGCC-3') (Rainey *et al.*, 1996), and the PCR products were sequenced by Invitrogen. Similarity searches with the derived sequence were done in the EzTaxon database (Chun *et al.*, 2007). Phylogenetic analysis was

performed by using MEGA, version 3.1 (Kumar et al., 2004), after multiple alignment of the data via CLUSTAL_X (Thompson et al., 1997). Distances were obtained using options according to Kimura's two-parameter model (Kimura, 1980) and clustering was performed by using the neighbour-joining method (Saitou & Nei, 1987). Bootstrap values from 1000 replications were used to determine the confidence level of the branches (Felsenstein, 1985).

Analysis of the 16S rRNA gene sequence of strain XM415^T resulted in a sequence of 1458 bp. 16S rRNA gene sequence similarities to recognized members of the genus *Lysobacter* were 96.0 % (*L. niastensis* GH41-7^T), 95.8 % (*L. enzymogenes* DSM 2043^T), 95.5 % (*L. brunescens* KCTC 12130^T), 95.4 % (*L. antibioticus* DSM 2044^T), 95.4 % (*L. gummosus* KCTC 12132^T) and 95.2 % (*L. capsici* YC5194^T), i.e. below the threshold for demarcating bacterial species (Stackebrandt & Goebel, 1994). No other recognized bacterial species showed more than 96.0 % 16S rRNA gene sequence similarity to the new isolate. As shown in the phylogenetic tree (Fig. 1), the novel isolate XM415^T forms a distinct subline within the genus *Lysobacter*, branching together with the type strain of *L. brunescens* by the




Fig. 1. Neighbour-joining phylogenetic tree, showing the position of strain XM415^T within the radiation of the order *Xanthomonadales*. The tree was based on a comparison of 16S rRNA gene sequences that were at least 90 % complete. Numbers at nodes are bootstrap percentages, based on 1000 resamplings. Bar, 0.005 sequence divergence.

Table 1. Cellular fatty acid profiles of members of the genus Lysobacter

Strains: 1, strain XM415^T; 2, *L. antibioticus* DSM 2044^T; 3, *L. brunescens* DSM 6979^T; 4, *L. capsici* KCTC 22007^T; 5, *L. concretionis* KCTC 12205^T; 6, *L. daejeonensis* DSM 17634^T; 7, *L. enzymogenes* DSM 2043^T; 8, *L. gummosus* DSM 6980^T; 9, *L. koreensis* KCTC 12204^T; 10, *L. niabensis* DSM 18244^T; 11, *L. niastensis* DSM 18481^T; 12, *L. spongiicola* JCM 14760^T; 13, *L. yangpyeongensis* DSM 17635^T. Data are expressed as percentages of total fatty acids and were taken from Weon *et al.* (2006), Park *et al.* (2008), Romanenko *et al.* (2008) and this study; –, <1% or not detected.

Fatty acid	1	2	3	4	5	6	7	8	9	10	11	12	13
C _{10:0} 3-OH	_	_	_	_	_	_	1.1	_	_	_	_	_	_
$iso-C_{11:0}$	3.7	3.1	5.9	2.3	5.7	3.7	3.4	3.8	5.3	6.4	4.1	9.5	4.3
iso-C _{11:0} 3-OH	5.2	8.0	7.2	3.8	6.9	6.0	6.6	9.7	9.0	9.3	8.0	15.5	5.5
C _{14:0}	1.7	1.1	_	1.9	_	_	1.0	_	_	_	_	_	_
iso-C _{14:0}	6.1	1.3	3.7	_	2.3	11.2	1.4	_	4.0	8.7	4.2	3.3	4.5
iso-C _{15:0}	22.6	24.9	19.6	23.3	33.6	13.1	20.5	25.2	12.5	12.7	21.9	23.0	14.5
iso-C _{15:1} at 5	_	1.0	_	_	_	_	_	1.7	4.4	2.4	1.6	_	3.1
iso-C _{15:1} F	1.8	_	1.7	_	3.2	3.2	_	_	_	_	_	_	_
anteiso-C _{15:0}	6.3	3.8	2.6	_	1.2	3.2	3.8	5.5	_	5.9	3.8	_	5.1
C _{16:0}	7.0	8.0	1.5	10.8	1.5	1.4	8.6	6.0	_	1.1	_	_	3.1
C _{16:1} ω7c alcohol	_	1.6	_	_	_	_	_	1.7	10.8	7.8	4.5	_	8.8
$C_{16:1}\omega 11c$	_	4.1	_	2.2	_	_	_	4.5	_	1.0	_	_	2.2
iso-C _{16:0}	24.0	10.3	23.5	_	20.4	33.7	13.8	5.7	26.3	23.7	23.3	32.5	27.5
iso-C _{16:1} H	_	_	1.5	_	_	2.6	_	_	2.1	1.0	1.3	_	1.1
C _{17:0} cyclo	_	7.24	_	_	1.9	_	6.2	1.0	_	_	_	_	_
iso-C _{17:0}	1.5	3.4	2.3	3.7	4.1	_	2.9	7.8	1.8	1.6	1.3	2.8	1.9
iso- $C_{17:1}\omega 9c$	6.7	6.4	15.5	9.3	15.1	6.7	4.7	12.2	16.7	10.0	10.9	_	6.7
anteiso-C _{17:0}	_	_	_	_	_	_	_	1.4	_	_	_	_	1.1
$C_{18:1}\omega 7c$	1.2	1.7	_	6.5	_	_	3.3	2.5	_	_	_	_	_
Unknown 11.799	_	2.0	_	_	_	_	1.5	1.8	_	_	1.4	_	_
Summed feature 3*	2.9	8.3	9.5	20.4	_	6.1	15.8	6.4	1.4	2.0	6.5	_	3.3

^{*}Summed feature 3 comprises iso- $C_{15:0}$ 2-OH and/or $C_{16:1}\omega$ 7c.

http://ijs.sgmjournals.org 787

Table 2. Differential phenotypic characteristics of strain XM415^T and type strains of species of the genus Lysobacter

Strains: 1, strain XM415^T; 2, *L. antibioticus* DSM 2044^T; 3, *L. brunescens* DSM 6979^T; 4, *L. capsici* KCTC 22007^T; 5, *L. concretionis* KCTC 12205^T; 6, *L. daejeonensis* DSM 17634^T; 7, *L. defluvii* DSM 18482^T; 8, *L. enzymogenes* DSM 2043^T; 9, *L. gummosus* DSM 6980^T; 10, *L. koreensis* KCTC 12204^T; 11, *L. niabensis* DSM 18244^T; 12, *L. niastensis* DSM 18481^T; 13, *L. spongiicola* JCM 14760^T; 14, *L. yangpyeongensis* DSM 17635^T. All data are from the present study. All strains are positive in API 20NE tests for gelatin hydrolysis and negative for indole production, glucose acidification, arginine dihydrolase, urease and assimilation of potassium gluconate, capric acid, adipic acid and phenylacetic acid.

Characteristic	1	2	3	4	5	6	7	8	9	10	11	12	13	14
Catalase/oxidase	+/-	+/+	+/+	+/+	+/+	+/+	+/+	+/+	+/+	+/-	+/+	+/+	+/+	-/+
Nitrate reduction	_	+	_	_	_	+	_	_	_	_	_	+	_	_
Aesculin hydrolysis	+	+	+	+	_	+	_	+	+	_	_	+	_	_
Assimilation of:														
D-Glucose	+	+	_	+	_	+	_	+	+	_	_	_	_	_
L-Arabinose	_	_	_	_	_	_	_	+	_	+	_	_	_	_
D-Mannose	+	+	_	+	_	_	_	+	+	_	_	_	_	_
D-Mannitol	_	_	_	_	_	_	+	+	_	+	_	_	_	_
N-Acetylglucosamine	+	+	_	_	_	_	_	+	+	_	_	+	_	_
Maltose	+	+	_	+	_	+	+	+	+	_	_	+	_	_
Malic acid	_	+	_	_	_	_	_	+	+	_	_	_	_	+
Trisodium citrate	_	_	_	_	_	_	_	+	_	+	_	_	_	_
Enzyme activities														
α-Galactosidase	_	_	_	+	_	_	_	_	_	_	_	_	_	_
α-Glucosidase	+	_	+	+	+	+	+	+	+	_	+	_	_	+
N -Acetyl- β -glucosaminidase	+	_	_	_	_	_	_	_	+	_	+	_	_	+
β -Glucosidase	_	+	+	+	_	_	_	+	+	_	_	_	_	_
β -Galactosidase	+	+	_	-	_	_	_	+	+	_	_	+	_	_

neighbour-joining method. The DNA G+C content of XM415^T was determined by HPLC (UltiMate 3000; Dionex) (Mesbah *et al.*, 1989) as 63.5 mol%.

The respiratory quinone system was extracted and determined by HPLC as described by Xie & Yokota (2003). Analysis of the respiratory lipoquinones indicated that the isolate contained ubiquinone-8 (Q-8), corresponding to the characteristic feature of members of the genus Lysobacter (Bae et al., 2005). Strain XM415^T was grown on R2A medium (Difco) for 48 h at 30 °C and cellular fatty acids were then analysed as methyl esters by GC (Agilent 6890N) according to the instructions of the Sherlock Microbial Identification System (MIDI). The major fatty acids (>5%) detected (percentages of the total cellular fatty acids) in strain XM415^T were iso-C_{16:0} (24.0%), iso- $C_{15:0}$ (22.6%), $C_{16:0}$ (7.0%), iso- $C_{17:1}\omega 9c$ (6.7%), iso- $C_{14:0}$ (6.1%) and iso- $C_{11:0}$ 3-OH (5.2%). The presence of iso- $C_{15:0}$, iso- $C_{16:0}$ and iso- $C_{17:1}\omega 9c$ as the major fatty acids is a characteristic of genera in the Xanthomonadaceae, including the genera Xanthomonas, Xylella Pseudoxanthomonas, Stenotrophomonas, Luteimonas (Assih et al., 2002; Roumagnac et al., 2004; Yang et al., 2005). A comparison of fatty acid profiles among related Lysobacter species is shown in Table 1.

The new isolate XM415^T exhibits characteristics consistent with those of the genus *Lysobacter*, but appears not to conform to any recognized species. The low level of 16S rRNA gene sequence similarity (<97%) among *Lysobacter*

species with validly published names indicates that their genomic DNA relatedness is less than 70 % (Stackebrandt & Goebel, 1994). In addition, some biochemical and physiological differences confirm that isolate XM415^T is different from recognized members of the genus *Lysobacter* (Table 2).

Based on the polyphasic taxonomic approach described here, we propose that strain XM415^T should be classified within a novel species of the genus *Lysobacter*, with the name *Lysobacter ximonensis* sp. nov.

Description of Lysobacter ximonensis sp. nov.

Lysobacter ximonensis (xi.mo.nen'sis. N.L. masc. adj. ximonensis pertaining to Ximo, a village in China, from where the type strain was isolated).

Cells are aerobic, Gram-negative, gliding and rod-shaped $(0.5 \times 1-3 \mu m)$. Colonies on $0.1 \times$ TSB agar are mucoid and yellow in colour. Growth occurs at 10-37 °C, 0-1 % (w/ v) NaCl and pH 5–10. Oxidase-negative and catalase-positive. Hydrolyses starch, casein, tyrosine and o-nitrophenyl β -D-galactopyranoside, but not chitin or CM-cellulose. Positive for alkaline phosphatase, leucine arylamidase, valine arylamidase, acid phosphatase, naphthol-AS-BI-phosphohydrolase, α -glucosidase and N-acetyl- β -glucosaminidase and weakly positive for esterase C4, esterase lipase C8, cystine arylamidase, α -chymotrypsin and β -galactosidase (API ZYM). Utilizes α -cyclodextrin, dextrin, glycogen, D-fructose,

α-D-glucose, maltose, D-mannose, L-alaninamide, L-alanyl glycine, L-glutamic acid, glycyl L-aspartic acid, glycyl L-glutamic acid, D-glucose 6-phosphate, cellobiose, pyruvic acid methyl ester, β -hydroxybutyric acid, L-alanine and L-aspartic acid, but not Tween 80 or other substances contained in the Biolog GN2 microplate. Other phenotypic characteristics are summarized in Table 2. Contains iso-C_{16:0}, iso-C_{15:0}, C_{16:0}, iso-C_{17:1} ω 9c, iso-C_{14:0} and iso-C_{11:0} 3-OH as major fatty acids and ubiquinone Q-8 as the respiratory quinone. The G+C content of the genomic DNA of the type strain is 63.5 mol%.

The type strain, $XM415^{T}$ (=CCTCC AB 207091^T =NRRL B-51263^T), was isolated from a soil sample taken from Ximo, Tibet, China.

Acknowledgements

This work was supported by the R & D Infrastructure and Facility Development Program from the Ministry of Science and Technology of the People's Republic of China (grant no. 2005DKA21208).

References

- Assih, E. A., Ouattara, A. S., Thierry, S., Cayol, J.-L., Labat, M. & Macarie, H. (2002). *Stenotrophomonas acidaminiphila* sp. nov., a strictly aerobic bacterium isolated from an upflow anaerobic sludge blanket (UASB) reactor. *Int J Syst Evol Microbiol* **52**, 559–568.
- Bae, H.-S., Im, W.-T. & Lee, S.-T. (2005). Lysobacter concretionis sp. nov., isolated from anaerobic granules in an upflow anaerobic sludge blanket reactor. Int J Syst Evol Microbiol 55, 1155–1161.
- **Bowman, J. P. (2000).** Description of *Cellulophaga algicola* sp. nov., isolated from the surfaces of Antarctic algae, and reclassification of *Cytophaga uliginosa* (ZoBell and Upham 1944) Reichenbach 1989 as *Cellulophaga uliginosa* comb. nov. *Int J Syst Evol Microbiol* **50**, 1861–1868.
- **Christensen, P. (2005).** Genus IV. *Lysobacter* Christensen and Cook 1978, 372^{AL}. In *Bergey's Manual of Systematic Bacteriology*, 2nd edn, vol. 2, part B, pp. 95–101. Edited by D. J. Brenner, N. R. Krieg, J. T. Staley & G. M. Garrity. New York: Springer.
- **Christensen, P. & Cook, F. D. (1978).** *Lysobacter*, a new genus of nonfruiting, gliding bacteria with a high base ratio. *Int J Syst Bacteriol* **28**, 367–393.
- Chun, J., Lee, J.-H., Jung, Y., Kim, M., Kim, S., Kim, B. K. & Lim, Y. W. (2007). EzTaxon: a web-based tool for the identification of prokaryotes based on 16S ribosomal RNA gene sequences. *Int J Syst Evol Microbiol* 57, 2259–2261.
- **Felsenstein, J. (1985).** Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* **39**, 783–791.
- **Kimura, M. (1980).** A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* **16**, 111–120.
- Kimura, H., Miyashita, H. & Sumino, Y. (1996). Organization and expression in *Pseudomonas putida* of the gene cluster involved in cephalosporin biosynthesis from *Lysobacter lactamgenus* YK90. *Appl Microbiol Biotechnol* **45**, 490–501.
- Kumar, S., Tamura, K. & Nei, M. (2004). MEGA3: integrated software for molecular evolutionary genetics analysis and sequence alignment. *Brief Bioinform* 5, 150–163.

- Lee, J. W., Im, W.-T., Kim, M. K. & Yang, D.-C. (2006). Lysobacter koreensis sp. nov., isolated from a ginseng field. Int J Syst Evol Microbiol 56, 231–235.
- Mesbah, M., Premachandran, U. & Whitman, W. B. (1989). Precise measurement of the G+C content of deoxyribonucleic acid by high-performance liquid chromatography. *Int J Syst Bacteriol* 39, 159–167.
- Ono, H., Nozaki, Y., Katayama, N. & Okazaki, H. (1984). Cephabacins, new cephem antibiotics of bacterial origin. I. Discovery and taxonomy of the producing organisms and fermentation. *J Antibiot (Tokyo)* 37, 1528–1535.
- Park, J. H., Kim, R., Aslam, Z., Jeon, C. O. & Chung, Y. R. (2008). *Lysobacter capsici* sp. nov., with antimicrobial activity, isolated from the rhizosphere of pepper, and emended description of the genus *Lysobacter*. *Int J Syst Evol Microbiol* 58, 387–392.
- Rainey, F. A., Ward-Rainey, N., Kroppenstedt, R. M. & Stackebrandt, E. (1996). The genus *Nocardiopsis* represents a phylogenetically coherent taxon and a distinct actinomycete lineage: proposal of *Nocardiopsiaceae* fam. nov. *Int J Syst Bacteriol* 46, 1088–1092.
- Romanenko, L. A., Uchino, M., Tanaka, N., Frolova, G. M. & Mikhailov, V. V. (2008). *Lysobacter spongiicola* sp. nov., isolated from a deep-sea sponge. *Int J Syst Evol Microbiol* 58, 370–374.
- Roumagnac, P., Gagnevin, L., Gardan, L., Sutra, L., Manceau, C., Dickstein, E. R., Jones, J. B., Rott, P. & Pruvost, O. (2004). Polyphasic characterization of xanthomonads isolated from onion, garlic and Welsh onion (*Allium* spp.) and their relatedness to different *Xanthomonas* species. *Int J Syst Evol Microbiol* 54, 15–24.
- Saitou, N. & Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4, 406–425.
- Smibert, R. M. & Krieg, N. R. (1994). Phenotypic characterization. In *Methods for General and Molecular Bacteriology*, pp. 607–654. Edited by P. Gerhardt, R. G. E. Murray, W. A. Woods & N. R. Krieg. Washington, DC: American Society for Microbiology.
- **Stackebrandt, E. & Goebel, B. M. (1994).** Taxonomic note: a place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in bacteriology. *Int J Syst Bacteriol* **44**, 846–849.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. (1997). The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 25, 4876–4882.
- Weon, H.-Y., Kim, B.-Y., Baek, Y.-K., Yoo, S.-H., Kwon, S.-W., Stackebrandt, E. & Go, S.-J. (2006). Two novel species, *Lysobacter daejeonensis* sp. nov. and *Lysobacter yangpyeongensis* sp. nov., isolated from Korean greenhouse soils. *Int J Syst Evol Microbiol* 56, 947–951.
- Weon, H.-Y., Kim, B.-Y., Kim, M.-K., Yoo, S.-H., Kwon, S.-W., Go, S.-J. & Stackebrandt, E. (2007). Lysobacter niabensis sp. nov. and Lysobacter niastensis sp. nov., isolated from greenhouse soils in Korea. Int J Syst Evol Microbiol 57, 548–551.
- Xie, C. H. & Yokota, A. (2003). Phylogenetic analysis of *Lampropedia hyalina* based on the 16S rRNA gene sequence. *J Gen Appl Microbiol* 49, 345–349.
- Yang, D.-C., Im, W.-T., Kim, M. K. & Lee, S.-T. (2005). Pseudoxanthomonas koreensis sp. nov. and Pseudoxanthomonas daejeonensis sp. nov. Int J Syst Evol Microbiol 55, 787–791.
- Yassin, A. F., Chen, W.-M., Hupfer, H., Siering, C., Kroppenstedt, R. M., Arun, A. B., Lai, W.-A., Shen, F.-T., Rekha, P. D. & Young, C. C. (2007). Lysobacter defluvii sp. nov., isolated from municipal solid waste. Int J Syst Evol Microbiol 57, 1131–1136.

http://ijs.sgmjournals.org 789