

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/268795407>

# *Lysobacter terrae* sp. Nov. isolated from *Aglaia odorata* rhizosphere soil

Article in *International Journal of Systematic and Evolutionary Microbiology* · November 2014

DOI: 10.1099/ijs.0.067397-0 · Source: PubMed

CITATIONS

18

READS

87

9 authors, including:



**Hien T.T. Ngo**

Kyung Hee University

69 PUBLICATIONS 778 CITATIONS

[SEE PROFILE](#)



**Juan Du**

Zhengzhou University of Light Industry

62 PUBLICATIONS 1,777 CITATIONS

[SEE PROFILE](#)



**Moochang Kook**

Anyang University

59 PUBLICATIONS 575 CITATIONS

[SEE PROFILE](#)



**Tae-Hoo Yi**

Kyung Hee University

274 PUBLICATIONS 4,666 CITATIONS

[SEE PROFILE](#)

## *Lysobacter terrae* sp. nov. isolated from *Aglaia odorata* rhizosphere soil

Hien T. T. Ngo,<sup>1†</sup> KyungHwa Won,<sup>1†</sup> Juan Du,<sup>1</sup> Heung-Min Son,<sup>1</sup> YongJin Park,<sup>1</sup> MooChang Kook,<sup>2</sup> Ki-Young Kim,<sup>1</sup> Feng-Xie Jin<sup>3</sup> and Tae-Hoo Yi<sup>1</sup>

Correspondence  
Tae-Hoo Yi  
drhoo@khu.ac.kr

<sup>1</sup>College of Life science, Kyung Hee University, 1 Seocheon, Kihung Yongin, Gyeonggi 446-701, Republic of Korea

<sup>2</sup>Department of Marine Biotechnology, Anyang University, Incheon 417-833, Republic of Korea

<sup>3</sup>College of Bio and Food Technology, Dalian Polytechnic University, Qinggong-yuan No. 1, Ganjingzi-qu, Dalia 116034, PR China

A Gram-stain negative, facultatively anaerobic, non-motile, rod-shaped bacterium, designated strain THG-A13<sup>T</sup>, was isolated from *Aglaia odorata* rhizosphere soil in Gyeonggi-do, Republic of Korea. Based on 16S rRNA gene sequence comparisons, strain THG-A13<sup>T</sup> had close similarity with *Lysobacter niabensis* GH34-4<sup>T</sup> (98.5%), *Lysobacter oryzae* YC6269<sup>T</sup> (97.9%) and *Lysobacter yangpyeongensis* GH19-3<sup>T</sup> (97.3%). Chemotaxonomic data revealed that strain THG-A13<sup>T</sup> possesses ubiquinone-8 (Q8) as the predominant isoprenoid quinone and iso-C<sub>15:0</sub>, iso-C<sub>16:0</sub> and iso-C<sub>17:1ω9c</sub> as the major fatty acids. The major polar lipids were phosphatidylethanolamine, phosphatidylglycerol and diphosphatidylglycerol. The G + C content was 66.3 mol%. The DNA–DNA relatedness values between strain THG-A13<sup>T</sup> and its closest phylogenetic neighbours were below 18.0%. These data corroborated the affiliation of strain THG-A13<sup>T</sup> to the genus *Lysobacter*. These data suggest that the isolate represents a novel species for which the name *Lysobacter terrae* sp. nov. is proposed, with THG-A13<sup>T</sup> as the type strain (=KACC 17646<sup>T</sup>=JCM 19613<sup>T</sup>).

The genus *Lysobacter* was first established by Christensen & Cook (1978), and later emended by Park *et al.* (2008). The species of the genus *Lysobacter* are characterized by a high G + C content (61.7–70.7%) and a lack of flagella (Lee *et al.*, 2006; Wei *et al.*, 2012). At the time of writing, the genus *Lysobacter* contains 26 species with validly published names (<http://www.bacterio.net/lysobacter.html>), with *Lysobacter enzymogenes* as the type species. Species of the genus *Lysobacter* have been mostly isolated from soil and water samples. Some species of the genus *Lysobacter* are potential biocontrol agents for plant fungal pathogens due to their antifungal and antimicrobial activities (Hashizume *et al.*, 2004; Park *et al.*, 2008; Wei *et al.*, 2012). In this study, a novel species, *Lysobacter terrae* sp. nov. THG-A13<sup>T</sup>, was characterized using a polyphasic taxonomic approach.

†These authors contributed equally to this paper.

Abbreviations: RP-HPLC, reversed phase HPLC; SIM, sulfide-indole-motility medium.

The NCBI GenBank accession number for the 16S rRNA gene sequence of strain THG-A13<sup>T</sup> is KF483861.

Four supplementary figures are available with the online Supplementary Material.

Strain THG-A13<sup>T</sup> was isolated from *Aglaia odorata* rhizosphere soil (in the spring of 2013) in Gyeonggi-do, South Korea (37° 30' N 127° 15' E). Approximately 1 g soil was suspended thoroughly in 10 ml sterile 0.85% NaCl (w/v; saline solution). Serially diluted samples were spread on R2A agar (Difco) and incubated at 28 °C for a week. Single colonies were purified by transferring to new R2A plates. One isolate, THG-A13<sup>T</sup>, was cultured routinely on R2A agar at 28 °C and stored in R2A broth suspension containing glycerol (25%, w/v) at –70 °C.


For the sequencing of the 16S rRNA gene, genomic DNA was obtained using a Solgent genomic DNA extraction kit and the 16S rRNA gene was amplified according to the methods of Weisburg *et al.* (1991). The 16S rRNA gene sequencing was performed by Solgent; sequences of related taxa were obtained from the EzTaxon-e server (<http://eztaxon-e.ezbiocloud.net>; Kim *et al.*, 2012) and GenBank database. Multiple alignments were performed using the program CLUSTAL X (Thompson *et al.*, 1997), followed by gap editions in the BioEdit program (Hall, 1999). The Kimura two-parameter model (Kimura, 1983) was used to calculate evolutionary distances. The neighbour-joining (Saitou & Nei, 1987) and the maximum-likelihood methods were used

to reconstruct phylogenetic trees, as implemented in MEGA 5.2 (Kumar *et al.*, 2008). The bootstrap values were calculated based on 1000 replications (Felsenstein, 1985).

The 16S rRNA gene sequence of strain THG-A13<sup>T</sup> (1430 bp) was analysed. Sequence similarities indicated that the closest relatives of strain THG-A13<sup>T</sup> were *Lysobacter niabensis* GH34-4<sup>T</sup> (98.5%), *Lysobacter oryzae* YC6269<sup>T</sup> (97.9%), *Lysobacter yangpyeongensis* GH19-3<sup>T</sup> (97.3%) and *L. enzymogenes* DSM 2043<sup>T</sup> (95.5%). Evidence of this relationship between strain THG-A13<sup>T</sup> and other recognized species of the genus *Lysobacter* was derived from the phylogenetic tree (Figs 1 and S1, available in the online Supplementary Material).

Gram-staining was determined using a bioMérieux Gram stain kit. Cell morphology was observed with a transmission electron microscope (JEOL; model JEM1010) at 11 000× magnification. Anaerobic growth was performed in R2A broth supplemented with thioglycollate [0.1% (w/v), Sigma]. Motility was tested in sulfide-indole-motility medium (SIM; Difco). Growth at different temperatures

(4, 10, 15, 20, 25, 28, 30, 37, 40 and 45 °C) was assessed after 7 days of incubation on R2A agar. Salt tolerance was tested in R2A broth containing 0–1% (w/v) NaCl (at 0.2% intervals), and 1–5% (w/v) NaCl (at 1.0% intervals). Growth at various pH values (pH 4–10, at intervals of 0.5 pH unit) was determined in R2A broth and evaluated by monitoring the OD<sub>600</sub> after 5 days of incubation. For the experiments on pH, two different buffers were used (final concentration 100 mM): acetate buffer for pH 4.0–6.5 and phosphate buffer for pH 7.0–10.0. Production of flexirubin-type pigments was determined by the reversible colour shift to red, purple or brown when yellow or orange colonies are covered with aqueous 20% (w/v) KOH solution (Fautz & Reichenbach, 1980). Oxidase and catalase activities were tested with 1% (w/v) *N*, *N*, *N*'-tetramethyl-1,4-phenylenediamine reagent and 3% (v/v) H<sub>2</sub>O<sub>2</sub>, respectively. Methyl red and Voges–Proskauer reactions were tested in Clark and Lubs medium (Scharlau). Tests for the degradation of starch [1% (w/v), Difco], casein [2% (w/v) skimmed milk, Oxoid], DNA (DNase agar, Oxoid), aesculin [0.1% (w/v) aesculin and



**Fig. 1.** Neighbour-joining phylogenetic tree based on 16S rRNA gene sequences showing the relationships of *Lysobacter terrae* sp. nov. THG-A13<sup>T</sup> with related species of the genus *Lysobacter*. Filled circles at the nodes indicate branches that were also recovered using the maximum-parsimony algorithm. Bootstrap values (expressed as percentages of 1000 replications) over 70% are shown at the branching points. *Rhodanobacter umsongensis* GR24-2<sup>T</sup> (FJ821731) was used as an outgroup. Bar, 0.005 substitutions per nucleotide position.

0.05 % (w/v) ferric citrate (Difco)], Tween 20 [1.0 % (w/v), Sigma], Tween 80 [1.0 % (w/v), Sigma], L-tyrosine [0.5 % (w/v), Sigma], CM-cellulose [0.1 % (w/v), Sigma] and chitin from crab shells [1.0 % (w/v), Sigma] were determined after 5 days of incubation at 28 °C. Growth on MacConkey agar (Oxoid), nutrient agar (NA, Oxoid) and tryptone soya agar (TSA, Oxoid) was also tested. In addition, API 20NE, ID 32 GN and API ZYM test kits (bioMérieux) were used according to manufacturer instructions to evaluate carbon source utilization and enzyme activities. Strains of the species *L. niabensis* KACC 11587<sup>T</sup>, *L. oryzae* KACC 14553<sup>T</sup>, *L. yangpyeongensis* KACC 11407<sup>T</sup> and *L. enzymogenes* KCTC 12131<sup>T</sup> were included in experiments, under the same laboratory conditions, to act as references for the investigation of biochemical properties of strain THG-A13<sup>T</sup>.

Strain THG-A13<sup>T</sup> was a Gram-stain negative, facultative anaerobic and rod-shaped bacterium. No gliding or flagella-induced motility was observed, but numerous thin fimbriae were present around the cells, often connecting the cells to each other (Fig. S2). The occurrence of long cells (0.40–0.45 × 1.5–5.1 µm) are characteristic of species of the genus *Lysobacter*. Strain THG-A13<sup>T</sup> was positive for the production of flexirubin-type pigments. Catalase activity was weakly positive, while oxidase activity was positive. The physiological characteristics of strain THG-A13<sup>T</sup> are summarized in the species description and a comparison of selected characteristics of strain THG-A13<sup>T</sup> and related type strains is given in Table 1.

In order to determine the G+C content and to perform DNA–DNA hybridization, genomic DNA of strain THG-A13<sup>T</sup> was prepared as described previously (Moore & Dowhan, 1995). The G+C content was analysed as described by Mesbah *et al.* (1989) using a reverse phase HPLC system. DNA–DNA hybridization was performed fluorometrically, according to the method developed by Ezaki *et al.* (1989), with photobiotin-labelled probes in microplate wells. DNA–DNA hybridization experiments were performed between strain THG-A13<sup>T</sup> and closely related type strains of species of the genus *Lysobacter*. The optimum hybridization temperature was 51.5 °C.

The DNA G+C content of strain THG-A13<sup>T</sup> was 66.3 mol%, which conforms to the expected range of G+C contents within the genus *Lysobacter*. The DNA–DNA relatedness values between strain THG-A13<sup>T</sup> and the other species of the genus *Lysobacter* were in the range of 13–18 % (*L. niabensis* KACC 11587<sup>T</sup>, 17.2 ± 0.5 %; *L. oryzae* KACC 14553<sup>T</sup>, 16.9 ± 0.5 %; *L. yangpyeongensis* KACC 11407<sup>T</sup>, 13.6 ± 0.9 %). These very low DNA relatedness values suggest that THG-A13<sup>T</sup> represents a novel species of the genus *Lysobacter* (Wayne *et al.*, 1987).

For fatty acid analysis, cells of strain THG-A13<sup>T</sup> and strains of the three most closely related species of the genus *Lysobacter* were harvested from R2A agar plates after incubation for 2 days at 28 °C. The cellular fatty acid profiles were prepared according to the protocol of the

Sherlock Microbial Identification System (MIDI), detected by GC (Hewlett Packard 6890) and analysed with the Sherlock Aerobic Bacterial Database (TSBA60) (Sasser, 1990). For the extraction of isoprenoid quinones and polar lipids of strain THG-A13<sup>T</sup>, cells were cultured on R2A broth for 3 days and freeze-dried after harvesting. The polar lipids were extracted (Minnikin *et al.*, 1977, 1984), and separated using two-dimensional TLC (Tindall, 1990). For the detection of polar lipids, TLC plates were sprayed with reagents: 5 % (v/v) molybdophosphoric acid was used for total polar lipids; 0.2 % (v/v) ninhydrin for amino lipids; molybdenum blue reagent for phospholipids and  $\alpha$ -naphthol sulphuric acid reagent for glycolipids. Isoprenoid quinones were analysed by RP-HPLC (reversed phase HPLC; Waters 2690 Alliance system) [solvent, methanol/2-propanol (7:5, v/v); flow rate, 1.0 ml min<sup>-1</sup>], as previously described (Collins & Jones, 1981; Hiraishi *et al.*, 1996; Tamaoka *et al.*, 1983).

The major cellular fatty acids (>10.0 %) of strain THG-A13<sup>T</sup> were iso-C<sub>15:0</sub> (18.6 %), iso-C<sub>16:0</sub> (16.4 %) and iso-C<sub>17:1ω9c</sub> (11.0 %), which was consistent with other species of the genus *Lysobacter*. A comparison of the fatty acid profiles of strain THG-A13<sup>T</sup> and selected closely related strains is shown in Table 2. The polar lipid profile of strain THG-A13<sup>T</sup> included phosphatidylethanolamine, phosphatidylmethylethanolamine, phosphatidylglycerol, diphosphatidylglycerol, unidentified phosphoglycolipids, an unidentified glycolipid, unidentified aminophospholipids, unidentified aminolipids, an unidentified phospholipid and an unidentified lipid (Fig. S3). Although the major polar lipids were similar to those found in other members of the genus *Lysobacter*, differences were found between selected closely related strains of species of the genus *Lysobacter* and the novel strain (Table 1, Fig. S4). Strain THG-A13<sup>T</sup> contained Q-8 as the major respiratory quinone, which is typical of the genus *Lysobacter* (Luo *et al.*, 2012; Wei *et al.*, 2012), with small amounts of Q-7 also present.

On the basis of 16S rRNA gene sequence analysis, fatty acid composition, G+C content and DNA–DNA relatedness values, it has been demonstrated that strain THG-A13<sup>T</sup> represents a novel species of the genus *Lysobacter*, for which the name *Lysobacter terrae* sp. nov. is proposed.

## Description of *Lysobacter terrae* sp. nov.

*Lysobacter terrae* (L. gen. n. *terrae* of the earth).

Cells are Gram-stain negative, facultatively anaerobic, non-motile, rod-shaped and long (0.40–0.45 × 1.5–5.1 µm). After culture on R2A agar for 3 days, colonies are circular, yellow, smooth and 1.0–1.5 mm in diameter. Growth occurs at 4–37 °C (optimum 28–30 °C), at pH 5.0–9.0 (optimum 7.0–7.5), and in 0–0.2 % (w/v) NaCl. Grows well on R2A agar, grows weakly on NA, but not on TSA and MacConkey agar. Catalase and oxidase activities are weakly positive and positive, respectively. Aesculin, casein, Tween 20, Tween 80, CM-cellulose and L-tyrosine are hydrolysed,

**Table 1.** Physiological characteristics and major polar lipids of *Lysobacter terrae* THG-A13<sup>T</sup> and related type strains of species of the genus *Lysobacter*

Strains: 1. THG-A13<sup>T</sup>; 2. *L. niabensis* KACC 11587<sup>T</sup>; 3. *L. oryzae* KACC 14553<sup>T</sup>; 4. *L. yangpyeongensis* KACC 11407<sup>T</sup>; 5. *L. enzymogenes* KACC 10127<sup>T</sup> (the type species of the genus *Lysobacter*). All data (except for the DNA G + C contents, which are taken from Christensen & Cook, 1978; Weon *et al.*, 2006, 2007; Aslam *et al.*, 2009) are from this study. All strains are non-motile and did not grow on MacConkey agar. All strains are negative for indole production, glucose acidification, arginine dihydrolase and the Voges–Prokauer test. All strains are positive for the hydrolysis of gelatin,  $\beta$ -galactosidase, aesculin and for the methyl red test. In API ZYM tests, all strains are positive for alkaline phosphatase, esterase (C4), esterase lipase (C8), lipase (C14), leucine arylamidase, valine arylamidase, cystine arylamidase, trypsin,  $\alpha$ -chymotrypsin, acid phosphatase, naphthol-AS-BI-phosphohydrolase and  $\alpha$ -glucosidase, but negative for  $\alpha$ -fucosidase. In API 20NE tests, all strains are positive for  $\beta$ -galactosidase,  $\beta$ -glucosidase, and protease and negative for D-glucose, L-arabinose, D-mannitol, potassium gluconate, adipic acid and phenylacetic acid. +, Positive; –, negative; w, weakly positive; ND, not detected; PE, phosphatidylethanolamine; PME phosphatidyl-N-methylethanolamine; PG, phosphatidylglycerol; DPG, diphosphatidylglycerol.

Characteristic	1	2	3	4	5
Growth temperature (°C)	4–37	5–37	15–42	15–40	10–40
Growth at pH	5.0–9.0	5.0–8.0	5.5–11.0	5.0–8.0	6.0–10.0
Salt tolerance at 1 % (w/v)	–	+	–	–	+
Oxidase	+	+	+	+	+
Catalase	w	+	+	–	+
Nitrate reduction	–	–	–	+	–
Facultative anaerobic growth	+	–	+	–	+
Assimilation of					
Capric acid	+	–	+	–	+
Malic acid	–	–	–	–	+
Trisodium citrate	–	–	–	–	+
N-Acetyl-glucosamine	–	–	–	–	+
D-Maltose	–	–	–	–	+
D-Mannose	–	–	–	–	w
Hydrolysis of					
DNA	–	w	–	–	–
Casein	+	–	+	+	+
CM-cellulose	+	–	–	+	+
L-tyrosine	+	–	+	+	+
Starch	–	–	–	–	+
Chitin	–	–	–	–	+
Tween 20	+	+	+	+	–
Tween 80	+	+	+	–	+
Urea	–	+	–	–	–
Enzyme activities					
$\alpha$ -Galactosidase	+	–	–	–	–
$\beta$ -Galactosidase	w	w	–	–	–
$\beta$ -Glucuronidase	–	w	–	–	–
$\beta$ -Glucosidase	w	+	–	w	–
N-Acetyl- $\beta$ -glucosaminidase	+	–	–	+	+
$\alpha$ -Mannosidase	w	–	–	w	–
DNA G + C content (mol%)	66.3	62.5	67.4	67.3	69.0
Major Polar lipids	DPG, PG, PE	DPG, PE	PG, PE, PME	DPG, PG, PE, PME	ND

but starch, chitin and DNA are not. The methyl red test is weakly positive, but the Voges–Proskauer test is negative. Positive for the flexirubin-type pigments test (the reversible colour shift from yellow to red). In API ZYM tests, positive results are obtained from alkaline phosphatase, esterase (C4), esterase lipase (C8), leucine arylamidase, valine arylamidase, cystine arylamidase, trypsin,  $\alpha$ -chymotrypsin, acid phosphatase, naphthol-AS-BI-phosphohydrolase,  $\alpha$ -galactosidase,  $\alpha$ -glucosidase and N-acetyl- $\beta$ -glucosaminidase;

weakly positive results are obtained from lipase (C14),  $\beta$ -galactosidase,  $\beta$ -glucosidase and  $\alpha$ -mannosidase; negative results are obtained from  $\beta$ -glucuronidase and  $\alpha$ -fucosidase. In API 20NE tests, nitrate reduction, indole production, glucose acidification, arginine dihydrolase, urease and assimilation of D-glucose, L-arabinose, D-mannose, D-mannitol, N-acetyl-glucosamine, D-maltose, potassium gluconate, adipic acid, malic acid, trisodium citrate, and phenylacetic acid is negative;  $\beta$ -galactosidase,  $\beta$ -glucosidase

**Table 2.** Cellular fatty acid profiles of strain THG-A13<sup>T</sup> and phylogenetically related species of the genus *Lysobacter*

Strains: 1. THG-A13<sup>T</sup>; 2. *L. niabensis* KACC 11587<sup>T</sup>; 3. *L. oryzae* KACC 14553<sup>T</sup>; 4. *L. yangpyeongensis* KACC 11407<sup>T</sup> and 5. *L. enzymogenes* DSM 2043<sup>T</sup>. Data for strains 1–4 are from this study, and data for strain 5 were taken from Aslam *et al.* (2009). Cells of all strains were cultured on R2A agar for 3 days at 28 °C. Fatty acids comprising less than 0.5 % are not listed. ND, Not detected; tr, trace (<1.0 %).

Fatty acid	1	2	3	4	5
Saturated					
C <sub>16:0</sub>	6.0	4.5	9.5	10.5	8.6
C <sub>18:0</sub>	1.2	1.0	1.7	2.2	ND
Unsaturated					
iso-C <sub>15:1</sub> ω9c	3.8	ND	2.6	2.6	ND
C <sub>16:1</sub> ω7c alcohol	7.7	7.8	2.8	6.8	tr
C <sub>16:1</sub> ω11c	2.0	1.0	1.6	2.1	tr
iso-C <sub>17:1</sub> ω9c	11.0	10.0	15.7	5.6	4.7
Branched-chain					
iso-C <sub>11:0</sub>	4.7	6.4	1.0	3.7	3.4
iso-C <sub>11:0</sub> 3OH	7.7	9.3	7.8	5.8	6.6
iso-C <sub>12:0</sub>	ND	1.3	tr	1.7	tr
iso-C <sub>14:0</sub>	3.2	8.7	2.6	4.6	1.4
iso-C <sub>15:0</sub>	18.6	12.7	19.6	12.9	20.5
anteiso-C <sub>15:0</sub>	5.9	5.9	4.2	4.6	3.8
iso-C <sub>16:0</sub>	16.4	23.7	16.9	23.9	13.8
iso-C <sub>17:0</sub>	3.8	1.6	12.3	1.9	2.9
*Summed feature 3	1.4	2.0	tr	2.9	15.8

\*Summed features represent groups of two or three fatty acids that could not be separated by GLC with the MIDI system. Summed feature 3 contained C<sub>16:1</sub>ω7c and/or C<sub>16:1</sub>ω6c in all strains except 5, where it contained C<sub>16:1</sub>ω7c and/or iso-C<sub>15:0</sub> 2OH.

and capric acid are positive; gelatin hydrolysis (protease) is weakly positive. The major fatty acids (>10 %) are iso-C<sub>15:0</sub>, iso-C<sub>16:0</sub> and iso-C<sub>17:1</sub>ω9c. The major polar lipids are phosphatidylethanolamine, phosphatidylglycerol and diphosphatidylglycerol. The predominant isoprenoid quinone is Q-8.

The type strain, THG-A13<sup>T</sup> (=KACC 17646<sup>T</sup>=JCM 19613<sup>T</sup>), was isolated from *Aglaia odorata* rhizosphere soil in Gyeonggi-do, Republic of Korea. The G+C content of the genomic DNA of the type strain is 66.3 mol%.

## Acknowledgements

This work was supported by the 2012 research program of a grant from Kyung Hee University in 2012 (20120596), Republic of Korea.

## References

Aslam, Z., Yasir, M., Jeon, C. O. & Chung, Y. R. (2009). *Lysobacter oryzae* sp. nov., isolated from the rhizosphere of rice (*Oryza sativa* L.). *Int J Syst Evol Microbiol* **59**, 675–680.

Christensen, P. & Cook, F. D. (1978). *Lysobacter*, a new genus of nonfruiting, gliding bacteria with a high base ratio. *Int J Syst Bacteriol* **28**, 367–393.

Collins, M. D. & Jones, D. (1981). Distribution of isoprenoid quinone structural types in bacteria and their taxonomic implication. *Microbiol Rev* **45**, 316–354.

Ezaki, T., Hashimoto, Y. & Yabuuchi, E. (1989). Fluorometric deoxyribonucleic acid-deoxyribonucleic acid hybridization in micro-dilution wells as an alternative to membrane filter hybridization in which radioisotopes are used to determine genetic relatedness among bacterial strains. *Int J Syst Bacteriol* **39**, 224–229.

Fautz, E. & Reichenbach, H. (1980). A simple test for flexirubin-type pigments. *FEMS Microbiol Ecol* **8**, 87–91.

Felsenstein, J. (1985). Confidence limit on phylogenies: an approach using the bootstrap. *Evolution* **39**, 783–791.

Hall, T. A. (1999). BioEdit: a user-friendly biological sequence alignment editor and analysis program for Windows 95/98/NT. *Nucl Acids Symp Ser* **41**, 95–98.

Hashizume, H., Hattori, S., Igarashi, M. & Akamatsu, Y. (2004). Tripropeptin E, a new tripropeptin group antibiotic produced by *Lysobacter* sp. BMK333-48F3. *J Antibiot (Tokyo)* **57**, 394–399.

Hiraishi, A., Ueda, Y., Ishihara, J. & Mori, T. (1996). Comparative lipoquinone analysis of influent sewage and activated sludge by high-performance liquid chromatography and photodiode array detection. *J Gen Appl Microbiol* **42**, 457–469.

Kim, O. S., Cho, Y. J., Lee, K., Yoon, S. H., Kim, M., Na, H., Park, S. C., Jeon, Y. S., Lee, J. H. & other authors (2012). Introducing EzTaxon-e: a prokaryotic 16S rRNA gene sequence database with phylotypes that represent uncultured species. *Int J Syst Evol Microbiol* **62**, 716–721.

Kimura, M. (1983). *The Neutral Theory of Molecular Evolution*. Cambridge: Cambridge University Press.

Kumar, S., Nei, M., Dudley, J. & Tamura, K. (2008). MEGA: a biologist-centric software for evolutionary analysis of DNA and protein sequences. *Brief Bioinform* **9**, 299–306.

Lee, J. W., Im, W. T., Kim, M. K. & Yang, D. C. (2006). *Lysobacter koreensis* sp. nov., isolated from a ginseng field. *Int J Syst Evol Microbiol* **56**, 231–235.

Luo, G., Shi, Z. & Wang, G. (2012). *Lysobacter arseniciresistens* sp. nov., an arsenite-resistant bacterium isolated from iron-mined soil. *Int J Syst Evol Microbiol* **62**, 1659–1665.

Mesbah, M., Premachandran, U. & Whitman, W. B. (1989). Precise measurement of the G+C content of deoxyribonucleic acid by high-performance liquid chromatography. *Int J Syst Bacteriol* **39**, 159–167.

Minnikin, D. E., Patel, P. V., Alshamaony, L. & Goodfellow, M. (1977). Polar lipid composition in the classification of *Nocardia* and related bacteria. *Int J Syst Bacteriol* **27**, 104–117.

Minnikin, D. E., O'Donnell, A. G., Goodfellow, M., Alderson, G., Athalye, M., Schaal, A. & Parleat, J. H. (1984). An integrated procedure for the extraction of bacterial isoprenoid quinones and polar lipids. *J Microbiol Methods* **2**, 233–241.

Moore, D. D. & Dowhan, D. (1995). Preparation and analysis of DNA. In *Current Protocols in Molecular Biology*, pp. 2–11. Edited by F. W. Ausubel, R. Brent, R. E. Kingston, D. D. Moore, J. G. Seidman, J. A. Smith & K. Struhl. New York: Wiley.

Park, J. H., Kim, R., Aslam, Z., Jeon, C. O. & Chung, Y. R. (2008). *Lysobacter capsici* sp. nov., with antimicrobial activity, isolated from the rhizosphere of pepper, and emended description of the genus *Lysobacter*. *Int J Syst Evol Microbiol* **58**, 387–392.

Saitou, N. & Nei, M. (1987). The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* **4**, 406–425.

- Sasser, M. (1990).** *Identification of bacteria by gas chromatography of cellular fatty acids*, MIDI Technical Note 101. Newark, DE: MIDI Inc.
- Tamaoka, J., Katayama-Fujiruma, Y. & Kuraishi, H. (1983).** Analysis of bacterial menaquinone mixtures by high performance liquid chromatography. *J Appl Bacteriol* **54**, 31–36.
- Thompson, J. D., Gibson, T. J., Plewniak, F., Jeanmougin, F. & Higgins, D. G. (1997).** The CLUSTAL\_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* **25**, 4876–4882.
- Tindall, B. J. (1990).** Lipid composition of *Halobacterium lacusprofundi*. *FEMS Microbiol Lett* **66**, 199–202.
- Wayne, L. G., Brenner, D. J., Colwell, R. R., Grimont, P. A. D., Kandler, O., Krichevsky, M. I., Moore, L. H., Moore, W. E. C., Murray, R. G. E. & other authors (1987).** International Committee on Systematic Bacteriology. Report of the ad hoc committee on reconciliation of approaches to bacterial systematics. *Int J Syst Bacteriol* **37**, 463–464.
- Wei, D. Q., Yu, T. T., Yao, J. C., Zhou, E. M., Song, Z. Q., Yin, Y.-R., Ming, H., Tang, S.-K. & Li, W.-J. (2012).** *Lysobacter thermophilus* sp. nov., isolated from a geothermal soil sample in Tengchong, southwest China. *Antonie van Leeuwenhoek* **102**, 643–651.
- Weisburg, W. G., Barns, S. M., Pelletier, D. A. & Lane, D. J. (1991).** 16S ribosomal DNA amplification for phylogenetic study. *J Bacteriol* **173**, 697–703.
- Weon, H. Y., Kim, B. Y., Baek, Y. K., Yoo, S. H., Kwon, S. W., Stackebrandt, E. & Go, S. J. (2006).** Two novel species, *Lysobacter daejeonensis* sp. nov. and *Lysobacter yangpyeongensis* sp. nov., isolated from Korean greenhouse soils. *Int J Syst Evol Microbiol* **56**, 947–951.
- Weon, H. Y., Kim, B. Y., Kim, M. K., Yoo, S. H., Kwon, S. W., Go, S. J. & Stackebrandt, E. (2007).** *Lysobacter niabensis* sp. nov. and *Lysobacter niastensis* sp. nov., isolated from greenhouse soils in Korea. *Int J Syst Evol Microbiol* **57**, 548–551.