




Lysobacter prati sp. nov., isolated from a plateau meadow sample

Bao-Zhu Fang · Yuan-Guo Xie · Xing-Kui Zhou · Xiao-Tong Zhang ·
Lan Liu · Jian-Yu Jiao · Min Xiao · Wen-Jun Li 

Received: 6 August 2019 / Accepted: 9 January 2020 / Published online: 24 January 2020
© Springer Nature Switzerland AG 2020

Abstract A novel proteobacterial strain designated SYSU H10001^T was isolated from a soil sample collected from plateau meadow in Hongyuan county, Sichuan province, south-western China. The taxonomic position of the strain was investigated using a polyphasic approach. On the basis of 16S rRNA gene sequence similarities and phylogenetic analysis, strain SYSU H10001^T was most closely related to *Lysobacter soli* KCTC 22011^T (98.6%, sequence similarity) and *Lysobacter panacisoli* JCM 19212^T (98.2%). The prediction result of secondary metabolites based on genome shown that the strain SYSU H10001^T contained 3 clusters of bacteriocins, 1 cluster of non-ribosomal peptide synthetase, 1 cluster of type 1 polyketide synthase and 1 cluster of arylpolyene. In

addition, the major isoprenoid quinone was Q-8 and the major fatty acids were identified as iso-C_{15:0}, iso-C_{17:0} and Summed feature 9. The polar lipids contained diphosphatidylglycerol, phosphatidylglycerol, phosphatidylethanolamine, and three unidentified phospholipids. The genomic DNA G + C content of strain SYSU H10001^T was 66.5% (genome). On the basis of phenotypic, genotypic and phylogenetic data, strain SYSU H10001^T represents a novel species of the genus *Lysobacter*, for which the name *Lysobacter prati* sp. nov. is proposed. The type strain is SYSU H10001^T (= KCTC 72062^T = CGMCC 1.16662^T).

Keywords *Lysobacter prati* sp. nov. · Secondary metabolites · Plateau grassland · Polyphasic taxonomy

Electronic supplementary material The online version of this article (<https://doi.org/10.1007/s10482-020-01386-6>) contains supplementary material, which is available to authorized users.

B.-Z. Fang · Y.-G. Xie · X.-T. Zhang · L. Liu ·
J.-Y. Jiao · M. Xiao · W.-J. Li (✉)
State Key Laboratory of Biocontrol and Guangdong Key
Laboratory of Plant Resources, School of Life Sciences,
Sun Yat-Sen University, Guangzhou 510275, People's
Republic of China
e-mail: liwenjun3@mail.sysu.edu.cn

X.-K. Zhou
State Key Laboratory for Conservation and Utilization of
Bio-Resources in Yunnan, Yunnan University,
Kunming 650091, People's Republic of China

Introduction

The genus *Lysobacter* was first proposed by Christensen and Cook (1978) as the type genus of the family *Lysobacteraceae* of the order *Lysobacterales* in the phylum *Proteobacteria* and was subsequently emended by Park et al. (2008). Members of this genus are generally characterized as Gram-staining negative, aerobic, rod-shaped with variable motility and chemoorganotrophic and have high DNA G + C contents (61–70 mol %) (de Bruijn et al. 2015; Hernández and Fernández 2017). Colonies are highly

muroid, cream-coloured, pink, or yellow–brown; many strains produce a brown, water-soluble pigment. The genus *Lysobacter* contains ubiquinone 8 (Q-8) as the major respiratory quinone and shows a predominance of iso-branched fatty acids (Weon et al. 2007; Srinivasan et al. 2010; Wei et al. 2012). Strains of this genus have been isolated from various environmental niches such as soil, sediments, aquatic habitats and rhizosphere (Fukuda et al. 2013; Choi et al. 2014; Lin et al. 2015; Chen et al. 2016; Siddiqi and Im 2016). At the time of writing, the genus *Lysobacter* comprises 56 species with validly published names (<https://www.namesforlife.com/10.1601/tx.2240>), of which six novel species were published recently (Chhetri et al. 2019; Kim et al. 2019; Luo et al. 2019; Xiao et al. 2019; Zhang et al. 2019). Furthermore, some species from the genus of *Lysobacter* has been seen as a high-efficiency producer of bioactive natural products including new antibiotics from species such as *L. antibioticus* 76, *L. capsici* 55, *L. enzymogenes* C3, *L. antibioticus* ATCC 29479^T that contain more secondary metabolite gene clusters than others in the genus. The secondary metabolites such as cyclodepsipeptides, cyclic lipodepsipeptides, cephem-type β -lactams and polycyclic tetramate macrolactams (PTM) have been reported from members of this genus (Xie et al. 2012; Panthee et al. 2016). During a study of the diversity of bacteria of the plateau grassland environment, strain SYSU H10001^T was isolated from a soil sample and its taxonomic status was investigated using a polyphasic taxonomy approach.

Materials and methods

Isolation and culture conditions

The soil sample used for isolation of strain SYSU H10001^T was collected from a meadow located in Hongyuan county (32° 82' N, 105° 58' E), Sichuan province during a trip in 2017. Sampling was done using a sterile spoon and the sample collected into a sterile sampling bag. It was then transported back to the laboratory under ambient condition and stored at 4 °C. Isolation of *Lysobacter* was done using the standard dilution plate method on Reasoner's 2A agar (Reasoner and Geldreich 1985). The colonies of strain SYSU H10001^T were obtained after incubation for 1 week at 28 °C. Selected colonies were then purified

on YIM 38 agar and maintained on YIM 38 agar slants at 4 °C and as glycerol suspensions (20%, w/v) at – 80 °C. The experimental control strains *Lysobacter panacisoli* JCM 19212^T and *Lysobacter soli* KCTC 22011^T were provided by Japan Collection of Microorganisms (JCM) and the Korean Collection for Type Cultures (KCTC), respectively. All the strains were maintained routinely on YIM 38 medium (28 °C, 5 days). Biomass of strain SYSU H10001^T and the experimental control strains for chemotaxonomic and molecular investigations were harvested from cultures grown on YIM 38 medium (28 °C, 5 days).

Phenotypic characterisation

Colony morphology of strain SYSU H10001^T was observed from cultures grown on YIM 38 medium. Cell morphology was observed by transmission electron microscopy (JEM-1400FLASH) with 3-day-old cultures grown on YIM 38 medium. Gram's reaction was determined by Solarbio's Gram staining kit (China) as per the manufacturer's instructions. Growth at various NaCl concentrations (up to 5%, w/v) and at different temperatures (0, 4, 10, 15, 28, 30, 37, 40, 45 °C) were examined on YIM 38 agar plates, while the pH range for growth (4.0–12.0, at intervals of 1 pH unit prepared using the buffer system as described by Xu et al. 2005) were tested for 4 weeks by culturing the strains in YIM 38 broth (Jiang et al. 2016). Activities of oxidase, catalase, and urease, gelatin liquefaction, milk coagulation and peptonisation, nitrate reduction, H₂S production, degradation of Tweens 20, 40, 60 and 80, starch and cellulose hydrolysis were investigated according to the conventional procedures described by Gordon et al. (1974) and Williams et al. (1989). Other enzyme activities and biochemical characteristics were determined by using API ZYM kits (BioMérieux) according to the manufacturer's instructions. Carbon-source utilization tests were performed according to the methods described by Shirling and Gottlieb (1966) and Athalye et al. (1985) Using modified basal medium recommended by Pridham and Gottlieb (1948), sole nitrogen source utilization was observed as described by Nie et al. (2012). Other physiological and biochemical characteristics were assessed by using the media and methods described by Gordon et al. (1974).

Chemotaxonomy

Chemotaxonomic characteristics of strain SYSU H10001^T and the experimental control strain were determined under similar conditions. Menaquinones were extracted from lyophilized cells as described by Collins et al. (1977) and Minnikin et al. (1984), and the extracts were purified and analyzed by HPLC (Kropenstedt 1982; Tamaoka et al. 1983). For analysis of fatty acids, the three strains were grown on tryptic soy agar for 3 days at 28 °C. The cellular fatty acids were extracted, methylated and analyzed following the instructions of Microbial Identification System (MIDI) (Sherlock Version 6.1; MIDI database: TSBA6) (Sasser 1990). Polar lipids were extracted as the method described by Bligh and Dyer 1959, and the individual polar lipids separated by two-dimensional TLC on silica gel G₆₀ plates (Merck; Germany), and the profile was identified using the described procedures (Minnikin et al. 1979; Collins and Jones 1980; Minnikin et al. 1984).

Molecular characterisation

Genomic DNA was extracted, and the 16S rRNA gene PCR-amplified and sequenced as described by Li et al. (2007). The amplicons were purified using a PCR purification kit (Sangon Biotech, China). The 16S rRNA gene sequences obtained were assembled using the SeqMan program (DNASTar software) and compared with the corresponding sequences of cultured species in the EzBioCloud web service (Yoon et al. 2017a) and the NCBI database by using BLAST search (Altschul et al. 1990). To determine the phylogenetic relationships of strain SYSU H10001^T and members of the genus *Lysobacter*, multiple alignments of their 16S rRNA gene sequences were performed using the CLUSTAL_X software package (Thompson et al. 1997). Phylogenetic and molecular evolutionary analyses were performed using the software package MEGA version 7.0 (Kumar et al. 2016). Phylogenetic dendrograms were generated with maximum-likelihood (Felsenstein 1981), neighbour-joining (Saitou and Nei 1987) and maximum-parsimony (Fitch 1971) algorithms. Kimura's two-parameter model (Kimura 1980, 1985) was used to calculate the evolutionary distances in the neighbour-joining and maximum-likelihood phylogenetic dendrograms. Bootstrap analysis was used to evaluate the

topology of each tree with 1000 replications (Felsenstein 1985).

Whole-genome sequencing of strain SYSU H10001^T and experimental control strain (*L. panacisoli* JCM 19212^T) were performed using paired-end sequencing method with Hiseq X platform (Illumina, San Diego, CA, USA) at Beijing Genomics Institute (Beijing, China). Reads of each data set were filtered, and high-quality paired-end reads assembled using Spades (Harrison and Strulo 2000). Contigs, with length greater than 500 bp, were kept for gene prediction by applying Prodigal (Hyatt et al. 2010). The predicted coding sequences of each genome were translated and annotated using the KEGG, COG and Pfam databases. For generation of phylogenomic tree, 16 syntenic genes that have been shown to undergo limited lateral gene transfer (*rpL2*, *rpL3*, *rpL4*, *rpL5*, *rpL6*, *rpL14*, *rpL15*, *rpL16*, *rpL18*, *rpL22*, *rpL24*, *rpS3*, *rpS8*, *rpS10*, *rpS17*, and *rpS19*) of the selected genomes of the genus *Lysobacter* in the NCBI database were extracted using AMPHORA2 (Wu and Scott 2012). Sequences of each of the marker genes were aligned separately using MUSCLE (Edgar 2004). Alignments were concatenated by using perl script (<https://github.com/nylander/catfasta2phyml>), prior to the generation of the tree. Poorly aligned regions were removed from the datasets using Gblocks (Castresana 2000). The phylogenomic tree was generated with software program RAXML (Stamatakis 2014) using the default parameters and visualized using the online Interactive Tree of Life program v.4.2 (Letunic and Bork 2019). The presence of gene clusters encoding secondary metabolites was predicted by using the antiSMASH 5.0 database (Blin et al. 2019). Furthermore, values of average nucleotide identity (ANI) were calculated by using the ANI Calculator (Yoon et al. 2017b).

Results and discussion

Phenotypic characteristics

Cells of strain SYSU H10001^T stained Gram-negative and were aerobic, non-motile, and rod-shaped (Supplementary Fig. S1). Colonies on YIM 38 medium were observed to be circular, convex, opaque and yellow in colour. No soluble pigments and melanin production were observed on the above tested media.

Detailed physiological and biochemical characteristics of strain SYSU H10001^T are summarised in the species description, and characteristics that differentiate strain SYSU H10001^T from its closely related type strains are listed in Table 1.

Molecular characteristics

Pairwise comparison of the 16S rRNA gene sequence of strain SYSU H10001^T (GenBank accession number MN181427) with the corresponding 16S rRNA gene sequences in the EzBioCloud database indicated that the strain shared 16S rRNA gene sequence similarities of 98.6% and 98.2% with *L. soli* KCTC 22011^T and *L. panacisoli* JCM 19212^T, respectively. In the

maximum-likelihood phylogenetic tree based on 16S rRNA gene sequences, strain SYSU H10001^T formed a close clade with the above two strains (Fig. 1). This relationship was also supported in the phylogenetic trees generated with maximum-parsimony and neighbour-joining methods (Supplementary Figs. S3 and S4).

The genome of strain SYSU H10001^T generated 2.58 Gb raw data of which 2.36 Gb was clean data following quality-filtering. The de novo read assembly produced 48 contigs with an N₅₀ size of 420,681 bp. The high-quality assembled draft genome sequence consisted of 4,127,215 bp with a DNA G + C content of 66.5% which is within the reported range of DNA G + C contents for species of the genus *Lysobacter*.

Table 1 Differential phenotypic characteristics between strain SYSU H10001^T and its closely related type strains of the genus *Lysobacter*

| Characteristic | 1 | 2 | 3 |
|---|-------------|-------------|------------------|
| Cell morphology | Short rods | Rods, cocci | Short rods |
| Temperature range for growth (°C) | 4–37 | 4–40 | 4–40 |
| NaCl tolerance (% w/v) | 0–1.5 | 0–1.5 | 0–2 |
| <i>Hydrolysis of</i> | | | |
| Tween 40 | + | – | – |
| Tween 60 | + | – | – |
| Tween 80 | + | – | – |
| Starch | – | – | + |
| <i>Assimilation as sources of carbon and nitrogen</i> | | | |
| D-Arabinose | – | – | + |
| Gentiobiose | – | – | + |
| D-Cellobiose | + | – | – |
| D-Maltose | + | – | + |
| D-Sorbitol | – | – | + |
| D-Trehalose | – | – | + |
| D-Xylose | – | – | + |
| L-Arginine | + | – | – |
| L-Cystine | + | – | – |
| L-Glutamic | + | – | – |
| L-Histidine | + | – | – |
| L-Lysine | + | + | – |
| L-Phenylalanine | + | – | – |
| <i>API ZYM</i> | | | |
| Chymotrypsin | + | + | – |
| Esterase lipase (C8) | + | + | – |
| Valine arylamidase | + | + | – |
| Chymotrypsin | + | + | – |
| α-Glucosidase | + | – | – |
| Major polar lipids | DPG, PE, PG | DPG, PE, PG | DPG, PE, PME, PG |

Strains: (1) SYSU H10001^T; (2) *L. panacisoli* JCM 19212^T; (3) *L. soli* KCTC 22011^T. All data are from this study unless indicated otherwise
+ positive; – negative; w weakly positive. ND not determined




Fig. 1 Maximum-likelihood phylogenetic tree showing the relationship between strain SYSU H10001^T and its closest relatives. Asterisks indicate branches that were also recovered using the maximum-parsimony and neighbour-joining methods.

Bootstrap values (expressed as percentages of 1000 replications) of above 50% are shown at the branch points. *Luteimonas aquatica* RIB1-20^T was used as an outgroup. Bar: 0.01 substitutions per nucleotide position

No contamination was detected for the 16S rRNA gene. The genome had a total of 3787 genes, including 3728 protein-coding genes, 53 tRNA genes, 6 rRNA genes (one 5S, three 16S and two 23S rRNA genes). The ANI values between the strain SYSU H10001^T and the type strains of the species *L. panacisoli* JCM 19212^T and *L. soli* KCTC 22011^T were 84.6% and 83.8%, respectively, which were lower than the threshold value recommended for distinguishing novel

species (Goris et al. 2007; Richter and Rosselló-Móra 2009). The general features of the genomes of *L. prati* SYSU H10001^T and *L. panacisoli* JCM 19212^T and *L. soli* KCTC 22011^T are shown in Table 2.

The analysis of secondary metabolic gene clusters revealed that strain SYSU H10001^T harboured fewer gene clusters than other species in the genus. This gene clusters include 3 clusters of bacteriocins, 1 cluster of non-ribosomal peptide synthetase, 1 cluster of type 1

polyketide synthase and 1 cluster of arylpolyene (Supplementary Data 1). Based on the genome analysis and the phylogenomic tree, strains in the genus of *Lysobacter* could be divided into two groups, Group I that contains the species *L. capsici*, *L. antibioticus* ATCC 29479^T and *L. enzymogenes* ATCC 29487^T with higher secondary metabolic genes cluster, and Group II the other species. Strain SYSU H10001^T form a subclade with the Group II members in the phylogenomic tree (Fig. 2 and Supplementary Data 2). Similar to certain members in Group II with high gene clusters such as *Lysobacter* sp. ZS60, *Lysobacter daejeonensis* GH1-9^T and *Lysobacter maris* HZ9B^T, the genome of strain SYSU H10001^T yielded 29 secondary metabolic genes (0.78%, secondary metabolic gene numbers/predicted gene numbers).

Chemotaxonomical characteristics

The ubiquinone Q-8 was detected as the predominant respiratory ubiquinone in strain SYSU H10001^T. The polar lipid profile consists of diphosphatidylglycerol, phosphatidylglycerol, phosphatidylethanolamine and three unidentified phospholipids (Supplementary Fig. S2). The major fatty acids (> 10%) are iso-C_{15:0}, iso-C_{17:0} and Summed Feature 9 (10-methyl C_{16:0} and/or iso-C_{17:1} ω9c). While the main fatty acids types between strains SYSU H10001^T and the related type strains remain the same, the observed difference is at the proportion of these fatty acids, particularly the Summed Feature 9. The detailed fatty acid compositions are listed in Supplementary Table S1.

The phylogenetic analyses, morphological and chemotaxonomic characteristics support the characterisation of strain SYSU H10001^T as a member of the genus *Lysobacter*. The differences in biochemical

Fig. 2 RAxML phylogenomic tree using genomes of selected members of the genus *Lysobacter* to determine the phylogenetic position of strain SYSU H10001^T. The phylogenomic tree was generated with the RAxML method using the default parameters. *Luteimonas rhizosphaerae* 4-12 (NRQR000000000) was used as an outgroup. Bar, 0.1 substitutions per nucleotide position. *Note* The heatmap showed the percentage of the secondary metabolic genes of each genome, the value was calculated by secondary metabolic genes/prediction genes

characteristics and fatty acid compositions distinguish strain SYSU H10001^T from its closely related strains. Based on the above characteristics, strain SYSU H10001^T is considered to represent a novel species of the genus *Lysobacter*, for which the name *Lysobacter prati* is proposed.


Description of *Lysobacter prati* sp. nov

Lysobacter. prati (pra'ti. L. gen. neut. n. *prati* of a meadow, the place from where the novel strain is isolated).

Cells are Gram-staining negative, aerobic, non-motile, and rod-shaped. Colonies on YIM 38 agar are smooth, circular, opaque and yellow in colour after 3 days of cultivation at 28 °C. Growth occurs at 4–37 °C (optimum 28 °C), pH 6.0–9.0, and in the presence of up to 1.5% (w/v) NaCl. Positive for nitrate reduction, milk coagulation and peptonisation, and catalase and oxidase activities, but not for urease activity. Give positive results in tests for hydrolysis of gelatin, esculin, and tweens 40, 60 and 80, but not in tests for production of H₂S or hydrolysis of cellulose, starch or tween 20. Utilizes D-cellobiose, D-galactose, D-glucose, glycerol, D-maltose, and D-sucrose as sole carbon sources but not L-arabinose, D-fructose, inositol, D-ribose, lactose, D-mannitol, D-mannose, D-

Table 2 General features of the genomes of *L. prati* SYSU H10001^T, *L. panacisoli* JCM 19212^T and *L. soli* KCTC 22011^T

| Attribute | <i>L. prati</i> | <i>L. panacisoli</i> | <i>L. soli</i> |
|-----------------------------|-----------------|----------------------|----------------|
| Genomic size (bp) | 4,127,215 | 3,879,713 | 3,953,742 |
| Number of contigs | 48 | 3 | 27 |
| Mean length (bp) | 85,983 | 1,293,234 | 146,434 |
| N ₅₀ length (bp) | 420,681 | 2,610,236 | 285,382 |
| G + C content (%) | 66.5 | 67.5 | 67.7 |
| Protein-coding genes | 3728 | 3586 | 3544 |
| Genes assigned to COGs | 1685 | 2694 | 3267 |
| Genes assigned to KEGG | 1830 | 3284 | 1808 |



raffinose, D-ribose, L-rhamnose, D-sorbitol, D-sucrose, xylitol, or D-xylose. Utilizes L-alanine, L-aspartic acid, glycine, L-glutamic acid, L-glutamine, L-histidine, L-methionine, L-phenylalanine and L-serine, L-threonine, L-tyrosine and L-valine as sole nitrogen sources but not L-tyrosine, L-alanine, L-valine, L-tryptophan, L-glutamine, L-ornithine, adenine or hypoxanthine. In the API ZYM tests, positive for activities of acid phosphatase, alkaline phosphatase, cysteine arylamidase, esterase (C4), esterase lipase (C8), naphthol-AS-BI-phosphohydrolase, chymotrypsin, and valine arylamidase, α -glucosidase; and negative for activities of cysteine arylamidase, leucine arylamidase, lipase (C14), β -fucosidase, α -galactosidase, β -galactosidase, β -glucuronidase, *N*-acetyl- β -glucosaminidase, α -mannosidase, and trypsin. The predominant respiratory quinone is Q-8. The cellular polar lipids are diphosphatidylglycerol, phosphatidylglycerol, phosphatidylethanolamine and three unidentified phospholipids. The major fatty acids are iso-C_{15:0}, iso-C_{17:0} and Summed Feature 9 (10-methyl C_{16:0} and/or iso-C_{17:1} ω 9c).

The type strain SYSU H10001^T (= KCTC 72062^T = CGMCC 1.16662^T) was isolated using a soil sample collected from a meadow in Hongyuan county, Szechwan province, P. R. China. The DNA G + C content of the type strain is 66.5% (genome). The 16S rRNA gene and genome sequences of strain SYSU H10001^T were submitted to GenBank with accession numbers MN181427 and VLVN00000000, respectively.

Acknowledgements The authors are grateful to Prof. Jung-Sook Lee (KCTC, Korea) and Prof. Takuji Kudo (JCM, Japan) for kindly providing the experimental control strains. This research was supported by the National Key R&D Program of China (2017YFD0200503), Fundamental research funds for the central universities, SYSU (No. 19lgpy173) and National Fundamental Fund Project Subsidy Funds of Personnel Training of China (No. J1310025). W-J Li was also supported by Guangdong Province Higher Vocational Colleges and Schools Pearl River Scholar Funded Scheme (2014).

Author's contributions BZF and WJL designed research and project outline. BZF, ZXX, and LL performed isolation, deposition, and identification. YGX, JYJ and XTZ performed genome analysis. BZF, MW and WJL drafted the manuscript. All authors read and approved the final manuscript.

Compliance with ethical standards

Conflict of interest All the authors have declared no conflict of interest.

Ethical approval This article does not contain any studies with human participants or animals performed by any of the authors.

References

- Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ (1990) Basic local alignment search tool. *J Mol Biol* 215:403–410
- Athalye M, Goodfellow M, Lacey J, White RP (1985) Numerical classification of *Actinomadura* and *Nocardopsis*. *Int J Syst Bacteriol* 35:86–98
- Bligh EG, Dyer WJ (1959) A rapid method of total lipid extraction and purification. *Can J Biochem Physiol* 37:911–917
- Blin K, Shaw S, Steinke K, Villebro R, Ziemert N, Lee SY, Medema MH, Weber T (2019) antiSMASH 5.0: updates to the secondary metabolite genome mining pipeline. *Nucleic Acids Res.* <https://doi.org/10.1093/nar/gkz310>
- Castresana J (2000) Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Mol Biol Evol* 17:540–552
- Chen W, Zhao YL, Cheng J, Zhou XK, Salam N, Fang BZ, Li QQ, Hozzein WN, Li WJ (2016) *Lysobacter cavernae* sp. nov., a novel bacterium isolated from a cave sample. *Antonie Van Leeuwenhoek* 109:1047–1053
- Chhetri G, Kim J, Kim I, Seo T (2019) *Lysobacter caseinilyticus*, sp nov, a casein hydrolyzing bacterium isolated from sea water. *Antonie Van Leeuwenhoek.* <https://doi.org/10.1007/s10482-019-01267-7>
- Choi JH, Seok JH, Cha JH, Cha CJ (2014) *Lysobacter panacisoli* sp. nov., isolated from ginseng soil. *Int J Syst Evol Microbiol* 64:2193–2197
- Christensen P, Cook FD (1978) *Lysobacter*, a new genus of nonfruiting, gliding bacteria with a high base ratio. *Int J Syst Evol Microbiol* 28:367–393
- Collins MD, Jones D (1980) Lipids in the classification and identification of coryneform bacteria containing peptidoglycan based on 2,4-diaminobutyric acid. *J Appl Bacteriol* 48:459–470
- Collins MD, Pirouz T, Goodfellow M, Minnikin DE (1977) Distribution of menaquinones in actinomycetes and corynebacteria. *J Gen Microbiol* 100:221–230
- De Bruijn I, Cheng X, de Jager V, Expósito RG, Watrous J, Patel N, Postma J, Dorrestein PC, Kobayashi D, Raaijmakers JM (2015) Comparative genomics and metabolic profiling of the genus *Lysobacter*. *BMC Genom* 16:991
- Edgar RC (2004) MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792–1797
- Felsenstein J (1981) Evolutionary trees from DNA sequences: a maximum likelihood approach. *J Mol Evol* 17:368–376

- Felsenstein J (1985) Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 39:783–791
- Fitch WM (1971) Toward defining the course of evolution: minimum change for a specific tree topology. *Syst Biol* 20:406–416
- Fukuda W, Kimura T, Araki S, Miyoshi Y, Atomi H, Imanaka T (2013) *Lysobacter oligotrophicus* sp. nov., isolated from an Antarctic freshwater lake in Antarctica. *Int J Syst Evol Microbiol* 63:3313–3318
- Gordon RE, Barnett DA, Handerman JE, Pang CHN (1974) *Nocardia coeliaca*, *Nocardia autotrophica*, and the nocardin strain. *Int J Syst Bacteriol* 24:54–63
- Goris J, Konstantinidis KT, Klappenbach JA, Coenye T, Vandamme P, Tiedje JM (2007) DNA–DNA hybridization values and their relationship to whole-genome sequence similarities. *Int J Syst Evol Microbiol* 57:81–91
- Harrison PG, Strulo B (2000) SPADES—a process algebra for discrete event simulation. *J Logic Comput* 10:3–42
- Hernández I, Fernández C (2017) Draft genome sequence and assembly of a *Lysobacter enzymogenes* strain with biological control activity against root knot nematodes. *Genome Announc* 5:e00271–17
- Hyatt D, Chen GL, Locascio PF, Land ML, Larimer FW (2010) Prodigal: prokaryotic gene recognition and translation initiation site identification. *BMC Bioinformatics* 11:119
- Jiang Y, Li QY, Chen X, Jiang CL (2016). Isolation and cultivation methods of Actinobacteria. In: Dhanasekaran D (ed) *Actinobacteria-basics and biotechnological applications*, pp 39–57, (Rijeka: InTech)
- Kim I, Choi J, Chhetri G, Seo T (2019) *Lysobacter helvus* sp. nov. and *Lysobacter xanthus* sp. nov., isolated from soil in South Korea. *Antonie Van Leeuwenhoek* 112:1253–1262
- Kimura M (1980) A simple method for estimating evolutionary rates of base substitutions through comparative studies of nucleotide sequences. *J Mol Evol* 16:111–120
- Kimura M (1985) *The neutral theory of molecular evolution*. Cambridge University Press, Cambridge
- Kroppenstedt RM (1982) Separation of bacterial menaquinones by HPLC using reverse phase (RP18) and a silver loaded ion exchanger as stationary phases. *J Liq Chromatogr* 5:2359–2367
- Kumar S, Stecher G, Tamura K (2016) MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol* 33:1870–1874
- Letunic I, Bork P (2019) Interactive Tree Of Life (iTOL) v4: recent updates and new developments. *Nucleic Acids Res* 47:256–259
- Li WJ, Xu P, Schumann P, Zhang YQ, Pukall R, Xu LH, Stackebrandt E, Jiang CL (2007) *Georgenia ruanii* sp. nov., a novel actinobacterium isolated from forest soil in Yunnan (China), and emended description of the genus *Georgenia*. *Int J Syst Evol Microbiol* 57:1424–1428
- Lin SY, Hameed A, Wen CZ, Liu YC, Hsu YH, Lai WA, Young CC (2015) *Lysobacter lycopersici* sp. nov., isolated from tomato plant *Solanum lycopersicum*. *Antonie Van Leeuwenhoek* 107:1261–1270
- Luo Y, Dong HQ, Zhou M, Huang YL, Zhang H, He WL, Sheng HM, An LZ (2019) *Lysobacter psychrotolerans* sp. nov., isolated from soil in the Tianshan Mountains, Xinjiang, China. *Int J Syst Evol Microbiol* 69:926–931
- Minnikin DE, Collins MD, Goodfellow M (1979) Fatty acid and polar lipid composition in the classification of *Cellulomonas*, *Oerskovia* and related taxa. *J Appl Bacteriol* 47:87–95
- Minnikin D, O'Donnell A, Goodfellow M, Alderson G, Athalye M, Schaal A, Parlett JH (1984) An integrated procedure for the extraction of bacterial isoprenoid quinones and polar lipids. *J Appl Bacteriol* 2:233–241
- Nie GX, Ming H, Li S, Zhou EM, Cheng J, Tang X, Feng HG, Tang SK, Li WJ (2012) *Amycolatopsis dongchuanensis* sp. nov., a novel actinobacterium isolated from dry-hot valley in Yunnan, south-west China. *Int J Syst Evol Microbiol* 62:2650–2656
- Panthee S, Hamamoto H, Paudel A, Sekimizu K (2016) *Lysobacter*, species: a potential source of novel antibiotics. *Arch Microbiol* 198(9):839–845
- Park JH, Kim R, Aslam Z, Jeon CO, Chung YR (2008) *Lysobacter capsici* sp. nov. with antimicrobial activity, isolated from the rhizosphere of pepper, and emended description of the genus *Lysobacter*. *Int J Syst Evol Microbiol* 58:387–392
- Pridham TG, Gottlieb G (1948) The utilization of carbon compounds by some Actinomycetales as an aid for species determination. *J Bacteriol* 56:107–114
- Reasoner DJ, Geldreich EE (1985) A new medium for the enumeration and subculture of bacteria from potable water. *Appl Environ Microbiol* 49:1–7
- Richter M, Rosselló-Móra R (2009) Shifting the genomic gold standard for the prokaryotic species definition. *Proc Natl Acad Sci USA* 106:19126–19131
- Saitou N, Nei M (1987) The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 4:406–425
- Sasser M (1990) Identification of bacteria by gas chromatography of cellular fatty acids, MIDI technical note 101. Microbial ID Inc, Newark
- Shirling EB, Gottlieb D (1966) Methods for characterization of *Streptomyces* species. *Int J Syst Bacteriol* 16:313–340
- Siddiqi MZ, Im WT (2016) *Lysobacter hankyongensis* sp. nov., isolated from activated sludge and *Lysobacter sediminicola* sp. nov., isolated from freshwater sediment. *Int J Syst Evol Microbiol* 66:212–218
- Srinivasan S, Kim MK, Sathiyaraj G, Kim HB, Kim YJ, Yang DC (2010) *Lysobacter soli* sp. nov., isolated from soil of a ginseng field. *Int J Syst Evol Microbiol* 60:1543–1547
- Stamatakis A (2014) RAxML version 8: a tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* 30:1312–1313
- Tamaoka J, Katayama-Fujimura Y, Kuraishi H (1983) Analysis of bacterial menaquinone mixtures by high performance liquid chromatography. *J Appl Bacteriol* 54:31–36
- Thompson JD, Gibson TJ, Plewniak F, Jeanmougin F, Higgins DG (1997) The CLUSTAL_X windows interface: flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res* 25:4876–4882
- Wei DQ, Yu TT, Yao JC, Zhou EM, Song ZQ, Yin YR, Ming H, Tang SK, Li WJ (2012) *Lysobacter thermophilus* sp. nov., isolated from a geothermal soil sample in Tengchong, south-west China. *Antonie Van Leeuwenhoek* 102:643–651

- Weon HY, Kim BY, Kim MK, Yoo SH, Kwon SW, Go SJ, Stackebrandt E (2007) *Lysobacter niabensis* sp. nov. and *Lysobacter niastensis* sp. nov., isolated from greenhouse soils in Korea. *Int J Syst Evol Microbiol* 57:548–551
- Williams ST, Goodfellow M, Alderson G (1989) Genus *Streptomyces* Waksman and Henrici 1943, 339^{AL}. In: Williams ST, Sharpe ME, Holt JG (eds) *Bergey's manual of systematic bacteriology*, vol 4. Baltimore, Williams & Wilkins, pp 2453–2492
- Wu M, Scott AJ (2012) Phylogenomic analysis of bacterial and archaeal sequences with AMPHORA2. *Bioinformatics* 28:1033–1034
- Xiao M, Zhou XK, Chen X, Duan YQ, Alkhalifah MDH, Im WT, Hozzein WN, Chen W, Li WJ (2019) *Lysobacter tabacisoli* sp. nov., isolated from rhizosphere soil of *Nicotiana tabacum* L. *Int J Syst Evol Microbiol* 69:1875–1880
- Xie YX, Wright S, Shen YM, Du LC (2012) Bioactive natural products from *Lysobacter*. *Nat Prod Rep* 29:1277–1287
- Xu P, Li WJ, Tang SK, Zhang YQ, Chen GZ, Chen HH, Xu LH, Jiang CL (2005) *Naxibacter alkalitolerans* gen. nov., sp. nov., a novel member of the family ‘Oxalobacteraceae’ isolated from China. *Int J Syst Evol Microbiol* 55:1149–1153
- Yoon SH, Ha SM, Kwon S, Lim J, Kim Y, Seo H, Chun J (2017a) Introducing EzBioCloud: a taxonomically united database of 16S rRNA and whole genome assemblies. *Int J Syst Evol Microbiol* 67:1613–1617
- Yoon SH, Ha SM, Lim JM, Kwon SJ, Chun J (2017b) A large-scale evaluation of algorithms to calculate average nucleotide identity. *Antonie Van Leeuwenhoek* 110:1281–1286
- Zhang XJ, Yao Q, Wang YH, Yang SZ, Feng GD, Zhu HH (2019) *Lysobacter silvisoli* sp. nov., isolated from forest soil. *Int J Syst Evol Microbiol* 69:93–98

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.