

# Positivity Concepts on Holomorphic Line Bundles and Theories on Canonical Kähler Metrics

28 – 31 January, 2018, at Osaka City University

## Program

### January 28

**13:30 – 14:00 Registration**

**14:10 – 15:00 Martin Sera (Chalmers University of Technology and University of Gothenburg)**

Chern forms of singular metrics on vector bundles

**15:40 – 16:30 Genki Hosono (The University of Tokyo)**

T.B.A.

**16:40 – 17:30 Song-Yan Xie (Max Planck Institute for Mathematics)**

On the Kobayashi hyperbolicity of certain surfaces in  $\mathbb{CP}^3$

### January 29

**10:00 – 11:20 Mini-Lecture by Ryosuke Takahashi (Tohoku University)**

Lectures on extremal Kähler metrics

**11:30 – 12:20 Satoshi Nakamura (Tohoku University)**

Generalized Kähler Einstein metrics and uniform stability for toric Fano manifolds

**14:10 – 15:00 Thibaut Delcroix (ENS Paris)**

Mabuchi's K-energy functional on horosymmetric varieties

**15:10 – 16:00 Tomoyuki Hisamoto (Nagoya University)**

T.B.A.

**16:10 – Discussion**

### January 30

**10:00 – 11:20 Mini-Lecture by Ryosuke Takahashi (Tohoku University)**

Lectures on extremal Kähler metrics

**11:30 – 12:20 Kento Fujita (Kyoto University)**

Openness results for uniform K-stability

**14:10 – 15:00 Ryosuke Takahashi (Tohoku University)**

The inverse Monge-Ampère flow and optimal degenerations on toric Fano manifolds

**15:40 – 16:30 Ryosuke Nomura (The University of Tokyo)**

Miyaoka-Yau inequality for compact Kähler manifolds with semi-positive canonical bundle

**16:40 – 17:30 Shin Kikuta (Kogakuin University)**

Degeneration of positivity for log-canonical bundle and Kähler-Einstein metric

## **January 31**

**11:30 – 12:20 Masanori Adachi (Shizuoka University)**

T.B.A.

**14:10 – 15:00 Dinh Tuan Huynh (Osaka University)**

On the truncation level in Cartan's Second Main Theorem for entire holomorphic curves into projective plane

**15:10 – 16:00 Takato Uehara (Saga University)**

On a construction of non-projective K3 surfaces

**16:10 – Discussion**

# Abstracts

**Masanori Adachi (Shizuoka University)**

T.B.A.

**Thibaut Delcroix (ENS Paris)**

Mabuchi's K-energy functional on horosymmetric varieties

Spherical varieties form a very large class of almost-homogeneous varieties and as such a promising class on which one can study existence of canonical metrics. However it is very difficult to do Kähler geometry in a uniform way on all spherical varieties. I will introduce a subclass containing homogeneous toric bundles and group compactifications on which I am able to adapt toric geometry techniques to obtain a combinatorial sufficient criterion of properness of Mabuchi's K-energy functional. I will also present applications to existence of cscK and log-Kähler-Einstein metrics.

**Kento Fujita (Kyoto University)**

Openness results for uniform K-stability

It is expected that several “openness” results holds for uniform K-stability of polarized varieties. In this talk, I will discuss certain partial results for this problem.

**Tomoyuki Hisamoto (Nagoya University)**

T.B.A.

**Genki Hosono (The University of Tokyo)**

T.B.A.

**Dinh Tuan Huynh (Osaka University)**

On the truncation level in Cartan's Second Main Theorem for entire holomorphic curves into projective plane

Let  $f : \mathbb{C} \rightarrow \mathbb{P}^2(\mathbb{C})$  be an entire holomorphic curve and let  $\{L_i\}_{1 \leq i \leq q}$  be a family of  $q \geq 4$  lines in general position in projective plane. If  $f$  is linearly nondegenerate, i.e. its image is not contained in any line, then the classical Second Main Theorem of Cartan states that the following inequality holds true outside a subset of  $(0, \infty)$  of finite Lebesgue measure:

$$(q-3)T_f(r) \leq \sum_{i=1}^q N_f^{[2]}(r, L_i) + o(T_f(r)).$$

Here  $T_f(r)$  and  $N_f^{[2]}(r, L_i)$  stand for the order function and the 2-truncated counting functions in Nevanlinna theory. It is conjectured that in the above estimate, the truncation level of the counting functions can be decreased to 1, provided that that  $f$  is algebraically nondegenerate (i.e. its image is not contained in any algebraic curve). In this talk, we will provide a partial answer to this conjecture and propose a strategy to achieve the full proof.

[1] Henri Cartan, Sur les zéros des combinaisons linéaires de  $p$  fonctions holomorphes données, *Mathematica*, **7**, 5–31, 1933.

[2] Michael McQuillan, Diophantine approximations and foliations, *Inst. Hautes Études Sci. Publ. Math.*, **87** 121–174, 1998.

- [3] Marco Brunella, Courbes entières et feuilletages holomorphes, *Ens. Math.*, **45**, 195–216, 1999.
- [4] Julien Duval and Dinh Tuan Huynh, A geometric second main theorem, *Math. Ann.*(2017), DOI:10.1007/s00208-017-1538-y.

**Shin Kikuta (Kogakuin University)**

Degeneration of positivity for log-canonical bundle and Kähler–Einstein metric

We would like to discuss some relations between the positivity of the log-canonical bundle and the boundary behavior of the Kähler-Einstein metric over a quasi-projective manifold. In particular, via its volume growth, we characterize when the boundary is of general type. We will also talk about our recent progress in the case when the boundary is Calabi–Yau.

**Satoshi Nakamura (Tohoku University)**

Generalized Kähler Einstein metrics and uniform stability for toric Fano manifolds

Generalized Kähler Einstein metrics, introduced by Mabuchi in 2000, is a generalization of Kähler Einstein metrics for Fano manifolds with non vanishing Futaki invariant. (Some people called them Mabuchi metrics or Mabuchi Einstein metrics or Mabuchi soliton... We should fix the name.) In this talk, We give a complete criterion for the existence of generalized Kähler Einstein metrics on toric Fano manifolds from view points of a uniform stability in a sense of GIT and the properness of a functional on the space of Kähler metrics. This talk is based on arXiv:1706.01608 (to appear in Tohoku Math J.).

**Ryosuke Nomura (The University of Tokyo)**

Miyaoka-Yau inequality for compact Kähler manifolds with semi-positive canonical bundle

Miyaoka-Yau inequality is an inequality for Chern classes which is known to hold for minimal projective varieties. In this talk, we prove the inequality for compact Kähler manifolds with semi-positive canonical bundle.

**Martin Sera (Chalmers University of Technology and University of Gothenburg)**

Chern forms of singular metrics on vector bundles

After recalling the notion of Segre and Chern forms for smooth metrics, we would like to present some results of a joint work with Richard Lärkäng, Hossein Raufi and Jean Ruppenthal, where we study singular Hermitian metrics on holomorphic vector bundles. Although it is in general not possible to define the curvature as a current with measure coefficients, we give a definition of Chern forms as closed currents of order 0. For Griffiths positive/negative singular metrics, we can define these Chern currents up to the degree less or equal than the codimension of the singular locus using pluripotential theoretical methods. Finally, we will discuss some properties of these Chern currents, for instance that they represent the Chern classes of the vector bundle.

**Ryosuke Takahashi (Tohoku University)**

The inverse Monge-Ampère flow and optimal degenerations on toric Fano manifolds

We discuss the gradient flow of the Ding energy functional, referred as the inverse Monge-Ampère flow on toric Fano manifolds. We prove that the inverse Monge-Ampère flow produces a “most destabilizing degeneration” with respect to the Ding-stability. This research is based on a joint work with T.C. Collins (Harvard Univ.) and T. Hisamoto (Nagoya Univ.).

**Takato Uehara (Saga University)**

On a construction of non-projective K3 surfaces

In this talk, we construct K3 surfaces by gluing two 9-point blowups of the complex projective plane, where the existence of overlaps is guaranteed by Arnol’d’s theorem. A calculation of their period maps

shows that such K3 surfaces constitute a large family, including non-projective K3 surfaces. Furthermore, we comment on a relation with automorphisms of K3 surfaces having positive entropy. This is a joint work in progress with T. Koike.

**Song-Yan Xie (Max Planck Institute for Mathematics)**

On the Kobayashi hyperbolicity of certain surfaces in  $\mathbb{CP}^3$

The Green-Griffiths-Lang Conjecture in  $\mathbb{CP}^3$  stipulates that, for a general surface of degree  $\geq 5$ , there should be no holomorphic entire curve on it. In this talk, we discuss some recent progress towards this conjecture.