


# What is Low-Cost Side Scan Sonar Habitat Mapping?

Adam J. Kaeser and Thomas L. Litts  
U.S. Fish and Wildlife Service  
Georgia Department of Natural Resources

Inspecting substrate with a drop camera




## A Low-Cost Remote Sensing Technique is Needed for Waterscape Research in Navigable Systems

- Producing detailed, landscape level habitat maps of turbid, non-wadeable systems is costly or impossible using traditional methods
- Side scan sonar is an active remote sensing device that uses reflected sound to produce images of underwater features
- The inexpensive Humminbird® Side Imaging system (\$2000-2700) generates high resolution imagery and employs a small, adjustable transducer
- Over the last 5 years we have pursued the development of a complete method for mapping with the SI system, incorporating several complementary objectives (listed below)
- Our work has focused on streams and rivers of Georgia and Florida

## Objectives of this Initiative

- Develop approaches for field sonar surveys
- Develop techniques for georeferencing and transformation (ie. geoprocessing) of sonar imagery for use in a GIS
- Produce detailed maps of instream habitat features (eg. substrates, LWD, depth) via image interpretation and manual digitization
- Evaluate/validate the techniques and map accuracies through field studies
- Develop and offer the tools, products, and training to interested professionals
- Continue testing and developing new applications of low-cost sonar habitat mapping

## The Process of Sonar Habitat Mapping



## What Do I Need to Get Started?



### Software & Training

ArcGIS ArcView level 9.2+, or 10.0- image geoprocessing and map production


Irfanview & ET Geowizards- freeware used during geoprocessing

Sonar processing tools- VBA scripts created by Thom Litts and delivered in a GIS project (provided freely at your request)


A copy of our "Sonar Imagery Geoprocessing Workbook"

Method training offered in workshop format, soon delivered as a free, web available "Guide to Low-Cost Sonar Habitat Mapping"

Next workshop offered- Southern Division AFS in Biloxi MS, Jan 26-29, 2012




## A Variety of Habitat Features Are Revealed



## Validation/Accuracy Assessment Studies

- Kaeser, A. J., and T. L. Litts. 2008. An assessment of deadhead logs and large woody debris using side scan sonar and field surveys in streams of Southwest Georgia. *Fisheries* 33:589-597.
- Kaeser, A. J., and T. L. Litts. 2010. A novel technique for mapping habitat in navigable streams using low-cost side scan sonar. *Fisheries* 35:163-174.
- Kaeser, A. J., T. L. Litts, and T. W. Tracy. *In review*. Using low-cost side scan sonar to map habitat throughout the lower Flint River, Georgia, USA. *River Research and Applications*.



## Applications are Widespread & Diverse

- Organism-habitat research in systems and at scales not previously feasible
- Studies of individual habitat use and behavioral patterns (eg. via radiotelemetry), identification/quantification/prediction of critical habitat (eg. sturgeon spawning habitat)
- Landuse associations with in-stream habitat (eg. patterns of LWD distribution with respect to riparian landuse)
- Monitoring habitat change over time (eg. sediment redistribution)
- Similar applications in lakes and reservoirs possible (eg. littoral zone mapping)
- Habitat map layers can be viewed in Google Earth and on smart phones

## The Future of this Initiative

- To demonstrate the utility of low-cost sonar mapping we are applying the method in ongoing studies of turtle, fish, and freshwater mussel habitat use and availability, and time-lapse detection of changes in substrate deposition
- To receive email announcements of new training products, or make inquiries contact Adam at adam\_kaeser@fws.gov

**\*Affordability, speed, flexibility, ease of training, and access to software are key traits of low-cost sonar habitat mapping. The future is now for waterscape research.**

## Acknowledgements

• Funding provided by:



• Support provided by:

The Joseph W. Jones Ecological Research Center at Ichauway

Center for Remote Sensing and Mapping Science, Department of Geography, University of Georgia

Student interns Wes Tracy, Philip Marley, and Joshua Hubbell

