

W2W: A Python package that injects WUDAPT's Local Climate Zone information in WRF

Matthias Demuzere^{*1}, Daniel Argüeso², Andrea Zonato³, and Jonas Kittner¹

¹ Urban Climatology Group, Department of Geography, Ruhr-University Bochum, Bochum, Germany ² Physics Department, University of the Balearic Islands, Palma, Spain ³ Atmospheric Physics Group, Department of Civil, Environmental and Mechanical Engineering, University of Trento, Trento, Italy

DOI: [10.21105/joss.0XXXX](https://doi.org/10.21105/joss.0XXXX)

Software

- [Review](#) ↗
- [Repository](#) ↗
- [Archive](#) ↗

Editor: [Editor Name](#) ↗

Submitted: 01 January XXXX

Published: 01 January XXXX

License

Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License ([CC BY 4.0](#)).

Summary

An important objective of WUDAPT, the World Urban Database and Access Portals Tools community project, is 1) to acquire and make accessible coherent and consistent information on form and function of urban morphology relevant to climate weather, and environment studies, and 2) to provide tools that extract relevant urban parameters and properties for models and model applications at appropriate scales for various climate, weather, environment, and urban planning purposes (Ching et al., 2018).

The Python-based WUDAPT-to-WRF (W2W) package is developed in this context, and translates Local Climate Zone (LCZ) maps into urban canopy parameters readable by WRF, the community “Weather Research and Forecasting” model (Skamarock et al., 2021). It is the successor of the Fortran-based W2W package developed by Brousse et al. (2016) and Martilli et al. (2016), and provides an improved, more simple, and more efficient procedure to use LCZ information in WRF. Some important changes include a direct manipulation of the geogrid files (without the creation of temporary files), and the use of average LCZ-based urban morphological parameters instead of assigning them to the modal LCZ class.

Statement of need

Since the pioneering work of Brousse et al. (2016) and Martilli et al. (2016), the level-0 WUDAPT information, the Local Climate Zone maps, have been used increasingly in WRF.

We expect this trend to continue, because of two recent developments: 1) the creation of city-wide LCZ maps is now easier than ever with the launch of the LCZ Generator web application (Demuzere et al., 2021), and 2) WRF versions > 4.3 (Skamarock et al., 2021) are able to ingest 10 or 11 built classes (corresponding to WUDAPT's LCZs) by default, whereas previous WRF versions required manual code changes (see Martilli et al. (2016), Zonato & Chen (2021) and Zonato et al. (2021) for more information).

Because of these developments, an improved, Python-based, WUDAPT-to-WRF (W2W) routine is presented here, so as to make the translation of LCZ-based parameters better and more simple.

*corresponding author

Initial data requirements

In order to use the tool, two input files are required:

1. A **geo_em.d0X** (.nc) file for the inner WRF model domain in which one would like to use the LCZ-based information. This file can be produced by WRF's `geogrid.exe` tool as part of the WRF Preprocessing System (WPS), without additional modifications of the standard procedure.
2. A **Local Climate Zone map** (.tif) file that is slightly bigger than the domain extent of the `geo_em.d0X.nc` file. There are a number of ways to obtain an LCZ map for your region of interest (ROI):
 - Extract your ROI from the continental-scale LCZ maps for Europe (Demuzere et al., 2019) or the United States (Demuzere et al., 2020) (see [here](#) for more info).
 - Check if your ROI is already covered by the many LCZ maps available in the [submission table](#) of the LCZ Generator.
 - Use the [LCZ Generator](#) to make a LCZ map for your ROI. See also [here](#) for more information. When using LCZ maps produced with the LCZ Generator, by default the Gaussian filtered LCZ map is used ($-l = 1$).

Workflow

The goal of the Python-based `W2W` tool is to obtain an inner WRF domain file (`geo_em.d0X.nc`) that contains the built LCZ classes and their corresponding urban canopy parameters relevant for all urban parameterizations embedded in WRF: the single layer urban canopy model (Noah/SLUCM, Kusaka et al. (2001)), the Building Environment Parameterization (BEP, Martilli et al. (2002)), and BEP+BEM (Building Energy Model, Salamanca et al. (2010)).

To get to that point, a number of sequential steps are required:

- *Step 1: Remove the default urban land cover*

The default urban land cover from MODIS is replaced with the dominant surrounding vegetation category, as done in Li et al. (2020). This procedure affects WRF's parameters `LU_INDEX`, `LANDUSEF` and `GREENFRAC`. `LU_INDEX` is selected as the dominant category from the $-n$ (default = 45) nearest grid points (excluding ocean, urban and lakes). `GREENFRAC` is calculated as the mean over all grid points with that dominant vegetation category among the $-n$ nearest points. For each grid point, if `LANDUSEF` had any percentage of urban, it is set to zero and the percentage is added to the dominant vegetation category assigned to that grid point.

Resulting output: **geo_em.d0X_NoUrban.nc**

- *Step 2: Define the LCZ-based urban extent*

LCZ-based impervious fraction values (`FRC_URB2D`, available from `LCZ_UCP_default.csv`) are assigned to the original 100 m resolution LCZ map, and are aggregated to the WRF resolution. Areas with `FRC_URB2D` < 0.2 ($-f$) are currently considered non-urban. This choice has been made to avoid the use of the urban schemes in areas where the majority of the landuse is vegetated, since the impact of the impervious surfaces is low. The `FRC_URB2D` field is also used to mask all other urban parameter fields, so that their extent is consistent.

Resulting output: **geo_em.d0X_LCZ_extent.nc**

77 ▪ *Step 3: Introduce modal built LCZ classes*

78 For each WRF grid cell, the mode of the underlying built LCZ classes is added to LU_INDEX
 79 (numbered from 31-41). See [here](#) for more info. Note that the W2W routine by default
 80 considers LCZ classes 1-10 as built classes (*-b*). In some cases, also LCZ E (or 15 - Bare rock
 81 or paved) can be considered as a built LCZ class, as it might reflect large asphalt surfaces
 82 such as big parking lots or airstrips. In that case, the user must make sure the *-b* argument
 83 is set appropriately.

84 ▪ *Step 4: Assign urban canopy parameters*

85 Two procedures are followed when assigning the various urban canopy parameters to the LCZ
 86 map and translating this information onto WRF's grid:

87 **Procedure 1: Morphological parameters** are assigned directly to the high-resolution LCZ
 88 map, and are afterwards aggregated to the lower-resolution WRF grid. As a result, the method
 89 produces a unique urban morphology parameter value for each WRF grid cell. This was found
 90 to be more efficient in reproducing urban boundary layer features, especially in the outskirts of
 91 the city (Zonato et al., 2020), and is in line with the [WUDAPT-to-COSMO](#) routine (Varentsov
 92 et al., 2020).

93 Morphological urban canopy parameter values are provided in `LCZ_UCP_default.csv`, and
 94 are generally based on values provided in Stewart & Oke (2012) and Stewart et al. (2014).
 95 Note however that the values of `MH_URB2D_MIN`, `MH_URB2D`, `MH_URB2D_MAX` for
 96 LCZ 7 are set to 4, 5 and 6 m instead of 2, 3 and 4 m, because the minimum building height
 97 that can be assigned to BEP-BEM is 5m if $dz_u = 5m$ (standard value) is used.

98 In addition:

- 99 ▪ While `URBPARAM_LCZ.TBL` (stored in WRF's `run/` folder) has values on street width
 100 (`SW`), W2W derives street width from the mean building height (`MH_URB2D`) and the
 101 Height-to-Width ratio (`H2W`), to have these fields consistent.
- 102 ▪ Building width (`BW`), is derived from $(BLDFR_URB2D / (FRC_URB2D - BLDFR_URB2D))$
 103 * `SW`, these values being available from the look-up table `LCZ_UCP_default.csv`.
- 104 ▪ Plan (`LP_URB2D`), frontal (`LF_URB2D`) and total (`LB_URB2D`) area indices are based
 105 on formulas in Zonato et al. (2020).
- 106 ▪ `HI_URB2D` is obtained by fitting a bounded normal distribution to the minimum
 107 (`MH_URB2D_MIN`), mean (`MH_URB2D`), and maximum (`MH_URB2D_MAX`)
 108 building height, as provided in `LCZ_UCP_default.csv`. The building height
 109 standard deviation is also required, and is approximated as $(MH_URB2D_MAX -$
 110 $MH_URB2D_MIN) / 4$.
- 111 ▪ For computational efficiency, `HI_URB2D` values lower than 5% were set to 0 after
 112 resampling, the remaining `HI_URB2D` percentages are re-scaled to 100%.

113 **Procedure 2:** In line with the former Fortran-based W2W procedure, **radiative and thermal**
 114 **parameters** are assigned to the modal LCZ class that is assigned to each WRF grid cell
 115 (see *Step 3*). These parameter values are not stored in the netcdf output, but are read from
 116 `URBPARAM_LCZ.TBL` and assigned automatically to the modal LCZ class when running the
 117 model.

118 ▪ *Step 5: Adjust global attributes*

119 In a final step, some global attributes are adjusted in the resulting netcdf files:

- NBUI_MAX is added as a global attribute, reflecting the maximum amount of HI_URB2D classes that are not 0 across the model domain. This parameter can be used when compiling WRF, to optimize memory storage.
- NUM_LAND_CAT is set to 41, to reflect the addition of 10 (or 11) built LCZ classes. This is not only done for the highest resolution domain file (e.g. d04), but also for **all of its lower-resolution parent domain files (e.g. d01, d02, d03)**. As such, make sure these files are also available in the input data directory. In case the parent domain files have NUM_CAT_LAND \neq 41, new parent domain files will be written to your drive with the extension _41.

Resulting output: `geo_em.d0X_LCZ_params.nc` (and `geo_em.d0X_41.nc`)

Integration in WRF's preprocessing

The current tool is designed to work with the `geo_em.d0X` files produced by `geogrid.exe`, which is available in the WRF Preprocessing System (WPS). WPS needs to be at a version >3.8 , in order to incorporate the urban geometrical parameters in the `URB_PARAM` matrix (Glotfelty et al., 2013). The user should run `geogrid.exe` using its default settings, which will provide the various `geo_em.d0X.nc` files containing the static data fields. No additional variables are required, neither in the `namelist.wps` nor within the `GEOGRID.TBL` table. The `W2W` tool (Figure 1) reads the standard `geo_em.d0X.nc` files (for all the domains) and produces the aforementioned `geo_em.d0X_LCZ_params.nc` files. The user should then simply rename these files to the standard name for each of the domains (e.g. rename `geo_em.d01_41.nc` to `geo_em.d01.nc`, `geo_em.d04_LCZ_params.nc` to `geo_em.d04.nc`, ...), which will serve as input to the `metgrid.exe` module (Figure 1).




Figure 1: Modified workflow to set-up and run a WRF simulations including urban parameters derived from LCZs using W2W.

Potential use cases

The files provided as output by W2W allow a wide range of applications, including - but not limited to - addressing the impact of:

- urbanization, by running WRF with the default `geo_em.d0X.nc` and the `geo_em.d0X_NoUrban.nc` files (see for example Li et al. (2020) and Hirsch et al. (2021)).
- an improved urban land cover extent description, by running WRF with the default `geo_em.d0X.nc` and the `geo_em.d0X_LCZ_extent.nc` files (similar to for example Bhati & Mohan (2018) and Mallard et al. (2018)).
- a more detailed (LCZ-based) urban description, by running WRF with the default `geo_em.d0X.nc` and the `geo_em.d0X_LCZ_params.nc` files (see for example Brousse et al. (2016), Hammerberg et al. (2018), Molnár et al. (2019), Wong et al. (2019), Patel et al. (2020), Zonato et al. (2020), Ribeiro et al. (2021), Hirsch et al. (2021) and Patel et al. (2022)).

Important notes

- The LCZ-based urban canopy parameter values provided in `LCZ_UCP_default.csv` and `URBPARAM_LCZ.TBL` are universal and generic, and might not be appropriate for your ROI. If available, please adjust the urban canopy parameters values according to the characteristics of your ROI. A custom csv file can be specified using the `--lcz-ucp path/to/custom_file.csv` flag.
- It is advised to use this tool with urban parameterization options BEP or BEP+BEM (`sf_urban_physics = 2` or `3`, respectively). In case you use this tool with the SLUCM model (`sf_urban_physics = 1`), make sure your lowest model level is above the highest building height. If not, `real.exe` will provide the following error message: `ZDC + ZOC + 2m is larger than the 1st WRF level - Stop in subroutine urban - change ZDC and ZOC`.
- It is advised to use WRF versions > 4.3 , that are able to ingest 10 or 11 built classes (corresponding to WUDAPT's LCZs) by default (Skamarock et al., 2021), and WPS versions > 3.8 , in order to incorporate the urban geometrical parameters in the `URBPARAM` matrix (Glotfelty et al., 2013).

Acknowledgements

We acknowledge contributions and support from Alberto Martilli, Alejandro Rodriguez Sanchez and Oscar Brousse.

References

- Bhati, S., & Mohan, M. (2018). WRF-urban canopy model evaluation for the assessment of heat island and thermal comfort over an urban airshed in India under varying land use/land cover conditions. *Geoscience Letters*, 5(1). <https://doi.org/10.1186/s40562-018-0126-7>
- Brousse, O., Martilli, A., Foley, M., Mills, G., & Bechtel, B. (2016). WUDAPT, an efficient land use producing data tool for mesoscale models? Integration of urban LCZ in WRF over madrid. *Urban Climate*, 17, 116–134. <https://doi.org/10.1016/j.uclim.2016.04.001>
- Ching, J., Mills, G., Bechtel, B., See, L., Feddema, J., Wang, X., Ren, C., Brorousse, O., Martilli, A., Neophytou, M., Mouzourides, P., Stewart, I., Hanna, A., Ng, E., Foley, M., Alexander, P., Aliaga, D., Niyogi, D., Shreevastava, A., ... Theeuwesits, N. (2018). WUDAPT: An urban weather, climate, and environmental modeling infrastructure for the anthropocene. *Bulletin of the American Meteorological Society*, 99(9), 1907–1924. <https://doi.org/10.1175/BAMS-D-16-0236.1>

- Demuzere, M., Bechtel, B., Middel, A., & Mills, G. (2019). Mapping Europe into local climate zones. *PLOS ONE*, 14(4), e0214474. <https://doi.org/10.1371/journal.pone.0214474>
- Demuzere, M., Hankey, S., Mills, G., Zhang, W., Lu, T., & Bechtel, B. (2020). Combining expert and crowd-sourced training data to map urban form and functions for the continental US. *Scientific Data*, 7(1), 264. <https://doi.org/10.1038/s41597-020-00605-z>
- Demuzere, M., Kittner, J., & Bechtel, B. (2021). LCZ Generator: A Web Application to Create Local Climate Zone Maps. *Frontiers in Environmental Science*, 9(April). <https://doi.org/10.3389/fenvs.2021.637455>
- Glotfelty, T., Tewari, M., Sampson, K., Duda, M., Chen, F., & Ching, J. (2013). *NUDAPT 44: How to use NUDAPT dataset in WRF/SLUCM/MLUCM models* (p. 9). National Center for Atmospheric Research. <https://www.yumpu.com/en/document/read/26871494/how-to-use-nudapt-dataset-in-wrf-slucm-mlucm-models>
- Hammerberg, K., Brousse, O., Martilli, A., & Mahdavi, A. (2018). Implications of employing detailed urban canopy parameters for mesoscale climate modelling: a comparison between WUDAPT and GIS databases over Vienna, Austria. *International Journal of Climatology*, 38, e1241–e1257. <https://doi.org/10.1002/joc.5447>
- Hirsch, A. L., Evans, J. P., Thomas, C., Conroy, B., Hart, M. A., Lipson, M., & Ertler, W. (2021). Resolving the influence of local flows on urban heat amplification during heatwaves. *Environmental Research Letters*, 16(6), 064066. <https://doi.org/10.1088/1748-9326/ac0377>
- Kusaka, H., Kondo, H., Kikegawa, Y., & Kimura, F. (2001). A Simple Single-Layer Urban Canopy Model For Atmospheric Models: Comparison With Multi-Layer And Slab Models. *Boundary-Layer Meteorology*, 101(3), 329–358. <https://doi.org/10.1023/A:1019207923078>
- Li, Y., Fowler, H. J., Argüeso, D., Blenkinsop, S., Evans, J. P., Lenderink, G., Yan, X., Guerreiro, S. B., Lewis, E., & Li, X. F. (2020). Strong Intensification of Hourly Rainfall Extremes by Urbanization. *Geophysical Research Letters*, 47(14), 1–8. <https://doi.org/10.1029/2020GL088758>
- Mallard, M. S., Spero, T. L., & Taylor, S. M. (2018). Examining WRF's sensitivity to contemporary land-use datasets across the contiguous united states using dynamical downscaling. *Journal of Applied Meteorology and Climatology*, 57(11), 2561–2583. <https://doi.org/10.1175/JAMC-D-17-0328.1>
- Martilli, A., Brousse, O., & Ching, J. (2016). *Urbanized WRF modeling using WUDAPT*. <http://www.wudapt.org/wudapt-to-wrf/>. [https://www.wudapt.org/wp-content/uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.pdf%20\(Accessed%20on%2011%20August%202021](https://www.wudapt.org/wp-content/uploads/2016/05/Urbanized-WRF-modeling-using-WUDAPT-web-version-March2016.pdf%20(Accessed%20on%2011%20August%202021)
- Martilli, A., Clappier, A., & Rotach, M. W. (2002). An urban surface exchange parameterisation for mesoscale models. *Boundary-Layer Meteorology*, 104(2), 261–304. <https://doi.org/10.1023/A:1016099921195>
- Molnár, G., Gyöngyösi, A. Z., & Gál, T. (2019). Integration of an LCZ-based classification into WRF to assess the intra-urban temperature pattern under a heatwave period in Szeged, Hungary. *Theoretical and Applied Climatology*, 138(1-2), 1139–1158. <https://doi.org/10.1007/s00704-019-02881-1>
- Patel, P., Jamshidi, S., Nadimpalli, R., Aliaga, D. G., Mills, G., Chen, F., Demuzere, M., & Niyogi, D. (2022). Modelling Large-Scale Heatwave by Incorporating Enhanced Urban Representation. *Journal of Geophysical Research : Atmospheres*, 127, 1–33. <https://doi.org/10.1029/2021JD035316>

- Patel, P., Karmakar, S., Ghosh, S., & Niyogi, D. (2020). Improved simulation of very heavy rainfall events by incorporating WUDAPT urban land use/land cover in WRF. *Urban Climate*, 32(July 2019), 100616. <https://doi.org/10.1016/j.uclim.2020.100616>
- Ribeiro, I., Martilli, A., Falls, M., Zonato, A., & Villalba, G. (2021). Highly resolved WRF-BEP/BEM simulations over Barcelona urban area with LCZ. *Atmospheric Research*, 248(August 2020), 105220. <https://doi.org/10.1016/j.atmosres.2020.105220>
- Salamanca, F., Krpo, A., Martilli, A., & Clappier, A. (2010). A new building energy model coupled with an urban canopy parameterization for urban climate simulations—part I. formulation, verification, and sensitivity analysis of the model. *Theoretical and Applied Climatology*, 99(3-4), 331–344. <https://doi.org/10.1007/s00704-009-0142-9>
- Skamarock, W. C., Klemp, J. B., Dudhia, J. B., Gill, D. O., Liu, Z., Berner, J., Wang, W., Powers, J. G., Duda, M. G., Barker, D. M., & Huang, X.-Y. (2021). *A Description of the Advanced Research WRF Model Version 4.3* (July). National Center for Atmospheric Research. <https://doi.org/10.5065/1dfh-6p97>
- Stewart, I. D., & Oke, T. R. (2012). Local Climate Zones for Urban Temperature Studies. *Bulletin of the American Meteorological Society*, 93(12), 1879–1900. <https://doi.org/10.1175/BAMS-D-11-00019.1>
- Stewart, I. D., Oke, T. R., & Krayenhoff, E. S. (2014). Evaluation of the 'local climate zone' scheme using temperature observations and model simulations [Journal Article]. *International Journal of Climatology*, 34(4), 1062–1080. <https://doi.org/10.1002/joc.3746>
- Varentsov, M., Samsonov, T., & Demuzere, M. (2020). Impact of Urban Canopy Parameters on a Megacity's Modelled Thermal Environment. *Atmosphere*, 11(12), 1349. <https://doi.org/10.3390/atmos11121349>
- Wong, M. M. F., Fung, J. C. H., Ching, J., Yeung, P. P. S., Tse, J. W. P., Ren, C., Wang, R., & Cai, M. (2019). Evaluation of uWRF performance and modeling guidance based on WUDAPT and NUDAPT UCP datasets for Hong Kong. *Urban Climate*, 28(June 2018), 100460. <https://doi.org/10.1016/j.uclim.2019.100460>
- Zonato, A., & Chen, F. (2021). *Updates of WRF-urban in WRF 4.3: Local Climate Zones, Mitigation Strategies, building materials permeability and new buildings drag coefficient*. <http://www.wudapt.org/wudapt-to-wrf/>. https://ral.ucar.edu/sites/default/files/public/product-tool/urban-canopy-model/WRF_urban_update_Readme_file_WRF4.3.pdf
- Zonato, A., Martilli, A., Di Sabatino, S., Zardi, D., & Giovannini, L. (2020). Evaluating the performance of a novel WUDAPT averaging technique to define urban morphology with mesoscale models. *Urban Climate*, 31(May 2019), 100584. <https://doi.org/10.1016/j.uclim.2020.100584>
- Zonato, A., Martilli, A., Gutierrez, E., Chen, F., He, C., Barlage, M., Zardi, D., & Giovannini, L. (2021). Exploring the effects of rooftop mitigation strategies on urban temperatures and energy consumption. *Journal of Geophysical Research: Atmospheres*, 1–30. <https://doi.org/10.1029/2021JD035002>