

Hierarchical Continuous Time Dynamic Models with ctsem

July, 2018


Charles Driver

Max Planck Institute for Human Development






- Common approaches to dynamic modelling (latent change, autoregressive) assume that the relation between observations at different occasions are equal.






- Common approaches to dynamic modelling (latent change, autoregressive) assume that the relation between observations at different occasions are equal.
- This is fine in some cases, however problems can arise when time intervals between measurements are not equal.





- Common approaches to dynamic modelling (latent change, autoregressive) assume that the relation between observations at different occasions are equal.
- This is fine in some cases, however problems can arise when time intervals between measurements are not equal.
- Continuous time models determine the relations between each measurement occasion via a deterministic function of the continuous time parameters and the time interval between measurements.

$$\eta_t = A^* \eta_{t-1} + b^* + \zeta^*(t)$$







- More accuracy by incorporating time explicitly.



- More accuracy by incorporating time explicitly.
- More theoretically justifiable - things happen between measurement occasions.



- More accuracy by incorporating time explicitly.
- More theoretically justifiable - things happen between measurement occasions.
- More accurate than embedding approaches to differential equations, and no need to set embedding dimension.



- More accuracy by incorporating time explicitly.
- More theoretically justifiable - things happen between measurement occasions.
- More accurate than embedding approaches to differential equations, and no need to set embedding dimension.
- Tangible benefits:



- More accuracy by incorporating time explicitly.
- More theoretically justifiable - things happen between measurement occasions.
- More accurate than embedding approaches to differential equations, and no need to set embedding dimension.
- Tangible benefits:
 - More flexible data collection - no longer a need for consistent measurement intervals within or between individuals.



- More accuracy by incorporating time explicitly.
- More theoretically justifiable - things happen between measurement occasions.
- More accurate than embedding approaches to differential equations, and no need to set embedding dimension.
- Tangible benefits:
 - More flexible data collection - no longer a need for consistent measurement intervals within or between individuals.
 - Consistent parameter comparisons between different data collection schedules - between waves, studies, or both.



- More accuracy by incorporating time explicitly.
- More theoretically justifiable - things happen between measurement occasions.
- More accurate than embedding approaches to differential equations, and no need to set embedding dimension.
- Tangible benefits:
 - More flexible data collection - no longer a need for consistent measurement intervals within or between individuals.
 - Consistent parameter comparisons between different data collection schedules - between waves, studies, or both.
 - Differential equations:



- More accuracy by incorporating time explicitly.
- More theoretically justifiable - things happen between measurement occasions.
- More accurate than embedding approaches to differential equations, and no need to set embedding dimension.
- Tangible benefits:
 - More flexible data collection - no longer a need for consistent measurement intervals within or between individuals.
 - Consistent parameter comparisons between different data collection schedules - between waves, studies, or both.
 - Differential equations:
 - Parsimonious specification of higher order dynamics such as oscillations and slow changing trends.



- More accuracy by incorporating time explicitly.
- More theoretically justifiable - things happen between measurement occasions.
- More accurate than embedding approaches to differential equations, and no need to set embedding dimension.
- Tangible benefits:
 - More flexible data collection - no longer a need for consistent measurement intervals within or between individuals.
 - Consistent parameter comparisons between different data collection schedules - between waves, studies, or both.
 - Differential equations:
 - Parsimonious specification of higher order dynamics such as oscillations and slow changing trends.
 - Direct interpretation re causality – given true model!



- More accuracy by incorporating time explicitly.
- More theoretically justifiable - things happen between measurement occasions.
- More accurate than embedding approaches to differential equations, and no need to set embedding dimension.
- Tangible benefits:
 - More flexible data collection - no longer a need for consistent measurement intervals within or between individuals.
 - Consistent parameter comparisons between different data collection schedules - between waves, studies, or both.
 - Differential equations:
 - Parsimonious specification of higher order dynamics such as oscillations and slow changing trends.
 - Direct interpretation re causality – given true model!
 - Helps (forces?) us to think more coherently in terms of time.



- More accuracy by incorporating time explicitly.
- More theoretically justifiable - things happen between measurement occasions.
- More accurate than embedding approaches to differential equations, and no need to set embedding dimension.
- Tangible benefits:
 - More flexible data collection - no longer a need for consistent measurement intervals within or between individuals.
 - Consistent parameter comparisons between different data collection schedules - between waves, studies, or both.
 - Differential equations:
 - Parsimonious specification of higher order dynamics such as oscillations and slow changing trends.
 - Direct interpretation re causality – given true model!
 - Helps (forces?) us to think more coherently in terms of time.
- Downsides:



- More accuracy by incorporating time explicitly.
- More theoretically justifiable - things happen between measurement occasions.
- More accurate than embedding approaches to differential equations, and no need to set embedding dimension.
- Tangible benefits:
 - More flexible data collection - no longer a need for consistent measurement intervals within or between individuals.
 - Consistent parameter comparisons between different data collection schedules - between waves, studies, or both.
 - Differential equations:
 - Parsimonious specification of higher order dynamics such as oscillations and slow changing trends.
 - Direct interpretation re causality – given true model!
 - Helps (forces?) us to think more coherently in terms of time.
- Downsides:
 - More mathematically and computationally demanding than autoregressive / latent change approaches.



- Latent dynamic process with measurement model, 1+ subjects, each with 1+ obs, varying time intervals. Dynamics are a linear stochastic differential equation:

$$d\boldsymbol{\eta}(t) = \left(\mathbf{A}\boldsymbol{\eta}(t) + \mathbf{b} + \mathbf{M}\boldsymbol{\chi}(t) \right) dt + \mathbf{G}d\mathbf{W}(t) \quad (1)$$

Observations for each subject are described by:

$$\mathbf{y}(t) = \mathbf{A}\boldsymbol{\eta}(t) + \boldsymbol{\tau} + \boldsymbol{\epsilon}(t) \quad \text{where } \boldsymbol{\epsilon}(t) \sim N(\mathbf{0}_c, \boldsymbol{\Theta}) \quad (2)$$



- Latent dynamic process with measurement model, 1+ subjects, each with 1+ obs, varying time intervals. Dynamics are a linear stochastic differential equation:

$$d\eta(t) = \left(\mathbf{A}\eta(t) + \mathbf{b} + \mathbf{M}\chi(t) \right) dt + \mathbf{G}d\mathbf{W}(t) \quad (1)$$

Observations for each subject are described by:

$$\mathbf{y}(t) = \mathbf{A}\eta(t) + \boldsymbol{\tau} + \epsilon(t) \quad \text{where } \epsilon(t) \sim N(\mathbf{0}_c, \boldsymbol{\Theta}) \quad (2)$$

- The SDE may be solved, for any observation $u \in \mathbf{U}$:

$$\eta_u = \mathbf{A}_u^* \eta_{u-1} + \mathbf{b}_u^* + \mathbf{M}\mathbf{x}_u + \zeta_u^* \quad \zeta_u^* \sim N(\mathbf{0}_v, \mathbf{Q}_u^*) \quad (3)$$

$$\mathbf{A}_u^* = e^{\mathbf{A}(t_u - t_{u-1})} \quad (4)$$

$$\mathbf{b}_u^* = \mathbf{A}^{-1}(\mathbf{A}_u^* - \mathbf{I})\mathbf{b} \quad (5)$$

$$\mathbf{Q}_u^* = \mathbf{Q}_\infty - \mathbf{A}_u^* \mathbf{Q}_\infty \mathbf{A}_u^{*\top} \quad (6)$$

$$\mathbf{Q}_\infty = \text{irow}(-\mathbf{A}_\#^{-1} \text{row}(\mathbf{Q})) \quad (7)$$





Individual differences in dynamics?






Individual differences in dynamics?







- Complete pooling - estimate single fixed effect parameter for entire sample.




- Complete pooling - estimate single fixed effect parameter for entire sample.
 - Contaminates within subject model with between subjects differences, and not a great way to understand individual differences...




- Complete pooling - estimate single fixed effect parameter for entire sample.
 - Contaminates within subject model with between subjects differences, and not a great way to understand individual differences...
- No pooling - assume individuals have no similarities, estimate separate model for each.



- Complete pooling - estimate single fixed effect parameter for entire sample.
 - Contaminates within subject model with between subjects differences, and not a great way to understand individual differences...
- No pooling - assume individuals have no similarities, estimate separate model for each.
 - Simple and perfect if sufficient data exists – 'sufficient' may be extremely large – otherwise prone to finite sample biases and high variance.



- Complete pooling - estimate single fixed effect parameter for entire sample.
 - Contaminates within subject model with between subjects differences, and not a great way to understand individual differences...
- No pooling - assume individuals have no similarities, estimate separate model for each.
 - Simple and perfect if sufficient data exists – ‘sufficient’ may be extremely large – otherwise prone to finite sample biases and high variance.
- Partial pooling - estimate population distribution for individual models.



- Complete pooling - estimate single fixed effect parameter for entire sample.
 - Contaminates within subject model with between subjects differences, and not a great way to understand individual differences...
- No pooling - assume individuals have no similarities, estimate separate model for each.
 - Simple and perfect if sufficient data exists – ‘sufficient’ may be extremely large – otherwise prone to finite sample biases and high variance.
- Partial pooling - estimate population distribution for individual models.
 - More complex models but most flexible - parameters are not either ‘freely varying’ or ‘not varying at all’ but the extent of allowed variation is estimated.





$$p(\Phi, \mu, \mathbf{R}, \beta | \mathbf{Y}, \mathbf{Z}) = \frac{p(\mathbf{Y} | \Phi) p(\Phi | \mu, \mathbf{R}, \beta, \mathbf{Z}) p(\mu, \mathbf{R}, \beta)}{p(\mathbf{Y})} \quad (9)$$

Where subject specific parameters Φ_i are determined in the following manner:

$$\Phi_i = \text{tform}(\mu + \mathbf{R}\mathbf{h}_i + \beta\mathbf{z}_i) \quad (10)$$

$$\mathbf{h}_i \sim \mathbf{N}(\mathbf{0}, \mathbf{1}) \quad (11)$$

$$\mu \sim \mathbf{N}(\mathbf{0}, \mathbf{1}) \quad (12)$$

$$\beta \sim \mathbf{N}(\mathbf{0}, \mathbf{1}) \quad (13)$$



- ctsem - open source R software





- ctsem - open source R software
 - Originally only mixed effects models, using OpenMx as backend.



- ctsem - open source R software
 - Originally only mixed effects models, using OpenMx as backend.
 - Using Stan as backend, added support for fully random effects with Bayesian estimation.



- ctsem - open source R software
 - Originally only mixed effects models, using OpenMx as backend.
 - Using Stan as backend, added support for fully random effects with Bayesian estimation.
- ctModel function constructs a ctsem model based on specified matrices of free and fixed parameters.



- ctsem - open source R software
 - Originally only mixed effects models, using OpenMx as backend.
 - Using Stan as backend, added support for fully random effects with Bayesian estimation.
- ctModel function constructs a ctsem model based on specified matrices of free and fixed parameters.
- Modify between subject effects, covariates, priors etc as needed.



- ctsem - open source R software
 - Originally only mixed effects models, using OpenMx as backend.
 - Using Stan as backend, added support for fully random effects with Bayesian estimation.
- ctModel function constructs a ctsem model based on specified matrices of free and fixed parameters.
- Modify between subject effects, covariates, priors etc as needed.
- ctStanFit function constructs a Stan model and calls rstan for estimation, using either Kalman filter for continuous variables or direct sampling of states for other measurement models.



Latest (undocumented, experimental) developments





- Non-linear dynamics and measurement via unscented Kalman filter.



- Non-linear dynamics and measurement via unscented Kalman filter.
- Binary and ordinal measurement models.



- Non-linear dynamics and measurement via unscented Kalman filter.
- Binary and ordinal measurement models.
- Optimization followed by importance sampling for faster results.



- ctsem and vignettes

<https://cran.r-project.org/web/packages/ctsem/index.html>

- Other articles:

https://www.researchgate.net/profile/Charles_Driver

- Overview of provided R script:

- Generate some data.
- Fit a univariate linear growth curve with random effects and a covariate.
- Add in dynamics.
- Add in an intervention.
- And a second latent process.
- Drop the second latent process and try a state-dependent (non-linear) intervention.