

On a characterization of Cauchy-Stieltjes transforms of compactly supported complex measures and its applications

YAMAGAMI Shigeru

Nagoya University (Emeritus Professor)

Non-Commutative Probability and Related Topics 2024

2024/10/17, Nishijin Plaza in Fukuoka

Contents below are updated on 2024/10/22

Synopsis

Cauchy-Stieltjes transform is one of key notions in analysis of quantum probability and has been investigated from various points of views.

We shall here give a pragmatic characterization of the range of the transform applied to compactly supported complex measures in the real line, which enables us to identify certain holomorphic functions with Cauchy-Stieltjes transforms.

As a simple application of the characterization, we work out a spectral analysis of linear functionals of some polynomial hypergroups parametrized by $0 < r \leq 1/2$.

When $r = 1/2l$ with l a positive integer, it is isomorphic to the commutative algebra of radial functions on the free group of l generators and the spectral properties of Haagerup's positive definite functions are then described explicitly, which turns out to be useful in establishing Plancherel type formulae.

Cauchy-Stieltjes Transform

Given a complex measure $\mu(dt)$ on \mathbb{R} ,
its **Cauchy-Stieltjes transform** $C(z)$ is a holomorphic function of
 $z = x + iy \in \mathbb{C} \setminus \mathbb{R}$ ($y \neq 0$) defined by

$$C(z) = \int \frac{1}{t - z} \mu(dt) = \int \frac{1}{t - x - iy} \mu(dt).$$

From

$$\left| \frac{y}{t - x - iy} \right| \leq 1 \quad (y \neq 0), \quad \lim_{y \rightarrow \pm\infty} \frac{y}{t - x - iy} = i$$

and

$$\lim_{y \rightarrow 0} \frac{y}{t - x - iy} = \begin{cases} i & (t = x) \\ 0 & (t \neq x) \end{cases},$$

we have $|yC(x + iy)| \leq |\mu|(\mathbb{R})$ and

$$\lim_{y \rightarrow \pm\infty} yC(x + iy) = i\mu(\mathbb{R}), \quad \lim_{y \rightarrow 0} yC(x + iy) = i\mu(\{x\}).$$

When μ is a signed measure, C is **real** in the sense that

$$\overline{C(z)} = C(\bar{z}) \quad (z \in \mathbb{C} \setminus \mathbb{R}).$$

Theorem (Stieltjes Inversion Formula)

The complex measure μ is recovered from its Cauchy-Stieltjes transform $C(z)$ as a Radon measure on $C_c(\mathbb{R})$ by

$$2\pi i \mu(dx) = \lim_{y \rightarrow +0} \left(C(x + iy) - C(x - iy) \right) dx.$$

The support $[\mu]$ of μ can be read off from $C(z)$ easily:

For $x \in \mathbb{R}$,

$x \notin [\mu] \iff C(z)$ is analytic in a neighborhood of $x \in \mathbb{C}$.

Given a holomorphic function $\varphi(z)$ of $z \in \mathbb{C} \setminus \mathbb{R}$, its **support** is

$$[\varphi] = \{c \in \overline{\mathbb{C}}; \varphi(z) \text{ is not analytic at } z = c\}.$$

Thus $[C]$ is the closure of $[\mu] \subset \mathbb{R}$ in the Riemann sphere $\overline{\mathbb{C}}$.

Example

For a Dirac measure $\delta(t - c)$ at $c \in \mathbb{R}$,

$$C(z) = \frac{1}{c - z}, \quad [C] = \{c\},$$

$$\begin{aligned} 2\pi i \delta(x - c) dx &= \lim_{y \rightarrow +0} \left(\frac{1}{c - x - iy} - \frac{1}{c - x + iy} \right) dx \\ &= \lim_{y \rightarrow +0} \frac{2iy}{(c - x)^2 + y^2} dx \end{aligned}$$

Observation:

If C is the Cauchy-Stieltjes transform of a complex measure, isolated singularity of $C(z)$ at a real point must be a simple pole because higher order poles give rise to derivatives of $\delta(x - c)$.

Theorem (cf. Akhiezer, §3.1)

A real holomorphic function $C(z)$ of $z \in \mathbb{C} \setminus \mathbb{R}$ is the Cauchy-Stieltjes transform of a positive (non-zero) finite measure if and only if the following conditions are satisfied.

- 1 $\operatorname{Im} C(z) > 0$ ($\operatorname{Im} z > 0$) and
- 2 $\{yC(iy); y \geq 1\}$ is a bounded set in \mathbb{C} .

From the expression

$$C(x+iy) = \int \frac{t-x}{(t-x)^2 + y^2} \mu(dt) + i \int \frac{y}{(t-x)^2 + y^2} \mu(dt),$$

the conditions are clearly necessary and non-trivial is sufficiency, which can be deduced from the integral representation of $C(z)$ satisfying the positivity condition (i).

Theorem

A holomorphic function $C(z)$ of $z \in \mathbb{C} \setminus \mathbb{R}$ is the Cauchy-Stieltjes transform of a compactly supported complex measure if and only if the following conditions are satisfied.

- 1 $C(z)$ is analytically extended to a neighborhood of $\infty \in \overline{\mathbb{C}}$ so that $C(\infty) = 0$.
- 2 The limit $\lim_{y \rightarrow +0} (C(x + iy) - C(x - iy)) dx$ exists as a complex Radon measure (as a linear functional) on $C_c(\mathbb{R})$.

Notice that a complex measure on \mathbb{R} is determined by the integration on $C_c(\mathbb{R})$ in the second condition.

The first condition implies that $C(x + iy) - C(x - iy)$ is integrable as a function of $x \in \mathbb{R}$ for any $0 \neq y \in \mathbb{R}$ and the limit in the second condition can be strengthened to functions in $C_0(\mathbb{R})$.

Example

For the Dirac mass at $c \in \mathbb{R}$, $C(z) = \frac{1}{c-z}$ and

$$C(x + iy)dx = \frac{c - x + iy}{(c - x)^2 + y^2} dx$$

converges for $y \rightarrow \pm 0$ as a linear functional on $C_c^1(\mathbb{R})$
but not as a Radon measure on $C_c(\mathbb{R})$.

However, the combination

$$C(x + iy) - C(x - iy) = \frac{2iy}{(c - x)^2 + y^2}$$

is an integrable function of $x \in \mathbb{R}$ for each $y > 0$ and the
complex measure $(C(x + iy) - C(x - iy))dx$ on \mathbb{R} converges
to $2\pi\delta(x - c)$ as a linear functional on $C_c(\mathbb{R})$ for $y \rightarrow +0$.

$F = \langle f_1, \dots, f_l \rangle$: a free group of l -generators.

$F_n = \{g \in F; |g| = n\}$: the set of words of length n .

$F_1 = \{f_1^{\pm 1}, \dots, f_l^{\pm 1}\}$ and $|F_n| = 2l(2l-1)^{n-1}$ for $n \geq 1$.

$\mathbb{C}F = \sum_{g \in G} \mathbb{C}g$: the algebraic group algebra ($g^* = g^{-1}$).

Radial elements $h_n = h_n^* \in \mathbb{C}F$ are defined to be

$$h_n = \frac{1}{|F_n|} \sum_{g \in F_n} g \quad \text{for } n \geq 0$$

with $h_0 = e$ the unit element in $\mathbb{C}F$, which satisfy

$$h_1 h_n = \frac{1}{2l} h_{n-1} + \frac{2l-1}{2l} h_{n+1} = h_n h_1$$

and give a linear basis of a commutative $*$ -subalgebra

$$\sum_{n \geq 0} \mathbb{C}h_n \subset \mathbb{C}F.$$

Polynomial Hypergroups

Given a real $1 \neq r \in \mathbb{R}$, let $\mathcal{A} = \sum_{n \geq 0} \mathbb{C}h_n$ be a universal $*$ -algebra with h_0 the unit element and $h_n = h_n^*$ satisfying

$$h_1 h_n = r h_{n-1} + (1 - r) h_{n+1} \quad (n \geq 1),$$

which is $*$ -isomorphic to the polynomial algebra $\mathbb{C}[t]$ by the correspondence $h_n \leftrightarrow P_n(t)$.

Here $P_n(t)$ is a polynomial of degree n specified by the recurrence relation

$$t P_n(t) = r P_{n-1}(t) + (1 - r) P_{n+1}(t)$$

with the initial condition $P_0(t) = 1$ and $P_1(t) = t$.

Generally a $*$ -algebra with a distinguished basis $\{h_n = h_n^*\}_{n \geq 0}$ with h_0 the unit element is called a (hermitian) **hypergroup** if, in the expansion $h_j h_k = \sum_{n \geq 0} c_{j,k,n} h_n$, coefficients satisfy

$$c_{j,k,n} \geq 0, \quad \sum_n c_{j,k,n} = 1 \quad \text{and} \quad c_{j,k,0} > 0 \iff j = k.$$

For the $*$ -algebra \mathcal{A} , coefficients are computed easily to see that

$$\mathcal{A} = \sum_{n \geq 0} \mathbb{C} h_n \text{ is a hypergroup } \iff 0 < r \leq \frac{1}{2}.$$

Note that $0 < r = 1/2l \leq 1/2$ for radial elements in $\mathbb{C}F$.

The hypergroup \mathcal{A} is completed to a Banach $*$ -algebra with respect to the ℓ^1 -norm

$$\left\| \sum_{n \geq 0} \alpha_n h_n \right\|_1 = \sum_{n \geq 0} |\alpha_n|.$$

Let A be the accompanied commutative C^* -algebra.
In the case of radial elements, we see that $A \subset C^*(F)$.

Under the correspondence

a linear functional $\phi : \mathcal{A} \rightarrow \mathbb{C} \iff$ a sequence $(\phi_n = \phi(h_n))$,

- ① ϕ is ℓ^1 -bounded $\iff (\phi_n)$ is bounded.
- ② A multiplicative functional $\phi \iff c = \phi(h_1) \in \mathbb{C}$.
- ③ A $*$ -homomorphism $\phi \iff c = \phi(h_1) \in \mathbb{R}$.
- ④ A bounded $*$ -homomorphism $\phi \iff c = \phi(h_1) \in [-1, 1]$.

The spectrum $\sigma(h_1)$ of h_1 in A is therefore $[-1, 1]$ and

$$A = C^*(h_0, h_1) \cong C([-1, 1]).$$

Analytic Functionals

Definition

A linear functional $\phi : \mathcal{A} \rightarrow \mathbb{C}$ is **analytic** if the generating function $\phi(z) = \sum_{n \geq 0} \phi_n z^n$ converges at some $0 \neq z \in \mathbb{C}$.

Any C^* -bounded functional ϕ is analytic: $\|a\| \leq \|a\|_1$ ($a \in \mathcal{A}$), whence (ϕ_n) is bounded and $\sum_{n \geq 0} \phi_n z^n$ converges for $|z| < 1$. Moreover ϕ is represented by a complex Radon measure μ on $[-1, 1]$ in such a way that

$$\phi(h_n) = \int_{[-1,1]} P_n(t) \mu(dt).$$

The C^* -boundedness of a given sequence $(\phi(h_n))$ is then equivalent to finding μ .

Thanks to the recurrence relation, the generating function

$$P(z, t) = \sum_{n \geq 0} z^n P_n(t)$$

is expressed by

$$P(z, t) = \frac{1 - r - rzt}{1 - r - zt + rz^2}$$

and we see that

$$\begin{aligned} \phi(z) &= \sum_{n \geq 0} \phi(h_n) z^n = \int_{[-1,1]} \frac{1 - r - rzt}{1 - r - zt + rz^2} \mu(dt) \\ &= r\phi_0 + \frac{r^2 z^2 - (1 - r)^2}{z} \int_{[-1,1]} \frac{1}{t - rz - (1 - r)/z} \mu(dt). \end{aligned}$$

Introduce a new variable by $w = rz + (1 - r)/z \iff$

$$z = \frac{w - \sqrt{w^2 - 4r(1 - r)}}{2r} \sim \frac{1 - r}{w} \quad (w \rightarrow \infty).$$

so that

$$\int_{[-1,1]} \frac{1}{t - w} \mu(dt) = \frac{z}{r^2 z^2 - (1 - r)^2} (\phi(z) - r\phi_0).$$

Thus finding μ is reduced to the validity of the Stieltjes inversion formula for the right hand side.

Notice here that, given a complex Radon measure μ on $[-1, 1]$, the left hand side is analytic in $1/w$ and vanishes at $w = \infty$, which in turn determines an analytic function $\phi(z)$ of z around $z = 0$ by equating it with the right hand side.

Recall that the Cauchy-Stieltjes transform

$$C(w) = \int_{[-1,1]} \frac{1}{t-w} \mu(dt)$$

of μ is holomorphic on $\overline{\mathbb{C}} \setminus [-1, 1]$ and vanishes at $w = \infty$.

The Stieltjes inversion formula: In the weak* topology, we have

$$2\pi i \mu(dt) = \lim_{\epsilon \rightarrow +0} \left(C(t + i\epsilon) - C(t - i\epsilon) \right) dt.$$

Example

When $\phi(z) \equiv 1$, μ is the Kesten measure:

$$C(w) = \frac{(2r-1)w + \sqrt{w^2 - 4r(1-r)}}{2r(1-w^2)}$$
$$\iff \mu(dt) = \frac{1}{2\pi r} \frac{\sqrt{4r(1-r) - t^2}}{1-t^2} dt.$$

Functionals of Geometric Series

Given $v \in \mathbb{C}$, let

$$\phi(z) = \frac{1}{1-vz} = \sum_{n=0}^{\infty} v^n z^n \iff \phi_n = \phi(h_n) = v^n \ (n \geq 0).$$

Here comes into the Joukowski transform of v :

$$c_r(v) = r \frac{1}{v} + (1-r)v.$$

The “Cauchy-Stieltjes transform” $C(w)$ of ϕ then takes the form

$$\frac{v^{-1} - v}{2} \frac{(2r-1)w + \sqrt{w^2 - 4r(1-r)}}{(1-w^2)(c_r(v) - w)} + \frac{1}{c_r(v) - w},$$

which is reduced at $v = 0$ to

$$C(w) = \frac{(2r-1)w + \sqrt{w^2 - 4r(1-r)}}{2r(1-w^2)}.$$

Remark

- ① If $\mathcal{A} \subset \mathbb{C}F$ consists of radial elements ($r = 1/2l$) and $0 < v < 1$, ϕ is the restriction of the Haagerup's positive definite function (a state on the group algebra) to $\mathcal{A} \subset \mathbb{C}F$.
- ② When $v = 0$, C is the Cauchy-Stieltjes transform of the Kesten measure.
- ③ H. Yoshida found that

$$-C(w) = \frac{1}{w - v - \frac{r(1 - v^2)}{w + rv - \frac{r(1 - r)}{w - \frac{r(1 - r)}{w - \ddots}}}}.$$

Theorem

The functional ϕ of geometric series is C^* -bounded if and only if v falls into one of the following cases.

- ① $|v| < \sqrt{r/(1-r)}$ or $v = \pm\sqrt{r/(1-r)}$,

$$\mu_c(dt) = \frac{v^{-1} - v}{2\pi} \frac{\sqrt{4r(1-r) - t^2}}{(1-t^2)(c_r(v) - t)} dt.$$

- ② $v \in \mathbb{R}$ and $\sqrt{r/(1-r)} < |v| \leq 1$
($c_r(v) \in \mathbb{R}$ and $2\sqrt{r(1-r)} < |c_r(v)| \leq 1$ then),

$$\mu(dt) = \mu_c(dt) + \omega(v)\delta(t - c_r(v)), \quad \omega(v) = \frac{1 - c_r(v^2)}{1 - c_r(v)^2}.$$

A C^* -bounded ϕ is positive if and only if $-1 \leq v \leq 1$.




Figure: v -region of C^* -boundedness

Corollary (Free Poisson as a normalized limit)

In terms of the normalized parameter $s = \frac{t - v}{\sqrt{r(1 - v^2)}}$,
the normalized distribution converges to $\nu(ds)$ under the limit
 $(r, v) \rightarrow (0, 0)$ keeping $\gamma = v/\sqrt{r} \in \mathbb{R}$ constant:

- ① For $-1 \leq \gamma \leq 1$,

$$\nu(ds) = \frac{\sqrt{4 - (s + \gamma)^2}}{2\pi(1 - \gamma s)} ds.$$

- ② For $|\gamma| \geq 1$,

$$\nu(ds) = \frac{\sqrt{4 - (s + \gamma)^2}}{2\pi(1 - \gamma s)} ds + \left(1 - \frac{1}{\gamma^2}\right) \delta(s - 1/\gamma).$$

Sketch of the Proof

Introduce the regular spectrum σ_r by

$$\sigma_r = [-2\sqrt{r(1-r)}, 2\sqrt{r(1-r)}] \subset [-1, 1]$$

and observe that

$$\sqrt{w^2 - 4r(1-r)} = w \sqrt{1 - \frac{4r(1-r)}{w^2}}$$

vanishes at $w = \infty$ and is holomorphic on $\bar{\mathbb{C}} \setminus \sigma_r$.

The same property holds for the Kesten part

$$\frac{(2r-1)w + \sqrt{w^2 - 4r(1-r)}}{1-w^2}.$$

The possible singularity of $C(w)$ other than $\pm 2\sqrt{r(1-r)}$ is

$$c_r(v) = \frac{r}{v} + (1-r)v.$$

Recall some properties of the Joukowski transform $c_r(v)$ of v :

- ① Invariance under the inversion $v \mapsto \frac{r}{1-r} \frac{1}{v}$

with critical fixed points

$$v = \pm \sqrt{\frac{r}{1-r}} \iff c_r(v) = \pm 1.$$

- ② Both $|v| < \sqrt{\frac{r}{1-r}}$ and $|v| > \sqrt{\frac{r}{1-r}}$ (including $v = \infty$) are biholomorphically mapped onto a slit sphere $\bar{\mathbb{C}} \setminus \sigma_r$.

- ③ Circle $|v| = \rho \neq \sqrt{\frac{r}{1-r}}$ is mapped to an ellipse of vertices $\frac{r}{\rho} + (1-r)\rho$, $\frac{r}{\rho} - (1-r)\rho$,

which shrinks to the segment σ_r when ρ approaches $\sqrt{\frac{r}{1-r}}$.

- ④ $-1 \leq c_r(v) \leq 1$ and $|v| > \sqrt{\frac{r}{1-r}}$

$$\iff v \in \mathbb{R} \text{ and } \sqrt{\frac{r}{1-r}} < |v| \leq 1.$$

Singularity criterion of $c_r(v)$:

$$\lim_{w \rightarrow c_r(v)} (c_r(v) - w)C(w) = \begin{cases} 0 & (|v| < \sqrt{r/(1-r)}) \\ \omega(v) & (|v| > \sqrt{r/(1-r)}) \end{cases}.$$

Combined with properties of Joukowski transform,
 C is holomorphic on $\overline{\mathbb{C}} \setminus [-1, 1]$ if and only if the following
 alternatives hold:

- ① $|v| \leq \sqrt{\frac{r}{1-r}}$. Then C is holomorphic on $\overline{\mathbb{C}} \setminus \sigma_r$.
- ② $\sqrt{\frac{r}{1-r}} < \pm v \leq 1 \iff 2\sqrt{r(1-r)} < \pm c_r(v) \leq 1$.
 Then C is meromorphic on $\overline{\mathbb{C}} \setminus \sigma_r$ with $c_r(v)$ the unique
 simple pole.

The characterization of Cauchy-Stieltjes transform is applied to the following cases:

- ① $|v| < \sqrt{r/(1-r)}$. A continuous measure on σ_r appears.
- ② $\sqrt{r/(1-r)} < \pm v \leq 1$. Adding to the continuous measure on σ_r , an atomic measure appears at $c_r(v) \in [-1, 1] \setminus \sigma_r$.
- ③ $|v| = \sqrt{r/(1-r)}$ but $v \neq \pm\sqrt{r/(1-r)}$.
 $C(w)$ is not a Cauchy-Stieltjes transform.
- ④ In the critical case
 $v = \pm\sqrt{r/(1-r)} \iff c_r(v) = \pm 2\sqrt{r(1-r)}$,
the limit measure appears from (1) or (2).

The critical case: $c_r(v) = \pm c$ with $c = 2\sqrt{r(1-r)}$.

$$\frac{\sqrt{w^2 - 4r(1-r)}}{c_r(v) - w} = -\sqrt{\frac{w \pm c}{w \mp c}}.$$

We may assume $c_r(v) = c \iff v = \sqrt{\frac{r}{1-r}}$ by symmetry.

By an obvious estimate

$$\left| \frac{t + is + c}{t + is - c} \right| \leq \frac{|t + c| + |s|}{|s|},$$

$$s \left| \sqrt{\frac{t + is + c}{t + is - c}} \right| \leq \sqrt{|s|(|t + c| + |s|)}$$

is bounded for a bounded (t, s) and converges to 0 as $s \rightarrow 0$ for any $t \in \mathbb{R}$.

From another estimate

$$\left| \frac{t + is + c}{t + is - c} \right| \leq \frac{|t + c| + |s|}{|t - c|},$$

$\left| \sqrt{\frac{w+c}{w-c}} \right|$ is majorized uniformly in $0 < |s| \leq 1$ by a locally integrable function $\sqrt{(|t + c| + 1)/|t - c|}$ of $t \in \mathbb{R}$ and

$$\lim_{s \rightarrow \pm 0} \sqrt{\frac{t + is + c}{t + is - c}} = \begin{cases} \sqrt{\frac{t+c}{t-c}} & (t > c) \\ \mp i \sqrt{\frac{t+c}{c-t}} & (-c < t < c) , \\ \sqrt{\frac{-c-t}{c-t}} & (t < -c) \end{cases}$$

whence

$$\lim_{s \rightarrow +0} \left(\sqrt{\frac{t + is + c}{t + is - c}} - \sqrt{\frac{t - is + c}{t - is - c}} \right) dt = -2i \sqrt{\frac{t + c}{c - t}} dt$$

as a Radon measure (supported by σ_r) on $C_c(\mathbb{R})$.

References

N.I. Akhiezer, The Classical Moment Problem, Oliver & Boyd (1965).

A. Hora and N. Obata, Quantum Probability and Spectral Analysis of Graphs, Springer (2007).

J. Mingo and R. Speicher, Free Probability and Random Matrices, Springer (2017).

S. Yamagami, On a moment problem of analytic functionals of polynomial hypergroups, arXiv:1912.03895.

S. Yamagami and H. Yoshida, On shifting semicircular roots, arXiv:2209.04585.