

Numerical Linear Algebra Assignment 9

Exercise 1. (10 points)

Consider the simultaneous iteration:

Algorithm 1: Simultaneous iteration

Pick $\mathbf{Q}_n^{(0)} \in \mathbb{C}^{m \times n}$ with orthonormal columns
for $k = 1, 2, 3, \dots$,
 $\mathbf{Z}^{(k)} = \mathbf{A}\mathbf{Q}_n^{(k-1)}$
 $\mathbf{Q}_n^{(k)}\mathbf{R}_n^{(k)} = \mathbf{Z}^{(k)} \quad (\text{QR factorization})$
end

Assume $\mathbf{A} = \mathbf{S}\mathbf{\Lambda}\mathbf{S}^{-1}$ is diagonalizable with $\mathbf{\Lambda} = \text{diag}\{\lambda_1, \lambda_2, \dots, \lambda_m\}$ and

$$|\lambda_1| \geq \dots \geq |\lambda_n| > |\lambda_{n+1}| \geq \dots \geq |\lambda_m|.$$

Assume $\mathbf{X}_n^{(0)} := [\mathbf{I}_n \quad \mathbf{0}] \mathbf{S}^{-1} \mathbf{Q}_n^{(0)}$ has full rank. Prove that

$$\text{span}\{\mathbf{Q}_n^{(k)}\} = \text{span}\{\mathbf{A}^k \mathbf{Q}_n^{(0)}\}.$$

Exercise 2. (Zhihao Cao, 10 points)

Let $\mathbf{A}_1 = \begin{bmatrix} a & b \\ \varepsilon & c \end{bmatrix} \in \mathbb{R}^{2 \times 2}$. Let $\mathbf{A}_2 = \mathbf{R}_1 \mathbf{Q}_1 + c\mathbf{I} = \begin{bmatrix} a_{11} & a_{12} \\ a_{21} & a_{22} \end{bmatrix}$, where $\mathbf{Q}_1 \mathbf{R}_1 = \mathbf{A}_1 - c\mathbf{I}$ is a QR factorization of $\mathbf{A}_1 - c\mathbf{I}$. Prove:

- (a) if a and c are not close (i.e., $\exists \delta > 0, |a - c| > \delta$), then $a_{21} = \mathcal{O}(\varepsilon^2)$;
- (b) if further $b = \varepsilon$ (i.e., \mathbf{A}_1 is symmetric), then $a_{21} = \mathcal{O}(\varepsilon^3)$.

Exercise 3. (10 points)

Assume that $\mathbf{A} \in \mathbb{R}^{n \times n}$ is nonsingular. Let $\mathbf{A} = \mathbf{Q}\mathbf{R}$ be a QR factorization. Prove that if \mathbf{A} is symmetric and has bandwidth $2p + 1$ (i.e., $a_{ij} = 0$ for $|i - j| > p$), then $\mathbf{B} = \mathbf{R}\mathbf{Q}$ is also symmetric and has bandwidth $2p + 1$.

Compulsory requirement for programming: Use Matlab's publish to save all your code, comments, and results to a PDF file. You must use the programming format files: example.format.zip.

Programming 1. (10 points)

Write a function `[Q,H] = myhess(A)` that reduces a complex square matrix to upper Hessenberg form by unitary similarity transformations, i.e., $\mathbf{H} = \mathbf{Q}^* \mathbf{A} \mathbf{Q}$. Your program should use only elementary Matlab operations – not the function `hess`, for example. Apply your program to $\mathbf{A} = \text{randn}(5) + 1i * \text{randn}(5)$.

Programming 2. (TreBau Exercise 29.1, 10 points)