Lecture 6: Stationary iterative methods



School of Mathematical Sciences, Xiamen University

1. Splitting and stationary iterative method

Definition 1

A *splitting* of $\mathbf{A} \in \mathbb{C}^{n\times n}$ is a decomposition $\mathbf{A} = \mathbf{M} - \mathbf{K}$, with $\mathbf{M}$ nonsingular.

Remark 2

A splitting yields a stationary iterative method as follows. The equation

$$\mathbf{Ax} = (\mathbf{M} - \mathbf{K})\mathbf{x} = \mathbf{b}$$

implies

$$\mathbf{x} = \mathbf{M}^{-1}\mathbf{Kx} + \mathbf{M}^{-1}\mathbf{b} := \mathbf{Rx} + \mathbf{c}.$$

Given a starting vector $\mathbf{x}^{(0)}$*, we obtain a stationary iterative method*

$$\mathbf{x}^{(m)} = \mathbf{Rx}^{(m-1)} + \mathbf{c}, \quad m = 1, 2, \ldots$$

We note that $\mathbf{R} = \mathbf{M}^{-1}\mathbf{K} = \mathbf{M}^{-1}(\mathbf{M} - \mathbf{A}) = \mathbf{I} - \mathbf{M}^{-1}\mathbf{A}$.

2. Convergence criterion

Definition 3

The spectral radius of a matrix $\mathbf{A} \in \mathbb{C}^{n\times n}$ is $\rho(\mathbf{A}) = \max\limits_{\lambda\in\Lambda(\mathbf{A})} |\lambda|$.

Exercise. Assume that $\mathbf{A}$ is singular and has a splitting $\mathbf{A} = \mathbf{M} - \mathbf{K}$ with $\mathbf{M}$ nonsingular. Prove that the spectral radius $\rho(\mathbf{M}^{-1}\mathbf{K}) \geq 1$.

Proposition 4

Let $\|\cdot\|$ denote a matrix norm on $\mathbb{C}^{n\times n}$ induced by a vector norm $\|\cdot\|$ on $\mathbb{C}^n$. For any $\mathbf{A} \in \mathbb{C}^{n\times n}$, we have $\rho(\mathbf{A}) \leq \|\mathbf{A}\|$.

Lemma 5

For any given $\mathbf{A} \in \mathbb{C}^{n\times n}$ and $\varepsilon > 0$ there exists an induced matrix norm $\|\cdot\|_\star$ (induced by a vector norm $\|\cdot\|_\star$ on $\mathbb{C}^n$) such that

$$\rho(\mathbf{A}) \leq \|\mathbf{A}\|_\star \leq \rho(\mathbf{A}) + \varepsilon.$$

The norm $\|\cdot\|_\star$ depends on both $\mathbf{A}$ and ε.

Proof.

Let $\mathbf{A} = \mathbf{SJS}^{-1}$ be a Jordan form of $\mathbf{A}$. Let

$$\mathbf{D}_\varepsilon = \mathrm{diag}\{1, \varepsilon, \varepsilon^2, \cdots, \varepsilon^{n-1}\}.$$

Now for all $\mathbf{x} \in \mathbb{C}^n$ and for all $\mathbf{B} \in \mathbb{C}^{n\times n}$, define the vector norm

$$\|\mathbf{x}\|_\star := \|(\mathbf{SD}_\varepsilon)^{-1}\mathbf{x}\|_\infty$$

and the corresponding induced matrix norm

$$\begin{aligned}
\|\mathbf{B}\|_\star :&= \sup_{\mathbf{x}\in\mathbb{C}^n, \mathbf{x}\neq\mathbf{0}} \frac{\|\mathbf{Bx}\|_\star}{\|\mathbf{x}\|_\star} = \sup_{\mathbf{x}\in\mathbb{C}^n, \mathbf{x}\neq\mathbf{0}} \frac{\|(\mathbf{SD}_\varepsilon)^{-1}\mathbf{Bx}\|_\infty}{\|(\mathbf{SD}_\varepsilon)^{-1}\mathbf{x}\|_\infty} \\
&= \sup_{\mathbf{y}\in\mathbb{C}^n, \mathbf{y}\neq\mathbf{0}} \frac{\|(\mathbf{SD}_\varepsilon)^{-1}\mathbf{B}(\mathbf{SD}_\varepsilon)\mathbf{y}\|_\infty}{\|\mathbf{y}\|_\infty} \\
&= \|\mathbf{D}_\varepsilon^{-1}\mathbf{S}^{-1}\mathbf{BSD}_\varepsilon\|_\infty.
\end{aligned}$$

The statement follows from $\|\mathbf{A}\|_\star = \|\mathbf{D}_\varepsilon^{-1}\mathbf{JD}_\varepsilon\|_\infty \leq \rho(\mathbf{A}) + \varepsilon$. □

Theorem 6

The stationary iteration $\mathbf{x}^{(m)} = \mathbf{R}\mathbf{x}^{(m-1)} + \mathbf{c}$ *converges to the solution of* $(\mathbf{I} - \mathbf{R})\mathbf{x} = \mathbf{c}$ *for all starting vectors* $\mathbf{x}^{(0)}$ *if and only if* $\rho(\mathbf{R}) < 1$.

Proof.

For all $\mathbf{x}^{(0)}$, *we have* $\mathbf{x}^{(m)} - \mathbf{x} = \mathbf{R}(\mathbf{x}^{(m-1)} - \mathbf{x}) = \cdots = \mathbf{R}^m(\mathbf{x}^{(0)} - \mathbf{x})$.

" $\Leftarrow$ " : *If* $\rho(\mathbf{R}) < 1$, *by Lemma 5 there exists an induced matrix norm* $\|\cdot\|_\star$ *such that* $\|\mathbf{R}\|_\star < 1$, *then* $\|\mathbf{x}^{(m)} - \mathbf{x}\|_\star \leq \|\mathbf{R}\|_\star^m \|\mathbf{x}^{(0)} - \mathbf{x}\|_\star \to 0$ *for all* $\mathbf{x}^{(0)}$, *i.e.,* $\mathbf{x}^{(m)} \to \mathbf{x}$ *as* $m \to \infty$.

" $\Rightarrow$ " : *Proof by contradiction. Let* $\mathbf{x}^{(0)} - \mathbf{x}$ *be an eigenvector of* $\mathbf{R}$ *corresponding to eigenvalue* λ *with* $|\lambda| = \rho(\mathbf{R})$. *If* $\rho(\mathbf{R}) \geq 1$, *then* $\mathbf{x}^{(m)} - \mathbf{x} = \lambda^m(\mathbf{x}^{(0)} - \mathbf{x})$ *will not approach* $\mathbf{0}$. *This is a contradiction.* □

Remark 7

A good stationary iterative method satisfies both (1) $\rho(\mathbf{R})$ *is small* (< 1) *and (2) the action* $\mathbf{R}\mathbf{x}^{(m-1)} + \mathbf{c}$ *is computationally inexpensive.*

- (1) and (2) are conflicting goals, and need to be balanced.
- If $\Lambda(\mathbf{R}) \subset [-\rho(\mathbf{R}), \rho(\mathbf{R})]$, then Chebyshev acceleration technique can be used. See Demmel's book ANLA, section 6.5.6.
- The iterative matrix $\mathbf{R}$ is only theoretically useful and usually will not be formed explicitly.

3. Classical stationary iterative methods

- Let $\mathbf{A} = \mathbf{D} - \mathbf{L} - \mathbf{U}$, where

  $\mathbf{D}$ is the diagonal matrix with diagonal entries $d_{ii} = a_{ii}$,

  $-\mathbf{L}$ is the strictly lower triangular part of $\mathbf{A}$,

  $-\mathbf{U}$ is the strictly upper triangular part of $\mathbf{A}$.
- Assume that $\mathbf{A}$ has no zero diagonal entries. We will introduce

  (1) Jacobi's method,

  (2) Gauss–Seidel method,

  (3) Successive overrelaxation: SOR(ω),

  (4) Symmetric successive overrelaxation: SSOR(ω).

3.1. Jacobi's method

- The splitting of Jacobi's method is $\mathbf{A} = \mathbf{D} - (\mathbf{L} + \mathbf{U})$. The corresponding $\mathbf{R}$ and $\mathbf{c}$ are

$$\mathbf{R} = \mathbf{D}^{-1}(\mathbf{L} + \mathbf{U}) = \mathbf{I} - \mathbf{D}^{-1}\mathbf{A}, \qquad \mathbf{c} = \mathbf{D}^{-1}\mathbf{b}.$$

Algorithm 1: Jacobi's method (from $\mathbf{x}^{(m)}$ to $\mathbf{x}^{(m+1)}$)

---

for $j = 1$ **to** n

$$x_j^{(m+1)} = \frac{1}{a_{jj}} \left(b_j - \sum_{k \neq j} a_{jk} x_k^{(m)} \right)$$

end

Theorem 8

If $\mathbf{A}$ *is Hermitian and* $a_{ii} > 0$ *for all* i*, then Jacobi's method converges for all starting vectors if and only if* $\mathbf{A}$ *and* $2\mathbf{D} - \mathbf{A}$ *are both* HPD.

3.2. Gauss–Seidel method

- The splitting of Gauss–Seidel method is

$$\mathbf{A} = (\mathbf{D} - \mathbf{L}) - \mathbf{U}.$$

The corresponding $\mathbf{R}$ and $\mathbf{c}$ are

$$\mathbf{R} = (\mathbf{D} - \mathbf{L})^{-1}\mathbf{U}$$

and

$$\mathbf{c} = (\mathbf{D} - \mathbf{L})^{-1}\mathbf{b}.$$

Algorithm 2: Gauss–Seidel method (from $\mathbf{x}^{(m)}$ to $\mathbf{x}^{(m+1)}$)

---

for $j = 1$ **to** n

$$x_j^{(m+1)} = \frac{1}{a_{jj}} \left(b_j - \sum_{k=1}^{j-1} a_{jk} x_k^{(m+1)} - \sum_{k=j+1}^{n} a_{jk} x_k^{(m)} \right)$$

end

3.3. Successive overrelaxation: SOR(ω), $\omega \in \mathbb{R}$

- The splitting of SOR(ω) is $\omega\mathbf{A} = (\mathbf{D} - \omega\mathbf{L}) - ((1-\omega)\mathbf{D} + \omega\mathbf{U})$. The corresponding $\mathbf{R}$ and $\mathbf{c}$ are

$$\mathbf{R} = (\mathbf{D} - \omega\mathbf{L})^{-1}((1-\omega)\mathbf{D} + \omega\mathbf{U}),$$

and

$$\mathbf{c} = \omega(\mathbf{D} - \omega\mathbf{L})^{-1}\mathbf{b}.$$

(i) $\omega = 1$: Gauss–Seidel method;
(ii) $0 < \omega < 2$: necessary condition for convergence (Theorem 13);
(iii) Optimal ω: see Ciaramella and Gander's book (Chapter 2).

Algorithm 3: SOR(ω), here ω is the relaxation parameter

---

for $j = 1$ **to** n

$$x_j^{(m+1)} = (1-\omega)x_j^{(m)} + \frac{\omega}{a_{jj}}\left(b_j - \sum_{k=1}^{j-1} a_{jk}x_k^{(m+1)} - \sum_{k=j+1}^{n} a_{jk}x_k^{(m)}\right)$$

end

3.4. Symmetric successive overrelaxation: SSOR(ω), $\omega \in \mathbb{R}$

- This method uses two splittings:

$$\begin{aligned}\omega\mathbf{A} &= (\mathbf{D} - \omega\mathbf{L}) - ((1-\omega)\mathbf{D} + \omega\mathbf{U}) \\ &= (\mathbf{D} - \omega\mathbf{U}) - ((1-\omega)\mathbf{D} + \omega\mathbf{L}).\end{aligned}$$

The corresponding $\mathbf{R}$ and $\mathbf{c}$ are

$$\mathbf{R} = (\mathbf{D} - \omega\mathbf{U})^{-1}((1-\omega)\mathbf{D} + \omega\mathbf{L})(\mathbf{D} - \omega\mathbf{L})^{-1}((1-\omega)\mathbf{D} + \omega\mathbf{U}),$$

$$\mathbf{c} = \omega(2-\omega)(\mathbf{D} - \omega\mathbf{U})^{-1}\mathbf{D}(\mathbf{D} - \omega\mathbf{L})^{-1}\mathbf{b}.$$

Algorithm 4: SSOR(ω)

for $j = 1$ **to** n

$$x_j^{(m+1/2)} = (1-\omega)x_j^{(m)} + \frac{\omega}{a_{jj}}\left(b_j - \sum_{k=1}^{j-1} a_{jk}x_k^{(m+1/2)} - \sum_{k=j+1}^{n} a_{jk}x_k^{(m)}\right)$$

end

for $j = n$ **to** 1

$$x_j^{(m+1)} = (1-\omega)x_j^{(m+1/2)} + \frac{\omega}{a_{jj}}\left(b_j - \sum_{k=1}^{j-1} a_{jk}x_k^{(m+1/2)} - \sum_{k=j+1}^{n} a_{jk}x_k^{(m+1)}\right)$$

end

3.5. Convergence (see Demmel's book ANLA, section 6.5.5)

Definition 9

A is an irreducible matrix if there is no permutation matrix such that

$$\mathbf{PAP}^\top = \begin{bmatrix} \mathbf{A}_{11} & \mathbf{A}_{12} \\ \mathbf{0} & \mathbf{A}_{22} \end{bmatrix}.$$

Definition 10

A matrix $\mathbf{A} \in \mathbb{C}^{n\times n}$ is weakly row diagonally dominant if for all i,

$$|a_{ii}| \geq \sum_{j\neq i} |a_{ij}|$$

with strict inequality at least once. A matrix **A** is strictly row diagonally dominant if for all i:

$$|a_{ii}| > \sum_{j\neq i} |a_{ij}|.$$

Theorem 11

If $\mathbf{A}$ *is strictly row diagonally dominant, then both Jacobi's and Gauss–Seidel methods converge. In fact* $\|\mathbf{R}_{\mathrm{GS}}\|_\infty \leq \|\mathbf{R}_{\mathrm{J}}\|_\infty < 1$.

Theorem 12

If $\mathbf{A}$ *is irreducible and weakly row diagonally dominant, then both Jacobi's and Gauss–Seidel methods converge, and* $\rho(\mathbf{R}_{\mathrm{GS}}) < \rho(\mathbf{R}_{\mathrm{J}}) < 1$.

Theorem 13

For any matrix $\mathbf{A}$*, it holds* $\rho(\mathbf{R}_{\mathrm{SOR}(\omega)}) \geq |\omega - 1|$*. Therefore* $0 < \omega < 2$ *is required for the convergence of* $\mathrm{SOR}(\omega)$ *for all starting vectors.*

Theorem 14

If $\mathbf{A}$ *is Hermitian positive definite, then* $\rho(\mathbf{R}_{\mathrm{SOR}(\omega)}) < 1$ *for all* $0 < \omega < 2$*, i.e.,* $\mathrm{SOR}(\omega)$ *converges for all* $0 < \omega < 2$*. Gauss–Seidel* ($\mathrm{SOR}(1)$) *converges for Hermitian positive definite* $\mathbf{A}$.