

# Prey-Predator model by Volterra-Lotka





Introduction History Applications

(slides)

Model equations Model properties

(whiteboard)

**Model simulation** 

(Matlab)

Wrap-up

(slides)

## Introduction

- Ecological model describing predation dynamics within 2 species (prey-predator)
- Developed independently by Alfred J. Lotka (1925) and by Vito Volterra (1926)
- Highly simplified model, but it's the basis for more advanced and sophisticated models



Alfred J. Lotka



Vito Volterra





# **History**

- Volterra's son in law (D'Ancona, a biologist) was looking at fish catched before and during WWI.
- He noticed 2 aspects that he couldn't explain:
  - 1) CYCLICAL BEHAVIOUR
  - 2) MORE SHARKS CATCHED THAN NORMAL FISHES



## Other observed cases








## **Examples of applications to physiology**

#### FLEXIBLE MODELS -> MANY DIFFERENT APPLICATIONS

## 1) Bone formation remodelling


**Osteoblasts** (PREY) promote the growth of **osteoclasts** (PREDATOR).



A. Nutini, Theoretical model of bone remodelling, RJBIO (2015)

## 2) Tumor-Immune System Dynamics


Immune cells (PREDATOR) "hunt" or suppress tumor cells (PREY).



L.G De Pillis, A Radunskaya, The dynamics of an optimally controlled tumor model: A case study, Mat Com Mod (2003)




# Model setup






## Volterra-Lotka model

$$\begin{cases} \frac{dx}{dt} = \alpha x - \beta xy \\ \frac{dy}{dt} = -\gamma y + \delta xy \end{cases}$$



## **Predicts oscillations in populations**






## Some model assumptions

**FOOD** 

#### **UNIFORMITY**

#### **UNLIMITED FOOD AVAILABILITY**

Preys can always find food



#### **EXCLUSIVE PREDATION**

Predators only eats preys



Predators will always eat when they can



#### **SPATIAL**

Preys/predators are distributed homogeneously

#### **TEMPORAL**

Preys and predators don't get old



#### **OTHER**

#### **NO EVOLUTION**

Predators will eat preys with the same efficiency over time




#### NO EXTINCTION


There can be fractional numbers of preys and predators






# Matlab model simulation











Predator population (y)



## Matlab model simulation

#### **GOALS**

- Compute equilibrium point (x\*,y\*)
- 2. Generate time-domain curves for several (x0, y0)
- 3. Generate phases-space orbits for several (x0, y0)
- 4. Compute ecological potential over time
- 8. Find duration of 1 cycle
- 9. Compute average values over 1 cycle
- 10. Generate surface ecological potential  $\phi$  vs x,y
- 11. Draw orbits as levels of  $\phi$

#### **NICE PARAMETERS**

$$x0=20:40:100$$
  $\alpha = 1$   
 $y0=10:10:30$   $\beta = 0.05$   
 $t0=0$   $\gamma = 1$   
 $tf=300$   $\delta = 0.01$ 

#### **SUGGESTIONS**

- 1. Create model as function with input: t, y,  $\alpha$ ,  $\beta$ ,  $\gamma$ ,  $\delta$
- 2. Solve using ODE45

## **MODEL**

$$\begin{cases} \frac{dx}{dt} = \alpha x - \beta xy \\ \frac{dy}{dt} = -\gamma y + \delta xy \end{cases}$$

## **EQUILIBRIUM**

$$x^* = \frac{\gamma}{\delta}, y^* = \frac{\alpha}{\beta}$$

## **ECOLOGICAL POTENTIAL**


$$\phi = -\gamma \ln(x) - \alpha \ln(y) + \delta x + \beta y$$


## Prey-Predator model (Volterra-Lotka)

# Extension to n species

$$\frac{dx_i}{dt} = x_i \sum_{j=1}^n A_{ij} (1 - x_j)$$

$$A = \begin{bmatrix} A_{11} & A_{12} & A_{13} \\ A_{21} & A_{22} & A_{23} \\ A_{31} & A_{32} & A_{33} \end{bmatrix}$$








Figure 12.3 Population levels for the three-species Lotka-Volterra system

riguer oron The Computational Dealay or nature. Computer Expositions of Tractics, Chaop, Compute Systems, and Respection. Copyrigin. (g) 1999 Gary William Flake, All rights reserved. Permission granted for educational, scholarly, and personal use provided that this notice remains intact and unaftered. I part of this work may be reproduced for commercial purposes without prior written permission from the MIT Press.




Figure 12.4 Period doublings in a three-species Lotka-Volterra system: phase space is on the left and  $x_1$  is plotted on the right. (a) spiral fixed point, (b) simple periodic orbit, (c) period-2 orbit, (d) period-4 orbit, (e) chaos

Figure from The Computational Beauty of Nature: Computer Explorations of Fractals, Chaos, Complex Systems, and Adaptation. Copyright © 1998–2000 b Garry William Flake, All rights reserved. Permission granted for educational, scholarly, and personal use provided that this notice remains intact and unaltered. N part of this work may be reproduced for commercial purposes without prior written permission from the MIT Press.