

COL703: Logic for Computer Science (Aug-Nov 2022)

Lectures 9 & 10 (SAT Solving, Binary Decision Diagrams)

Kumar Madhukar

madhukar@cse.iitd.ac.in

September 15th and 19th, 2022

Horn formulas

- a **literal** is a boolean variable or its negation
- for a variable x , we have a **positive literal** (x) and a **negative literal** ($\neg x$)
- a **horn clause** is a finite disjunction of literals with **at most one positive literal**
- a **horn formula** is a finite conjunction of horn clauses
- **example**
$$(x \vee \neg y \vee \neg z \vee \neg w) \wedge (\neg x \vee \neg y \vee \neg w) \wedge (\neg x \vee \neg z \vee w) \wedge (\neg x \vee y) \wedge (x) \wedge (\neg z) \wedge (\neg x \vee \neg y \vee w)$$

Horn-SAT

- if the formula contains a **unit clause**, say (ℓ)
 - all **clauses** containing (ℓ) **is removed**
 - from all clauses containing $(\neg \ell)$ have $(\neg \ell)$ **removed**
- this may generate new unit clauses, which are propagated similarly
- if there are no unit clauses left, the formula can be **satisfied by setting every remaining variable to false**
- formula is **unsat** if propagation generates an **empty clause**

2-SAT

- given a 2-CNF formula, is it satisfiable or not
- every clause has 2 literals
- example
 $(\neg x \vee y) \wedge (\neg y \vee z) \wedge (x \vee \neg z) \wedge (z \vee y)$

2-SAT satisfiability check

- create a graph with $2n$ vertices (for a formula with n variables)
- corresponding to positive and negative literals for every variable
- for every clause $(a \vee b)$, create directed edges $\neg a \rightarrow b$ and $\neg b \rightarrow a$

2-SAT satisfiability check

- create a graph with $2n$ vertices (for a formula with n variables)
- corresponding to positive and negative literals for every variable
- for every clause $(a \vee b)$, create directed edges $\neg a \rightarrow b$ and $\neg b \rightarrow a$
- **claim:** if the graph contains a path from α to β , then it also contains a path from $\neg\beta$ to $\neg\alpha$

2-SAT satisfiability check

- create a graph with $2n$ vertices (for a formula with n variables)
- corresponding to positive and negative literals for every variable
- for every clause $(a \vee b)$, create directed edges $\neg a \rightarrow b$ and $\neg b \rightarrow a$
- **claim:** if the graph contains a path from α to β , then it also contains a path from $\neg\beta$ to $\neg\alpha$
- **claim:** a 2-CNF formula is unsat iff there exists a variable x such that:
 - there is a path from x to $\neg x$
 - there is a path from $\neg x$ to x

2-SAT satisfying assignment

- pick an unassigned literal ℓ , with no path from ℓ to $\neg\ell$
- assign true to ℓ and all vertices reachable from ℓ (and assign false to their negations)
- repeat until all vertices are assigned

Example

$$(\neg x \vee y) \wedge (\neg y \vee z) \wedge (x \vee \neg z) \wedge (z \vee y)$$




image source: <https://www.iitg.ac.in/deepkesh/CS301/assignment-2/2sat.pdf>

Another example

$$(x_1 \vee \neg x_2) \wedge (\neg x_1 \vee \neg x_3) \wedge (x_1 \vee x_2) \wedge (x_4 \vee \neg x_3) \wedge (x_4 \vee \neg x_1)$$

Tseitin transformation

- we know that an arbitrary boolean formula can be converted to CNF
- using De Morgan's law and distributivity property
- but this may result in an exponential explosion of the formula
- example: $(x_1 \wedge y_1) \vee (x_2 \wedge y_2) \vee \dots \vee (x_n \wedge y_n)$
- Tseitin transformation is guaranteed to only linearly increase the size of the formula

Tseitin transformation: Example

(source: Wikipedia)

Consider the following formula ϕ .

$$\phi := ((p \vee q) \wedge r) \rightarrow (\neg s)$$

Consider all subformulas (excluding simple variables):

$$\neg s$$

$$p \vee q$$

$$(p \vee q) \wedge r$$

$$((p \vee q) \wedge r) \rightarrow (\neg s)$$

Introduce a new variable for each subformula:

$$x_1 \leftrightarrow \neg s$$

$$x_2 \leftrightarrow p \vee q$$

$$x_3 \leftrightarrow x_2 \wedge r$$

$$x_4 \leftrightarrow x_3 \rightarrow x_1$$

Conjoin all substitutions and the substitution for ϕ :

$$T(\phi) := x_4 \wedge (x_4 \leftrightarrow x_3 \rightarrow x_1) \wedge (x_3 \leftrightarrow x_2 \wedge r) \wedge (x_2 \leftrightarrow p \vee q) \wedge (x_1 \leftrightarrow \neg s)$$

All substitutions can be transformed into CNF, e.g.

$$\begin{aligned} x_2 \leftrightarrow p \vee q &\equiv (x_2 \rightarrow (p \vee q)) \wedge ((p \vee q) \rightarrow x_2) \\ &\equiv (\neg x_2 \vee p \vee q) \wedge (\neg(p \vee q) \vee x_2) \\ &\equiv (\neg x_2 \vee p \vee q) \wedge ((\neg p \wedge \neg q) \vee x_2) \\ &\equiv (\neg x_2 \vee p \vee q) \wedge (\neg p \vee x_2) \wedge (\neg q \vee x_2) \end{aligned}$$

1-SAT to 3-SAT

$$c = (\ell)$$

$$c' = (\ell \vee u \vee v) \wedge (\ell \vee \neg u \vee v) \wedge (\ell \vee u \vee \neg v) \wedge (\ell \vee \neg u \vee \neg v)$$

c' is satisfiable iff c is satisfiable.

2-SAT to 3-SAT

$$c = (l_1 \vee l_2)$$

$$c' = (l_1 \vee l_2 \vee u) \wedge (l_1 \vee l_2 \vee \neg u)$$

c' is satisfiable iff c is satisfiable.

$k(>3)$ -SAT to $(k-1)$ -SAT

$$c = (\ell_1 \vee \ell_2 \vee \dots \vee \ell_k)$$

$$c' = (\ell_1 \vee \ell_2 \vee \dots \vee \ell_{k-2} \vee u) \wedge (\ell_{k-1} \vee \ell_k \vee \neg u)$$

c' is satisfiable iff c is satisfiable.

breaking 3SAT similarly to get 2SAT fails!

$$c = (l_1 \vee l_2 \vee l_3)$$

$$c' = (l_1 \vee l_2 \vee u) \wedge (l_3 \vee \neg u) \quad (\text{still 3!})$$

$$c' = (l_1 \vee u) \wedge (l_2 \vee l_3 \vee \neg u) \quad (\text{still 3!})$$

Davis-Putnam Algorithm

- **unit-clause** – if there is a unit clause ℓ , delete all clauses containing ℓ , and delete all occurrences of $\neg\ell$ from other clauses
- **pure-literal** – if there is a pure literal ℓ , delete all clauses containing ℓ
- **eliminate a variable by resolution** – choose an atom p and perform all possible resolutions on clauses that clash on p and $\neg p$. Add these resolvents to the set of clauses and then delete all the clauses containing p or $\neg p$.
- use these repeatedly; but use resolution only if the the first two rules do not apply

Davis-Putnam Algorithm

- if **empty clause** is produced, the formula is **unsat**
- if no more rules are applicable, report **sat**
- why does this terminate?
- why is this correct?
- example: $(p) \wedge (\neg p \vee q) \wedge (\neg q \vee r) \wedge (\neg r \vee s \vee t)$

DPLL Algorithm

- creating all possible resolvents is very inefficient
- DPLL improves on the DP algorithm by replacing the variable elimination with a search for a model of the formula

Next class

- DPLL
- Binary Decision Diagrams

Thank you!