

Homework 13

Mengxiang Jiang
CSEN 5303 Foundations of Computer Science

November 14, 2022

Problem 1. Determine which rule of inference is used in each of the following arguments.

1. Kangaroos live in Australia and are marsupials. Therefore, kangaroos are marsupials.

Simplification: $(p \wedge q) \rightarrow p$

2. It is hotter than 100 degrees today or the pollution is dangerous. It is less than 100 degrees outside today. Therefore, the pollution is dangerous.

Disjunctive syllogism: $[(p \vee q) \wedge \neg p] \rightarrow q$

3. Linda is an excellent swimmer. If Linda is an excellent swimmer, then she can work as a lifeguard. Therefore, Linda can work as a lifeguard.

Modus ponens: $[p \wedge (p \rightarrow q)] \rightarrow q$

4. Steve will work at a computer company this summer. Therefore, this summer Steve will work at a computer company or he will be in beach bun.

Addition: $p \rightarrow (p \vee q)$

5. If I work all night on this homework, then I can answer all the exercises. If I answer all the exercises, I will understand the material. Therefore, if I work all night on this homework, then I will understand the material.

Hypothetical syllogism: $[(p \rightarrow q) \wedge (q \rightarrow r)] \rightarrow (p \rightarrow r)$

6. If George does not have eight legs, then he is not an insect. George is an insect. Therefore, George has eight legs.

Modus tollens: $[\neg q \wedge (p \rightarrow q)] \rightarrow \neg p$

Problem 2. Let $P(n)$ be the statement that $1^2 + 2^2 + \dots + n^2 = n(n+1)(2n+1)/6$ for the positive integer n .

1. What is the statement $P(1)$?

$$1^2 = 1(1+1)(2+1)/6$$

2. Show that $P(1)$ is true, completing the basis step of the proof.

$$1^2 = 1(1) = 1$$

$$1(1+1)(2+1)/6 = 1(2)(3)/6 = 6/6 = 1$$

$\therefore P(1)$ is true.

3. What is the inductive hypothesis?

Assume $P(k)$ is true for some integer $k \geq 1$.

4. What do you need to prove in the inductive step?

Prove $P(k+1)$ is true using $P(k)$.

5. Complete the inductive step.

$$P(k+1): 1^2 + 2^2 + \dots + k^2 + (k+1)^2 = (1^2 + 2^2 + \dots + k^2) + (k+1)^2$$

$$= \frac{k(k+1)(2k+1)}{6} + (k+1)^2 = (k+1) \left(\frac{2k^2 + k}{6} + k + 1 \right)$$

$$= (k+1) \left(\frac{2k^2 + 7k + 6}{6} \right) = (k+1) \left(\frac{(k+2)(2k+3)}{6} \right)$$

$$= \frac{(k+1)((k+1)+1)(2(k+1)+1)}{6}$$

6. Explain why these steps show that this formula is true whenever n is a positive integer.

Since we have shown $P(1)$ is true in step 2, we can let $k = 1$ in step 3 and show $P(2)$ is true in step 5. We can then set $k = 2$ in step 3 and show $P(3)$ is true in step 5, and so on for all the integers.

Problem 6. Prove each of the following statements.

1. 2 divides $n^2 + n$ whenever n is a positive integer.

Two cases for n :

Case 1: $n \equiv 0 \pmod{2} \rightarrow n^2 + n \equiv 0^2 + 0 \equiv 0 \pmod{2} \rightarrow 2|(n^2 + n)$

Case 2: $n \equiv 1 \pmod{2} \rightarrow n^2 + n \equiv 1^2 + 1 \equiv 2 \equiv 0 \pmod{2} \rightarrow 2|(n^2 + n)$

2. 3 divides $n^3 + 2n$ whenever n is a positive integer.

Three cases for n :

Case 1: $n \equiv 0 \pmod{3} \rightarrow n^3 + 2n \equiv 0^3 + 2(0) \equiv 0 \pmod{3} \rightarrow 3|(n^3 + 2n)$

Case 2: $n \equiv 1 \pmod{3} \rightarrow n^3 + 2n \equiv 1^3 + 2(1) \equiv 3 \equiv 0 \pmod{3} \rightarrow 3|(n^3 + 2n)$

Case 3: $n \equiv 2 \pmod{3} \rightarrow n^3 + 2n \equiv 2^3 + 2(2) \equiv 12 \equiv 0 \pmod{3} \rightarrow 3|(n^3 + 2n)$

3. 5 divides $n^5 - n$ whenever n is a nonnegative integer.

Five cases for n :

Case 1: $n \equiv 0 \pmod{5} \rightarrow n^5 - n \equiv 0^5 - 0 \equiv 0 \pmod{5} \rightarrow 5|(n^5 - n)$

Case 2: $n \equiv 1 \pmod{5} \rightarrow n^5 - n \equiv 1^5 - 1 \equiv 0 \pmod{5} \rightarrow 5|(n^5 - n)$

Case 3: $n \equiv 2 \pmod{5} \rightarrow n^5 - n \equiv 2^5 - 2 \equiv 30 \equiv 0 \pmod{5} \rightarrow 5|(n^5 - n)$

Case 4: $n \equiv 3 \pmod{5} \rightarrow n^5 - n \equiv n(n^4 - 1) \equiv n(n^2 - 1)(n^2 + 1) \equiv 3(3^2 - 1)(3^2 + 1) \equiv 3(8)(10) \equiv 3(8)(0) \equiv 0 \pmod{5} \rightarrow 5|(n^5 - n)$

Case 5: $n \equiv 4 \pmod{5} \rightarrow n^5 - n \equiv n(n^4 - 1) \equiv n(n^2 - 1)(n^2 + 1) \equiv n(n + 1)(n - 1)(n^2 + 1) \equiv 4(0)(3)(4^2 + 1) \equiv 0 \pmod{5} \rightarrow 5|(n^5 - n)$

4. 6 divides $n^3 - n$ whenever n is a nonnegative integer.

$n^3 - n = n(n^2 - 1) = n(n + 1)(n - 1) = (n - 1)(n)(n + 1)$

$\therefore n^3 - n$ is the product of 3 consecutive integers.

Given 3 consecutive integers, at least one must be a multiple of 3, and one (could be the same one) a multiple of 2 (pigeonhole principle). The product of the 3 integers therefore contains both 2 and 3 as factors, therefore the product must be divisible by 6.