

Quantum phases from semi-classical Monte Carlo

Exploring the phases of $J_1 - J_2$ Heisenberg model

Kunal Verma (MS18148)

Supervisor: Prof. Sanjeev Kumar

PRJ502 Thesis Presentation

April 27, 2023

$J_1 - J_2$ Heisenberg model

Quantum spin model with nearest neighbor and next nearest neighbor coupling

$$\hat{H} = J_1 \sum_{\langle i,j \rangle} \hat{\vec{S}}_i \cdot \hat{\vec{S}}_j + J_2 \sum_{\langle\langle i,j \rangle\rangle} \hat{\vec{S}}_i \cdot \hat{\vec{S}}_j$$

$J_1 - J_2$ Heisenberg model

Quantum spin model with nearest neighbor and next nearest neighbor coupling

$$\hat{H} = J_1 \sum_{\langle i,j \rangle} \hat{\mathbf{S}}_i \cdot \hat{\mathbf{S}}_j + J_2 \sum_{\langle\langle i,j \rangle\rangle} \hat{\mathbf{S}}_i \cdot \hat{\mathbf{S}}_j$$





Figure 1: Lattice structure

Quantum phases at $T = 0$


- a Néel-ordered phase ((π, π) AFM) for $J_2/J_1 \lesssim 0.4$.
- a quantum paramagnetic phase ($0.4 \lesssim J_2/J_1 \lesssim 0.6$).
- a collinear-ordered phase ($(\pi, 0)$ or $(0, \pi)$ AFM) for $J_2/J_1 \gtrsim 0.6$.

Quantum phases at $T = 0$

- a Néel-ordered phase ((π, π) AFM) for $J_2/J_1 \lesssim 0.4$.
- a quantum paramagnetic phase ($0.4 \lesssim J_2/J_1 \lesssim 0.6$).
- a collinear-ordered phase ($(\pi, 0)$ or $(0, \pi)$ AFM) for $J_2/J_1 \gtrsim 0.6$.




(a) Néel (π, π) AFM




(b) Collinear $(\pi, 0)$ AFM

Quantum phases at $T = 0$

- a Néel-ordered phase ((π, π) AFM) for $J_2/J_1 \lesssim 0.4$.
- a **quantum paramagnetic** phase ($0.4 \lesssim J_2/J_1 \lesssim 0.6$).
- a collinear-ordered phase ($(\pi, 0)$ or $(0, \pi)$ AFM) for $J_2/J_1 \gtrsim 0.6$.




(a) Néel (π, π) AFM




(b) Collinear $(\pi, 0)$ AFM

Quantum phases at $T = 0$

- a Néel-ordered phase ((π, π) AFM) for $J_2/J_1 \lesssim 0.4$.
- a **spin liquid** phase ($0.4 \lesssim J_2/J_1 \lesssim 0.6$).
- a collinear-ordered phase ($(\pi, 0)$ or $(0, \pi)$ AFM) for $J_2/J_1 \gtrsim 0.6$.



(a) Néel (π, π) AFM



(b) Collinear $(\pi, 0)$ AFM

Spin liquid phase

- Spin liquid \sim a sea of singlet-triplet dimers.
- A purely “quantum” effect.




Figure 3: Spin liquid phase.

Spin liquid phase

- Spin liquid \sim a sea of singlet-triplet dimers.
- A purely “quantum” effect.




Figure 3: Spin liquid phase.

Question: Emergent spin liquid by adding “quantum fluctuations” on top of a classical model?

Semi-classical mapping

$$\hat{\vec{S}}_i \cdot \hat{\vec{S}}_j = \underbrace{\hat{X}_i \hat{X}_j + \hat{Y}_i \hat{Y}_j}_{\text{fluctuation}} + \hat{Z}_i \hat{Z}_j$$

$\hat{X}_i \hat{X}_j + \hat{Y}_i \hat{Y}_j \sim$ acts like a fluctuation in the Z basis.

Semi-classical mapping

$$\vec{\hat{S}}_i \cdot \vec{\hat{S}}_j = \underbrace{\hat{X}_i \hat{X}_j + \hat{Y}_i \hat{Y}_j}_{\text{fluctuations}} + \hat{Z}_i \hat{Z}_j$$

$\hat{X}_i \hat{X}_j + \hat{Y}_i \hat{Y}_j \sim$ acts like a fluctuation in the Z basis.

\Rightarrow **Proposal:** $\vec{\hat{S}}_i \cdot \vec{\hat{S}}_j \approx \underbrace{\hat{Z}_i \hat{Z}_j}_{\text{Ising}} + \text{quantum fluctuations}$

Semi-classical mapping

$$\hat{\vec{S}}_i \cdot \hat{\vec{S}}_j = \underbrace{\hat{X}_i \hat{X}_j + \hat{Y}_i \hat{Y}_j}_{\text{fluctuation}} + \hat{Z}_i \hat{Z}_j$$

$\hat{X}_i \hat{X}_j + \hat{Y}_i \hat{Y}_j \sim$ acts like a fluctuation in the Z basis.

\Rightarrow **Proposal:** $\hat{\vec{S}}_i \cdot \hat{\vec{S}}_j \approx \underbrace{\hat{Z}_i \hat{Z}_j}_{\text{Ising}} + \text{quantum fluctuations}$

\therefore Semi-classical mapping of $J_1 - J_2$ Heisenberg model to a $J_1 - J_2$ classical Ising and add quantum fluctuations on top of it!

$$H = J_1 \sum_{\langle i,j \rangle} \hat{Z}_i \hat{Z}_j + J_2 \sum_{\langle\langle i,j \rangle\rangle} \hat{Z}_i \hat{Z}_j + \text{q.f.}$$

Quantum fluctuations

Source of quantum fluctuations: quantum dimer formation.

Quantum fluctuations

Source of quantum fluctuations: quantum dimer formation.

For a two-site Heisenberg interaction $J\hat{S}_1 \cdot \hat{S}_2$, we get the following singlet-triplet basis

$$\left. \begin{aligned} |s=1; m_s=+1\rangle &= |\uparrow\uparrow\rangle \\ |s=1; m_s=-1\rangle &= |\downarrow\downarrow\rangle \\ |s=1; m_s=0\rangle &= (|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)/\sqrt{2} \end{aligned} \right\} s=1 \text{ (triplet)}$$
$$|s=0; m_s=0\rangle = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)/\sqrt{2} \left. \vphantom{\begin{aligned} |s=1; m_s=+1\rangle &= |\uparrow\uparrow\rangle \\ |s=1; m_s=-1\rangle &= |\downarrow\downarrow\rangle \\ |s=1; m_s=0\rangle &= (|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)/\sqrt{2} \end{aligned}} \right\} s=0 \text{ (singlet)}$$

Quantum fluctuations

Source of quantum fluctuations: quantum dimer formation.

For a two-site Heisenberg interaction $J\hat{S}_1 \cdot \hat{S}_2$, we get the following singlet-triplet basis

$$\left. \begin{aligned} |s=1; m_s=+1\rangle &= |\uparrow\uparrow\rangle \\ |s=1; m_s=-1\rangle &= |\downarrow\downarrow\rangle \\ |s=1; m_s=0\rangle &= (|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)/\sqrt{2} \end{aligned} \right\} s=1 \text{ (triplet)}$$
$$|s=0; m_s=0\rangle = (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)/\sqrt{2} \left. \vphantom{\begin{aligned} |s=1; m_s=+1\rangle &= |\uparrow\uparrow\rangle \\ |s=1; m_s=-1\rangle &= |\downarrow\downarrow\rangle \\ |s=1; m_s=0\rangle &= (|\uparrow\downarrow\rangle + |\downarrow\uparrow\rangle)/\sqrt{2} \end{aligned}} \right\} s=0 \text{ (singlet)}$$

Energy eigenvalues

$$\hat{h} |s=1\rangle = \frac{+J}{4} |s=1\rangle \rightarrow \text{triplet}$$
$$\hat{h} |s=0\rangle = \frac{-3J}{4} |s=0\rangle \rightarrow \text{singlet}$$

Quantum fluctuations (contd.)

Energy of the dimer $\langle E \rangle = \frac{3J}{4} \left(\frac{e^{-\beta J} - 1}{3e^{-\beta J} + 1} \right)$

Magnetization of the dimer $\langle \vec{M} \rangle = 0.$

Quantum fluctuations (contd.)

Energy of the dimer $\langle E \rangle = \frac{3J}{4} \left(\frac{e^{-\beta J} - 1}{3e^{-\beta J} + 1} \right)$

Magnetization of the dimer $\langle \vec{M} \rangle = 0.$

Encode the *quantum effects* of the dimer as a *semi-classical dimer*.

Semi-classical dimer $\langle i, j \rangle$

$$M_{\text{dimer}}(T) = 0, \quad E_{\text{dimer}}(T) = \frac{3J}{4} \left(\frac{e^{-J/T} - 1}{3e^{-J/T} + 1} \right).$$



Two different degrees of freedom/units in the lattice:

- Classical **Ising spin** Hamiltonian:

$$E = J_1 \sum_{\langle i,j \rangle} S_i S_j + J_2 \sum_{\langle\langle i,j \rangle\rangle} S_i S_j$$

- Semi-classical **dimer** $\langle i,j \rangle$ (source of quantum fluctuations):

$$S_i = S_j = 0, \quad E_J(T) = \frac{3J}{4} \left(\frac{e^{-J/T} - 1}{3e^{-J/T} + 1} \right)$$

Two different degrees of freedom/units in the lattice:

- Classical **Ising spin** Hamiltonian:

$$E = J_1 \sum_{\langle i,j \rangle} S_i S_j + J_2 \sum_{\langle\langle i,j \rangle\rangle} S_i S_j$$

- Semi-classical **dimer** $\langle i,j \rangle$ (source of quantum fluctuations):

$$S_i = S_j = 0, \quad E_J(T) = \frac{3J}{4} \left(\frac{e^{-J/T} - 1}{3e^{-J/T} + 1} \right)$$

Combine the dynamics using a Markov chain Monte Carlo simulation!

Semi-classical Monte Carlo

Markov chain process now includes two different types of steps-

- random spin flips moves $(1 - p)$.
- dimer creation/annihilation moves (p) .

Standard Metropolis acceptance $\rightarrow P_{\text{accept}} = \min[1, e^{-\beta\Delta E}]$




Figure 5: Dimer creation on $\langle i, j \rangle$ with $S_i = S_j = 0$ and $E = E_J(T)$.

Order parameters

Three order parameters for four different phases:

- Staggered magnetization (π, π)

$$m_{(\pi, \pi)} = \frac{1}{L^2} \sum_{i=1}^{L^2} (-1)^{i_x + i_y} S_i$$

- Staggered magnetization $(\pi, 0)$ and $(0, \pi)$

$$m_{(\pi, 0)} + m_{(0, \pi)} = \frac{1}{L^2} \sum_{i=1}^{L^2} [(-1)^{i_x} + (-1)^{i_y}] S_i$$



- Number of dimers

$$N_{\text{dimers}} = N_{J_1} \text{ dimers} + N_{J_2} \text{ dimers}$$

Preliminary analysis of phases

Lattice geometry: 10×10

Phases:



Quantum phase transition




Figure 7: $SL \rightarrow (\pi, 0) + (0, \pi)$ AFM transition is first-order, whereas the (π, π) AFM \rightarrow SL transition is second-order.

Results ($p = 0.15$)




Figure 8: Order parameters at $p = 0.15$

Results ($p = 0.25$)




Figure 9: Order parameters at $p = 0.25$

Results ($p = 0.50$)




Figure 10: Order parameters at $p = 0.50$

Entropic stabilization




Figure 11: $T = 0$ phase structure argued from an energetic viewpoint.

Entropic stabilization




Figure 11: $T = 0$ phase structure argued from an energetic viewpoint.

- Broader SL region, but meta-stable at very small temperatures!
- AFM state entropically more favorable than a dimerized SL (even at extremely small temperatures).

Entropic stabilization (contd.)




Figure 12: Entropic stabilization of AFM and destabilization of the spin liquid state.




- Entropic stabilization pushes the boundaries of the phases,
- can be tuned via the parameter p to tune the phase transition points.

Conclusions:

- $J_1 - J_2$ Heisenberg $\xrightarrow{\text{semi-classical}}$ $J_1 - J_2$ Ising model + dimer fluctuations.
- Quantum phases do emerge as a semi-classical correction!

Future direction:

- Tune p to maximises the Monte Carlo acceptance ratios.
- Address critical properties of the model.
- Dimer structure factor as a direct indicator for the SL phase.
- Extension of the “semi-classical” method to other lattice geometries and spin models.

-  Janke, W. (2008). *Monte Carlo Methods in Classical Statistical Physics*. In: Fehske, H., Schneider, R., Weiße, A. (eds) *Computational Many-Particle Physics*. Lecture Notes in Physics, vol 739. Springer, Berlin, Heidelberg.
https://doi.org/10.1007/978-3-540-74686-7_4.
-  Yong-Zhi Ren, Ning-Hua Tong, and Xin-Cheng Xie. *Cluster mean-field theory study of $J_1 - J_2$ Heisenberg model on a square lattice*. In: *Journal of Physics: Condensed Matter* 26.11 (Mar. 2014), p. 115601. doi:
[10.1088/0953-8984/26/11/115601](https://dx.doi.org/10.1088/0953-8984/26/11/115601). url:
<https://dx.doi.org/10.1088/0953-8984/26/11/115601>.
-  A. W. Sandvik, A. Avella, F. Mancini. *Computational Studies of Quantum Spin Systems*. AIP Conference Proceedings. AIP, 2010. doi: [10.1063/1.3518900](https://doi.org/10.1063/1.3518900).