

Nonlinear Dynamical Systems:

Lecture 10

Madhu N. Belur & Harish K. Pillai
Control & Computing group
Department of Electrical Engineering
Indian Institute of Technology Bombay

(IIT Bombay)

Outline

- **Bendixson criterion and Poincaré-Bendixson criterion**
- **Examples: Lotka-Volterra predator prey model**
- **van der Pol oscillator**

Lotka Volterra Predator Prey Model

Predator \rightarrow Hunter

Population dynamics of two species: prey and hunter.

Equations of the model

Lotka Volterra Predator Prey Model

Predator \rightarrow Hunter

Population dynamics of two species: prey and hunter.

Equations of the model

$$\begin{aligned}\dot{x}_h &= -x_h + x_h x_p \\ \dot{x}_p &= x_p - x_h x_p\end{aligned}$$

x_h is the amount of hunter specimen in the model

Lotka Volterra Predator Prey Model

Predator \rightarrow Hunter

Population dynamics of two species: prey and hunter.

Equations of the model

$$\begin{aligned}\dot{x}_h &= -x_h + x_h x_p \\ \dot{x}_p &= x_p - x_h x_p\end{aligned}$$

x_h is the amount of hunter specimen in the model

x_p is the amount of prey specimen in the model

Lotka Volterra Predator Prey Model

Predator \rightarrow Hunter

Population dynamics of two species: prey and hunter.

Equations of the model

$$\begin{aligned}\dot{x}_h &= -x_h + x_h x_p \\ \dot{x}_p &= x_p - x_h x_p\end{aligned}$$

x_h is the amount of hunter specimen in the model

x_p is the amount of prey specimen in the model

More generally,

$$\begin{aligned}\dot{x}_h &= -ax_h + bx_h x_p \\ \dot{x}_p &= cx_p - dx_h x_p\end{aligned}$$

(parameters a , b , c and d are positive.)

Equilibrium points ?

Equilibrium points ?

Nature of equilibrium points for the linearized system ?

Equilibrium points ?

Nature of equilibrium points for the linearized system ?

Periodic orbits?

Equilibrium points ?

Nature of equilibrium points for the linearized system ?

Periodic orbits?

Bendixson criterion and Poincaré Bendixson criterion?

Another example

Consider the following:

$$\begin{aligned}\dot{x}_1 &= x_2 + (x_1 x_2^2) \\ \dot{x}_2 &= -x_1 + (x_2 x_1^2)\end{aligned}$$

$$\nabla \cdot f(x) = \frac{\partial f_1}{\partial x_1} + \frac{\partial f_2}{\partial x_2} = (x_2^2 + x_1^2)$$

$\nabla \cdot f(x)$ is always positive,

Another example

Consider the following:

$$\begin{aligned}\dot{x}_1 &= x_2 + (x_1 x_2^2) \\ \dot{x}_2 &= -x_1 + (x_2 x_1^2)\end{aligned}$$

$$\nabla \cdot f(x) = \frac{\partial f_1}{\partial x_1} + \frac{\partial f_2}{\partial x_2} = (x_2^2 + x_1^2)$$

$\nabla \cdot f(x)$ is always positive, except at equilibrium point.
Hence, by Bendixson criteria, there are no periodic orbits.

Another example

Consider

$$\dot{x}_1 = x_2 + x_1 x_2^2$$

$$\dot{x}_2 = -x_1 - x_2 x_1^2$$

Another example

Consider

$$\dot{x}_1 = x_2 + x_1 x_2^2$$

$$\dot{x}_2 = -x_1 - x_2 x_1^2$$

$$\nabla \cdot f(x) = \frac{\partial f_1}{\partial x_1} + \frac{\partial f_2}{\partial x_2} = (x_2^2 - x_1^2)$$

$\nabla \cdot f(x)$ is zero for $x_1 = x_2$, and changes sign.

Another example

Consider

$$\dot{x}_1 = x_2 + x_1 x_2^2$$

$$\dot{x}_2 = -x_1 - x_2 x_1^2$$

$$\nabla \cdot f(x) = \frac{\partial f_1}{\partial x_1} + \frac{\partial f_2}{\partial x_2} = (x_2^2 - x_1^2)$$

$\nabla \cdot f(x)$ is zero for $x_1 = x_2$, and changes sign.
Bendixson criterion ?

Periodic orbits

Linear case:

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & a \\ -2 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \mathbf{A} \mathbf{x}$$

Periodic orbits

Linear case:

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & a \\ -2 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \mathbf{A} \mathbf{x}$$

$$a = 5,$$

Periodic orbits

Linear case:

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & a \\ -2 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \mathbf{A} \mathbf{x}$$

$a = 5,$ $a = -3$?

Periodic orbits

Linear case:

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & a \\ -2 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \mathbf{A} \mathbf{x}$$

$a = 5$, $a = -3$?

$$\mathbf{A} = \begin{bmatrix} \epsilon & 2 \\ -2 & \epsilon \end{bmatrix}$$

for $\epsilon > 0$?

Periodic orbits

Linear case:

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & a \\ -2 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \mathbf{A} \mathbf{x}$$

$a = 5$, $a = -3$?

$$\mathbf{A} = \begin{bmatrix} \epsilon & 2 \\ -2 & \epsilon \end{bmatrix}$$

for $\epsilon > 0$? for $\epsilon < 0$

What if ϵ 'changes' sign ?

Consider

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} (25 - x_1^2 - x_2^2) & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

This can be written in matrix form as

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} \epsilon(r) & 1 \\ -1 & \epsilon(r) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Consider the cases

What if ϵ 'changes' sign ?

Consider

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} (25 - x_1^2 - x_2^2) & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

This can be written in matrix form as

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} \epsilon(r) & 1 \\ -1 & \epsilon(r) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Consider the cases

1. When $r = 5$, we have $\epsilon(r) = 0$.

What if ϵ 'changes' sign ?

Consider

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} (25 - x_1^2 - x_2^2) & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

This can be written in matrix form as

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} \epsilon(r) & 1 \\ -1 & \epsilon(r) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Consider the cases

1. When $r = 5$, we have $\epsilon(r) = 0$.
2. When $r < 5$, we have $\epsilon(r) > 0$.

What if ϵ 'changes' sign ?

Consider

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} (25 - x_1^2 - x_2^2) & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

This can be written in matrix form as

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} \epsilon(r) & 1 \\ -1 & \epsilon(r) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Consider the cases

1. When $r = 5$, we have $\epsilon(r) = 0$.
2. When $r < 5$, we have $\epsilon(r) > 0$.
3. When $r > 5$, we have $\epsilon(r) < 0$.

What if ϵ 'changes' sign ?

Consider

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} (25 - x_1^2 - x_2^2) & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

This can be written in matrix form as

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} \epsilon(r) & 1 \\ -1 & \epsilon(r) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Consider the cases

1. When $r = 5$, we have $\epsilon(r) = 0$.
2. When $r < 5$, we have $\epsilon(r) > 0$.
3. When $r > 5$, we have $\epsilon(r) < 0$.

Hence, the trajectories will approach the **periodic orbit**.

Example: (about Poincare Bendixson criterion)

Consider the case $\dot{x} = Ax$ where $A = \begin{bmatrix} 0 & 2 \\ -2 & \epsilon \end{bmatrix}$
for the cases $\epsilon > 0$,

Example: (about Poincare Bendixson criterion)

Consider the case $\dot{x} = Ax$ where $A = \begin{bmatrix} 0 & 2 \\ -2 & \epsilon \end{bmatrix}$
for the cases $\epsilon > 0$, $\epsilon = 0$,

Example: (about Poincare Bendixson criterion)

Consider the case $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x}$ where $\mathbf{A} = \begin{bmatrix} 0 & 2 \\ -2 & \epsilon \end{bmatrix}$
for the cases $\epsilon > 0$, $\epsilon = 0$, $\epsilon < 0$,

Example: (about Poincare Bendixson criterion)

Consider the case $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x}$ where $\mathbf{A} = \begin{bmatrix} 0 & 2 \\ -2 & \epsilon \end{bmatrix}$
for the cases $\epsilon > 0$, $\epsilon = 0$, $\epsilon < 0$,

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Converting to polar coordinates, we get

Example: (about Poincare Bendixson criterion)

Consider the case $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x}$ where $\mathbf{A} = \begin{bmatrix} 0 & 2 \\ -2 & \epsilon \end{bmatrix}$
for the cases $\epsilon > 0$, $\epsilon = 0$, $\epsilon < 0$,

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Converting to polar coordinates, we get

$$x_1 = r \cos \theta$$

$$x_2 = r \sin \theta$$

Example (continued)

$$x_1^2 + x_2^2 = r^2$$

Example (continued)

$$x_1^2 + x_2^2 = r^2$$

$$x_1 \dot{x}_1 + x_2 \dot{x}_2 = r \dot{r}$$

Example (continued)

$$x_1^2 + x_2^2 = r^2$$

$$x_1 \dot{x}_1 + x_2 \dot{x}_2 = r \dot{r}$$

Substituting $x_1, x_2, \dot{x}_1, \dot{x}_2$ using r, \dot{r} and θ , we get $r \dot{r} =$

Example (continued)

$$x_1^2 + x_2^2 = r^2$$

$$x_1 \dot{x}_1 + x_2 \dot{x}_2 = r \dot{r}$$

Substituting $x_1, x_2, \dot{x}_1, \dot{x}_2$ using r, \dot{r} and θ , we get $r \dot{r} =$

$$(r \cos \theta)(r \sin \theta) + (r \sin \theta)(-r \cos \theta + (25 - r^2)r \sin \theta)$$

This reduces to $\dot{r} = (25 - r^2)r \sin^2 \theta$

Therefore at $r = 5$ we get $\dot{r} = 0$ i.e circle with radius 5 is a periodic orbit.

Moreover, other trajectories 'converge' to this limit cycle:

Example (continued)

$$x_1^2 + x_2^2 = r^2$$

$$x_1 \dot{x}_1 + x_2 \dot{x}_2 = r \dot{r}$$

Substituting $x_1, x_2, \dot{x}_1, \dot{x}_2$ using r, \dot{r} and θ , we get $r \dot{r} =$

$$(r \cos \theta)(r \sin \theta) + (r \sin \theta)(-r \cos \theta + (25 - r^2)r \sin \theta)$$

This reduces to $\dot{r} = (25 - r^2)r \sin^2 \theta$

Therefore at $r = 5$ we get $\dot{r} = 0$ i.e circle with radius 5 is a periodic orbit.

Moreover, other trajectories 'converge' to this limit cycle: 'isolated stable limit cycle'.

van der Pol oscillator

This oscillator has a 'stable limit cycle'.
Differential equation

$$\ddot{x} - \epsilon(1 - x^2)\dot{x} + x = 0$$

where $\epsilon > 0$

Let $\dot{x} = y$ and $\dot{y} = -x + \epsilon(1 - x^2)y$