

# **Nonlinear Dynamical Systems:**

## **Lecture 12**

**Madhu N. Belur & Harish K. Pillai**  
**Control & Computing group**  
**Department of Electrical Engineering**  
**Indian Institute of Technology Bombay**  
  
**(IIT Bombay)**

# Outline

## Scilab simulation of

- Lotka-Volterra predator prey model
- van der Pol oscillator

# Lotka Volterra Predator Prey Model

**Predator  $\rightarrow$  Hunter**

**Population dynamics of two species: prey and hunter.**

**Equations of the model**

# Lotka Volterra Predator Prey Model

**Predator  $\rightarrow$  Hunter**

**Population dynamics of two species: prey and hunter.**

**Equations of the model**

$$\begin{aligned}\dot{x}_h &= -x_h + x_h x_p \\ \dot{x}_p &= x_p - x_h x_p\end{aligned}$$

$x_h$  is the amount of hunter specimen in the model

# Lotka Volterra Predator Prey Model

**Predator  $\rightarrow$  Hunter**

**Population dynamics of two species: prey and hunter.**

**Equations of the model**

$$\begin{aligned}\dot{x}_h &= -x_h + x_h x_p \\ \dot{x}_p &= x_p - x_h x_p\end{aligned}$$

$x_h$  is the amount of hunter specimen in the model

$x_p$  is the amount of prey specimen in the model

# Lotka Volterra Predator Prey Model

Predator  $\rightarrow$  Hunter

Population dynamics of two species: prey and hunter.

Equations of the model

$$\begin{aligned}\dot{x}_h &= -x_h + x_h x_p \\ \dot{x}_p &= x_p - x_h x_p\end{aligned}$$

$x_h$  is the amount of hunter specimen in the model

$x_p$  is the amount of prey specimen in the model

More generally,

$$\begin{aligned}\dot{x}_h &= -ax_h + bx_h x_p \\ \dot{x}_p &= cx_p - dx_h x_p\end{aligned}$$

(parameters  $a$ ,  $b$ ,  $c$  and  $d$  are positive.)

## Equilibrium points ?

**Equilibrium points ?**

**Nature of equilibrium points for the linearized system ?**



**Equilibrium points ?**

**Nature of equilibrium points for the linearized system ?**

**Periodic orbits?**

**Equilibrium points ?**

**Nature of equilibrium points for the linearized system ?**

**Periodic orbits?**

**Bendixson criterion and Poincaré Bendixson criterion?**

## Another example

Consider the following:

$$\begin{aligned}\dot{x}_1 &= x_2 + (x_1 x_2^2) \\ \dot{x}_2 &= -x_1 + (x_2 x_1^2)\end{aligned}$$

$$\nabla \cdot f(x) = \frac{\partial f_1}{\partial x_1} + \frac{\partial f_2}{\partial x_2} = (x_2^2 + x_1^2)$$

$\nabla \cdot f(x)$  is always positive,

## Another example

Consider the following:

$$\begin{aligned}\dot{x}_1 &= x_2 + (x_1 x_2^2) \\ \dot{x}_2 &= -x_1 + (x_2 x_1^2)\end{aligned}$$

$$\nabla \cdot f(x) = \frac{\partial f_1}{\partial x_1} + \frac{\partial f_2}{\partial x_2} = (x_2^2 + x_1^2)$$

$\nabla \cdot f(x)$  is always positive, except at equilibrium point.  
Hence, by Bendixson criteria, there are no periodic orbits.

## Another example

Consider

$$\dot{x}_1 = x_2 + x_1 x_2^2$$

$$\dot{x}_2 = -x_1 - x_2 x_1^2$$

## Another example

Consider

$$\dot{x}_1 = x_2 + x_1 x_2^2$$

$$\dot{x}_2 = -x_1 - x_2 x_1^2$$

$$\nabla \cdot f(x) = \frac{\partial f_1}{\partial x_1} + \frac{\partial f_2}{\partial x_2} = (x_2^2 - x_1^2)$$

$\nabla \cdot f(x)$  is zero for  $x_1 = x_2$ , and changes sign.

## Another example

Consider

$$\dot{x}_1 = x_2 + x_1 x_2^2$$

$$\dot{x}_2 = -x_1 - x_2 x_1^2$$

$$\nabla \cdot f(x) = \frac{\partial f_1}{\partial x_1} + \frac{\partial f_2}{\partial x_2} = (x_2^2 - x_1^2)$$

$\nabla \cdot f(x)$  is zero for  $x_1 = x_2$ , and changes sign.  
Bendixson criterion ?

# Periodic orbits

**Linear case:**

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & a \\ -2 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \mathbf{A} \mathbf{x}$$



# Periodic orbits

**Linear case:**

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & a \\ -2 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \mathbf{A} \mathbf{x}$$

$$a = 5,$$

# Periodic orbits

**Linear case:**

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & a \\ -2 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \mathbf{A} \mathbf{x}$$

**$a = 5$ ,       $a = -3$  ?**

# Periodic orbits

**Linear case:**

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & a \\ -2 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \mathbf{A} \mathbf{x}$$

**$a = 5$ ,  $a = -3$  ?**

$$\mathbf{A} = \begin{bmatrix} \epsilon & 2 \\ -2 & \epsilon \end{bmatrix}$$

**for  $\epsilon > 0$  ?**

# Periodic orbits

**Linear case:**

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & a \\ -2 & 0 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \mathbf{A} \mathbf{x}$$

**$a = 5$ ,      $a = -3$  ?**

$$\mathbf{A} = \begin{bmatrix} \epsilon & 2 \\ -2 & \epsilon \end{bmatrix}$$

**for  $\epsilon > 0$  ? for  $\epsilon < 0$**

## What if $\epsilon$ 'changes' sign ?

Consider

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} (25 - x_1^2 - x_2^2) & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

This can be written in matrix form as

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} \epsilon(r) & 1 \\ -1 & \epsilon(r) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Consider the cases

## What if $\epsilon$ 'changes' sign ?

Consider

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} (25 - x_1^2 - x_2^2) & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

This can be written in matrix form as

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} \epsilon(r) & 1 \\ -1 & \epsilon(r) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Consider the cases

1. When  $r = 5$ , we have  $\epsilon(r) = 0$ .

## What if $\epsilon$ 'changes' sign ?

Consider

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} (25 - x_1^2 - x_2^2) & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

This can be written in matrix form as

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} \epsilon(r) & 1 \\ -1 & \epsilon(r) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Consider the cases

1. When  $r = 5$ , we have  $\epsilon(r) = 0$ .
2. When  $r < 5$ , we have  $\epsilon(r) > 0$ .

## What if $\epsilon$ 'changes' sign ?

Consider

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} (25 - x_1^2 - x_2^2) & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

This can be written in matrix form as

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} \epsilon(r) & 1 \\ -1 & \epsilon(r) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Consider the cases

1. When  $r = 5$ , we have  $\epsilon(r) = 0$ .
2. When  $r < 5$ , we have  $\epsilon(r) > 0$ .
3. When  $r > 5$ , we have  $\epsilon(r) < 0$ .



## What if $\epsilon$ 'changes' sign ?

Consider

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} (25 - x_1^2 - x_2^2) & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

This can be written in matrix form as

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} \epsilon(r) & 1 \\ -1 & \epsilon(r) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Consider the cases

1. When  $r = 5$ , we have  $\epsilon(r) = 0$ .
2. When  $r < 5$ , we have  $\epsilon(r) > 0$ .
3. When  $r > 5$ , we have  $\epsilon(r) < 0$ .

Hence, the trajectories will approach the **periodic orbit**.

## Example: (about Poincare Bendixson criterion)

Consider the case  $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x}$  where  $\mathbf{A} = \begin{bmatrix} 0 & 2 \\ -2 & \epsilon \end{bmatrix}$   
for the cases  $\epsilon > 0$ ,

## Example: (about Poincare Bendixson criterion)

Consider the case  $\dot{x} = Ax$  where  $A = \begin{bmatrix} 0 & 2 \\ -2 & \epsilon \end{bmatrix}$   
for the cases  $\epsilon > 0$ ,  $\epsilon = 0$ ,

## Example: (about Poincare Bendixson criterion)

Consider the case  $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x}$  where  $\mathbf{A} = \begin{bmatrix} 0 & 2 \\ -2 & \epsilon \end{bmatrix}$   
for the cases  $\epsilon > 0$ ,  $\epsilon = 0$ ,  $\epsilon < 0$ ,

## Example: (about Poincare Bendixson criterion)

Consider the case  $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x}$  where  $\mathbf{A} = \begin{bmatrix} 0 & 2 \\ -2 & \epsilon \end{bmatrix}$   
for the cases  $\epsilon > 0$ ,  $\epsilon = 0$ ,  $\epsilon < 0$ ,

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Converting to polar coordinates, we get

## Example: (about Poincare Bendixson criterion)

Consider the case  $\dot{\mathbf{x}} = \mathbf{A}\mathbf{x}$  where  $\mathbf{A} = \begin{bmatrix} 0 & 2 \\ -2 & \epsilon \end{bmatrix}$   
for the cases  $\epsilon > 0$ ,  $\epsilon = 0$ ,  $\epsilon < 0$ ,

$$\begin{bmatrix} \dot{x}_1 \\ \dot{x}_2 \end{bmatrix} = \begin{bmatrix} 0 & 1 \\ -1 & (25 - x_1^2 - x_2^2) \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix}$$

Converting to polar coordinates, we get

$$x_1 = r \cos \theta$$

$$x_2 = r \sin \theta$$

## Example (continued)

$$x_1^2 + x_2^2 = r^2$$

## Example (continued)

$$x_1^2 + x_2^2 = r^2$$

$$x_1 \dot{x}_1 + x_2 \dot{x}_2 = r \dot{r}$$



## Example (continued)

$$x_1^2 + x_2^2 = r^2$$

$$x_1 \dot{x}_1 + x_2 \dot{x}_2 = r \dot{r}$$

Substituting  $x_1, x_2, \dot{x}_1, \dot{x}_2$  using  $r, \dot{r}$  and  $\theta$ , we get  $r \dot{r} =$

## Example (continued)

$$x_1^2 + x_2^2 = r^2$$

$$x_1 \dot{x}_1 + x_2 \dot{x}_2 = r \dot{r}$$

Substituting  $x_1, x_2, \dot{x}_1, \dot{x}_2$  using  $r, \dot{r}$  and  $\theta$ , we get  $r \dot{r} =$

$$(r \cos \theta)(r \sin \theta) + (r \sin \theta)(-r \cos \theta + (25 - r^2)r \sin \theta)$$

This reduces to  $\dot{r} = (25 - r^2)r \sin^2 \theta$

Therefore at  $r = 5$  we get  $\dot{r} = 0$  i.e circle with radius 5 is a periodic orbit.

Moreover, other trajectories 'converge' to this limit cycle:

## Example (continued)

$$x_1^2 + x_2^2 = r^2$$

$$x_1 \dot{x}_1 + x_2 \dot{x}_2 = r \dot{r}$$

Substituting  $x_1, x_2, \dot{x}_1, \dot{x}_2$  using  $r, \dot{r}$  and  $\theta$ , we get  $r \dot{r} =$

$$(r \cos \theta)(r \sin \theta) + (r \sin \theta)(-r \cos \theta + (25 - r^2)r \sin \theta)$$

This reduces to  $\dot{r} = (25 - r^2)r \sin^2 \theta$

Therefore at  $r = 5$  we get  $\dot{r} = 0$  i.e circle with radius 5 is a periodic orbit.

Moreover, other trajectories 'converge' to this limit cycle: 'isolated stable limit cycle'.

# van der Pol oscillator

This oscillator has a 'stable limit cycle'.  
Differential equation

$$\ddot{x} - \epsilon(1 - x^2)\dot{x} + x = 0$$

where  $\epsilon > 0$

Let  $\dot{x} = y$  and  $\dot{y} = -x + \epsilon(1 - x^2)y$