

Modelling of carving turns in alpine skiing

Serguei S.Komissarov*

Department of Applied Mathematics

The University of Leeds

Leeds, LS29JT, UK

(Dated: Received/Accepted)

In this paper we present a simple mathematical theory of carving turns in alpine skiing and snowboarding and its application to runs on slopes of constant gradient. We find that pure carving is possible only on relatively flat slopes, with the critical slope angle around 17° for slalom, 19° for giant slalom and 21° for downhill skis. In wiggly carving runs on slopes of subcritical gradient, the aerodynamic drag force remains relatively unimportant and the speed stops growing well below the one achieved in the case of fall-line gliding runs. This is because the increased g-force of carving turns leads to enhanced snow friction which can significantly exceed the one found in gliding. At the critical gradient, the g-force of carving turns becomes infinite. For such and even steeper slopes only hybrid turns, where at least some part of the turn is skidded, are possible. The carving turns of all alpine racing disciplines are approximately similar. This is because in the dimensionless equations of the model the sidecut radius defines only the coefficient of the aerodynamic drag term, which always remains relatively small. Simple modifications to the model are made to probe the roles of skier's angulation and skidding at the transition phase of hybrid turns. As expected the angulation gives certain control over the turn trajectory, by modifying the local turn radius, whereas the hybrid technique allows skiing on steeper slopes.

Keywords: applied physics, mechanics, biomechanics, sport science, alpine skiing, carving, mathematical modelling

* s.s.komissarov@leeds.ac.uk

I. INTRODUCTION

When making their way down the hill, expert skiers execute complex coordinated body movements, often within a fraction of a second, which allow them to ski at speed and yet to remain in control. Their decisions are dictated by many factors, such as terrain, snow condition, equipment etc. A ski racer faces additional challenges as a race course significantly reduces the freedom of choosing trajectory. There is a great deal of qualitative understanding of skiing techniques and race tactics based on the personal experiences of ski professionals, coaches and instructors. When designing a race course, the setter has a pretty good idea of how the turns have to be executed in order to excel. All this knowledge is based on countless previous runs in similar circumstances and trial-and-error attempts to figure out what works and what does not. Understandably, this empiric knowledge is very subjective, often imprecise and sometimes even subconscious and this keeps the door open to misunderstanding, misconceptions and controversies.

Science can help to solidify this knowledge and to raise the level of understanding. This road has already been followed by many scientifically-minded skiing experts as well as academics. For example, in the book “Ultimate Skiing” by Ron LeMaster [7], the mechanical and bio-mechanical principles are used to elucidate the reasons behind skiing techniques and in “The Physics of Skiing” by Lind and Sanders [5] a rudimentary mathematical theory of skiing is developed. Few studies have tried to identify quantitative parameters of skiing turns which could tell to racers how to ski in order to minimise the run time [8–10]. First attempts have been made to mathematically model ski runs. These studies were limited to the so-called steered (or skidded) turns where skis are pivoted to an angle with the direction of motion, thus introducing enhanced dissipation of the skier’s kinetic energy via snow-ploughing and ice-scraping [1, 6, 12].

The advance of modern shaped skis has moved the focus of competitive skiing from skidded turns to carving turns. The main feature of a carving turn is a near-perfect alignment of skis with their trajectory, which significantly reduces the energy dissipation and increases the speed. Nowadays even mass-produced skis are shaped, thus giving the opportunity to enjoy carving runs to all skiing enthusiasts. This has made an impact on the way the alpine skiing is taught by ski instructors [e.g. 2]. The key property of shaped skis is their ability to bend into an arc when put on edge and pressed down in the middle, the higher the

inclination angle of the ski the tighter is the arc. When pressed against hard snow (or ice), such a ski cuts in it an arc-like platform which has the following two main outcomes: First this helps to inhibit ski's lateral displacement and hence skidding. Second, the shape of the bended ski (or rather of its inside edge) becomes the local shape of its trail in the snow.

Naively, one may think that skiers are free to change the inclination angle of their skis as they like and hence fully control the local shape (curvature) of their trajectory. However, this angle is largely dictated by the inclination angle of the skier, and an arbitrary change of this angle may upset skier's lateral balance. Indeed, as we all know a stationary skier must stay more-or-less vertical to avoid falling to the one side or the other. The theoretical analysis of the lateral balance in a carving turn allowed Jentshura and Fahrbach [3] to derive the so-called Ideal Carving Equation (ICE), which shows that the inclination angle, and hence the local radius of curvature of skier's trajectory, are completely determined by his speed and direction of motion relative to the fall line. To a mathematician, this immediately tells that the speed and direction of motion at the initiation of a carving turn (initial conditions) should determine the rest of the turn. The skier is almost like a passenger. In our paper we continue this analysis and formulate the key dynamic equations of carving turns. We also present a number of solutions describing simple runs down a slope of constant gradient, which reveal some intriguing properties.

The inclination is only one of several body manoeuvres executed by advanced skiers during their runs in order to maintain their balance. Another one is the lateral angulation, mostly at the hip but also at the knees. The angulation breaks the rigid relationship between the ski's and skier's inclination angles imposed by ICE and allows skiers to regain some control over their motion in pure carving turns. We generalise ICE by taking the angulation into account and analyse its role in some details.

Because the carving technique minimises the losses of kinetic energy and allows to ski faster, one could naively expect that ski racers would execute pure carving turns from start to finish, but this is not the case. The deviations from pure carving often manifest themselves in the form of snow plumes, ejected from under their skis. In many cases, this is the result of mistakes made by skiers while deciding when to terminate one turn and to start next one, mistakes which are corrected by braking. However, in this paper we argue that it may not be possible in principle to get rid of the skidding completely. In fact, ICE implies an existence of maximum speed consistent with the lateral balance in carving turns – no mater

how strong and skilful the athlete is, she/he would not be able to perform a carving turn at a speed exceeding this limit [3]. In our paper, we presents a somewhat different analysis of this issue and extend it to the case of angulated skier. We find that the speed limit is robust and then show that, unless the slope is very flat, some amount of skidding has to be introduced just to stay on the run. We modify our mathematical model of skiing accordingly and present the results for runs with a certain degree of skidding during transitions between the carving phases of turns.

II. AERODYNAMICS DRAG AND CHARACTERISTIC SCALES OF ALPINE SKIING

A. Fall-line gliding

Although recreational skiing can be very relaxed and performance skiing physically most demanding, the dominant source of energy in both cases is the Earth's gravity. The total available gravitational energy is

$$U = mgH, \quad (1)$$

where m is skier's mass, g is the gravitational acceleration and H is the total vertical drop of the slope. If all this energy was converted into the kinetic energy of the skier, $K = mv^2/2$, then at the bottom of the slope the speed would reach

$$v = \sqrt{2gH} \approx 227 \left(\frac{H}{200 \text{ m}} \right)^{1/2} \text{ km/h}. \quad (2)$$

In fact, this not far from what is achieved in the speed skiing competitions, where skiers glide straight down the fall line. For example, the current speed record of the Riberal slope in Grandvalira (Andorra), which has exactly 200m vertical drop and is often used for speed skiing competitions, is just short of 200km/h[11]. However, the typical speeds in other alpine disciplines are significantly lower, indicating that only a fraction of the available gravitational energy is converted into the kinetic energy of the skiers. In other words, there are some forces working against gravity. Two of the candidates are the dynamic snow friction and the aerodynamic drag [5]. The friction force is antiparallel to the skier velocity vector \mathbf{v} and its magnitude relates to the normal reaction force \mathbf{F}_n via

$$F_f = \mu F_n, \quad (3)$$

where μ is the dynamic coefficient of friction. For a brick of mass m residing on a firm surface, the normal reaction force is normal to this surface and balances the normal component of the gravity force:

$$F_n = mg \cos \alpha, \quad (4)$$

where α is the angle between the surface and the horizontal plane. This result also applies to a skier gliding down the fall line, like in the speed skiing competitions. Because the friction force determined by equations (3) and (4) does not depend on the skier's speed, it cannot limit its growth but only to reduce the growth rate.

The aerodynamic drag force is also antiparallel to the velocity vector and has the magnitude

$$F_d = \kappa v^2 \quad \text{where} \quad \kappa = \frac{C_d A \rho}{2}, \quad (5)$$

where C_d is the drag coefficient, A is the cross-section area of the skier normal to the direction of motion and ρ is the mass density of the air. The fact that the drag force grows with speed without limit implies that at some speed it will balance the gravity along the fall line. Below this speed, the gravity wins and skier accelerates. Above this speed, the drag wins and the skier decelerates. Hence the equilibrium speed is also a growth saturation speed (an attractor). The value of the saturation speed v_s can be easily found from the energy principle.

In the saturation regime, the work carried out by the drag and friction forces over the distance L along the fall line is

$$W = (F_f + F_d)L. \quad (6)$$

This must be equal to the gravitational energy $U = mgH$ released over the same distance. This leads to

$$v_s^2 = \frac{mg}{\kappa} \sin \alpha (1 - \mu \cot \alpha) \quad (7)$$

[cf. 5]. Incidentally, the result shows that the slope angle has to exceed $\alpha_{\min} = \arctan(\mu)$, as otherwise $v_s^2 < 0$, indicating that skiing is impossible. For the realistic value $\mu = 0.04$ [5] this gives $\alpha_{\min} = 2.3^\circ$. Usually ski slopes are significantly steeper than this and the snow friction contribution is small. In this case, the saturation speed is determined mostly by the balance between the gravity and aerodynamic drag, which yields the speed

$$V_g = \sqrt{\frac{mg}{\kappa} \sin \alpha}. \quad (8)$$

From this analysis it follows that V_g is a characteristic speed for the problem of fall-line gliding and presumably for alpine skiing in general. We note here that it depends on the slope gradient – the steeper is the slope, the higher is the speed. The time required to reach this speed under the action of gravity,

$$T_g = \frac{V_g}{g \sin \alpha} = \sqrt{\frac{m}{kg \sin \alpha}}, \quad (9)$$

is the corresponding characteristic time scale. This scale is shorter for steeper slopes. However, the corresponding length scale

$$L_g = V_g T_g = \frac{m}{k} \quad (10)$$

does not depend of the slope gradient, which is not very intuitive (L_g is twice the distance required to reach V_g under the action of gravity alone.). Lind & Sanders [5] state the values $C_d = 0.5$, $m = 80$ kg, $A = 0.4$ m², and $\rho = 1.2$ kg/m³ as typical for downhill (DH) competitions. For these parameters

$$V_g = 148 \sqrt{\frac{\sin \alpha}{\sin 15^\circ}} \text{ km/h}, \quad T_g = 16 \sqrt{\frac{\sin 15^\circ}{\sin \alpha}} \text{ s}, \quad L_g = 0.667 \text{ km}. \quad (11)$$

Though rather large, L_g is definitely smaller than the typical length of downhill tracks, which are usually few kilometres long. Similarly, T_g is significantly shorter compared to the typical duration of competition runs. This indicates that V_g must be well within reach during the runs and indeed it is quite close to the typical speeds recorded in downhill – the fastest speed ever recorded with a speed gun in downhill is $v \approx 162$ km/h (skier Johan Clarey, Wengen, 2013). Although downhill courses are not entirely straight, they still include a number of long gliding sections and this is why the gliding model works quite well here. (The significantly higher speed of speed skiing is due to the extreme measures undertaken by its participants in order to reduce the aerodynamic drag, and hence the lower typical values of κ .)

B. Skiing with turns

Slalom (SL) race tracks are much shorter than the downhill ones – for example the Ganslern track in Kitzbuhel is only about 590 meters long which is about the same as the

value L_g estimated above. Moreover, for a lighter slalom skier (smaller m) in a less compact position (higher κ), m/κ can be about twice as low, leading to

$$V_g = 104 \sqrt{\frac{\sin \alpha}{\sin 15^\circ}} \text{ km/h}, \quad T_g = 11 \sqrt{\frac{\sin 15^\circ}{\sin \alpha}} \text{ s}, \quad L_g = 0.333 \text{ km}. \quad (12)$$

Thus even in slalom, tracks are few times longer than the distance required to hit the drag-imposed limit. However, the typical speeds of slalom runs stay well below this limit, near or below 40 km/h (e.g. www.scholastic.com). For such low speeds, the aerodynamic drag is about six times below of what is required to balance gravity [see also 10] and hence skiers must be losing a lot of kinetic energy in a way not accounted for in equation (6).

On average, slalom slopes are steeper compared to the downhill ones. For example, the mean inclination angle of the Ganslern track is $\langle \alpha \rangle \approx 19^\circ$. This makes the discrepancy slightly more pronounced as according to equation (11) V_g increases with α (by approximately 12% in the case of the Ganslern). The fact that a slalom course is not a straight line, contrary to what is assumed in the derivation of Eq.(7), but wanders from side to side does not make much difference either. The wandering makes the skier path longer compared to the length L of the slope. However, even when each turn is a full half-circle, the length of skier's path $L' = (\pi/2)L$ is only about 50% longer. Replacing L with L' in Eq.(6) one finds that this results in a decrease of the saturation speed by only about 20%, compensating the increase due to steeper gradients.

Equation (7) assumes that no snow-ploughing or ice-scraping by the ski edges is involved. In the case of skiing with turns, this implies ideal carving where the skis are bend to adopt the local shape of their trajectory and leave on the snow two very narrow tracks. Moreover, it assumes that the skier's weight is just mg , which is not necessarily true. Indeed, turning introduces the centrifugal force, which leads to an increased effective weight of the skier and hence higher normal reaction and friction forces compared to the case of fall-line gliding. As the centrifugal force depends on the speed and local radius of skier's trajectory, it varies during the run and its overall effect cannot be estimated easily. A detailed mathematical model of carving turns is required for this purpose.

When watching alpine skiing competitions or instruction videos, one can see that pure carving turns are executed mostly on flat slopes, whereas on steeper slopes skiers use hybrid turns which are partly carved and partly skidded. The skidding introduces additional losses kinetic energy and could be another reason behind the fact that the typical speeds in slalom

are significantly below V_g .

III. THE MATHEMATICAL MODEL

Most phenomena surrounding us are usually complicated with many connections and numerous factors, whereas our mathematical models of them are normally quite simplified. The simplifications are usually driven by the understandable desire to keep the resultant mathematical equations treatable by known mathematical techniques. The typical evolution of a mathematical model starts from the most basic form that still captures some key factors and progresses to a more complex form where some additional factors, recognised as potentially important, are included as well. Since the theory of carving is in a rather immature state, the model which we present here is rather simplistic and ignores a number of factors that may end up being important. Moreover, the accounted for factors might have been included in a rather simplistic way. Yet we believe the model is useful as it captures the nature of carving rather well and shows the way forward.

Instead of listing all the simplifications from the start, we rather describe them as they come into force. The first one concerns the geometry of the slope. Here we limit ourselves to the idealised case of a plane slope with constant gradient and introduce such system of Cartesian coordinates $\{x, y, z\}$ associated with the slope that on its surface $z = 0$. The unit vectors parallel to the coordinate axes will be denoted as $\{\mathbf{i}, \mathbf{j}, \mathbf{k}\}$ respectively. For convenience, we direct the y -axis along the fall line, pointing downwards. We also introduce the vertical unit vector $\mathbf{s} = -\sin(\alpha)\mathbf{j} + \cos(\alpha)\mathbf{k}$, so that the gravitational acceleration $\mathbf{g} = -g\mathbf{s}$ (see figure 1).

In the model, we will focus mostly of the motion of skier's centre of mass (CM). We will also consider the torques that may force the skier to rotate about CM, focusing on prevention of falling to the side.

A. Basic dynamics of alpine skiing

When only the gravity, normal reaction, dynamic friction and aerodynamic drag forces are taken into account, the second law of Newtonian mechanics governing the motion skier




FIG. 1. Geometry of the slope. *Left panel:* The vertical section of the slope along the fall line. *Right panel:* The slope as seen at an angle from above. The curved line in the middle is the skier trajectory.

CM reads

$$m \frac{d\mathbf{v}}{dt} = \mathbf{F}_g + \mathbf{F}_f + \mathbf{F}_d + \mathbf{F}_n, \quad (13)$$

where

$$\mathbf{F}_g = m\mathbf{g}, \quad \mathbf{F}_f = -\mu F_n \mathbf{u}, \quad \mathbf{F}_d = -kv^2 \mathbf{u} \quad (14)$$

are the total gravity, friction, aerodynamic drag, and snow reaction force acting on the skier respectively and \mathbf{F}_n is the normal reaction. In these expressions, \mathbf{u} is the unit vector in the direction of motion. It is convenient to introduce the *angle of traverse* β as the angle between $-\mathbf{i}$ and \mathbf{u} for the right turns and between \mathbf{i} and \mathbf{u} for the left turns of a run (see figure 1). With this definition,

$$\mathbf{u} = \mp \cos(\beta) \mathbf{i} + \sin(\beta) \mathbf{j} \quad (15)$$

where the upper sign of $\cos \beta$ corresponds to the right turns and the lower sign to the left turns (we will use this convention throughout the paper).

The normal reaction force \mathbf{F}_n is not as easy to describe as the other forces. First, it is not normal to the slope surface but to the surface of contact between the snow and the skis. When skis are put on edge they carve a platform (or a step) in the snow and the normal reaction force is normal to the surface (surfaces) of this platform [7]. The connection between the orientation of this platform and the CM motion is not obvious. It is easy to see that it is parallel to the skier velocity but the rest is not immediately clear. Second,




FIG. 2. The unit velocity vector \mathbf{u} , its infinitesimal variation $d\mathbf{u}$ and the local centripetal unit vector \mathbf{c} .

this force equals the total effective weight of the skier, which is determined not only by the gravity but also by the centrifugal force. The latter depends not only on the skier speed but also on the acceleration, (in other words the local curvature of the trajectory.) Thus the system of equations (13) is not closed and we need to find a way of closing it.

Since the velocity vector $\mathbf{v} = v\mathbf{u}$ we have

$$\frac{d\mathbf{v}}{dt} = \mathbf{u} \frac{dv}{dt} + v \frac{d\mathbf{u}}{dt}. \quad (16)$$

Ignoring the up and down motion of CM, we can write $d\mathbf{u}/dt = \mathbf{c}|d\beta/dt|$, where the unit vector \mathbf{c} is parallel to the slope plane, perpendicular to the instantaneous velocity direction \mathbf{u} and points in the same direction as $d\mathbf{u}$ (see figure 2). \mathbf{c} is called the centripetal unit vector as it points towards the local centre of curvature. Hence

$$\frac{d\mathbf{v}}{dt} = \mathbf{u} \frac{dv}{dt} + v \mathbf{c} \left| \frac{d\beta}{dt} \right|. \quad (17)$$

Since $dt = dl/v$, where l is the distance measured along the skier trajectory, the last equation can also be written as

$$\frac{d\mathbf{v}}{dt} = \mathbf{u} \frac{dv}{dt} + \frac{v^2}{R} \mathbf{c}, \quad (18)$$

where

$$R = \left| \frac{dl}{d\beta} \right|. \quad (19)$$

R is called as the (local) radius of curvature of the trajectory. Hence we can rewrite equation (13) as

$$m\mathbf{u} \frac{dv}{dt} + \frac{mv^2}{R} \mathbf{c} = \mathbf{F}_g + \mathbf{F}_f + \mathbf{F}_d + \mathbf{F}_n. \quad (20)$$

Scalar multiplication of equation (20) with \mathbf{u} delivers the equation governing the evolution of skier's speed. Since

$$\mathbf{u} \cdot \mathbf{c} = 0, \quad \mathbf{u} \cdot \mathbf{u} = 1, \quad \mathbf{u} \cdot \mathbf{s} = -\sin \alpha \sin \beta,$$

this equation reads

$$\frac{dv}{dt} = g \sin \alpha \sin \beta - \mu \frac{F_n}{m} - \frac{k}{m} v^2. \quad (21)$$

Like in the case of fall-line gliding the speed is govern the gravity, friction and aerodynamic drag but now it also depends on the traverse angle. Notice that we do not utilise the expression (4) for F_n here. This is because, as we are about to show, its area of application is limited to the case of fall-line gliding only.

The normal reaction force can be decomposed into components parallel to \mathbf{c} and \mathbf{k} :

$$\mathbf{F}_n = F_{n,c} \mathbf{c} + F_{n,k} \mathbf{k}. \quad (22)$$

The scalar multiplication of equation (20) and \mathbf{k} immediately yields

$$F_{n,k} = mg \cos \alpha. \quad (23)$$

Thus, the normal to the slope component of the snow reaction force \mathbf{F}_n is the same as in the case of the fall-line gliding. This is what is needed to match exactly the normal to the slope component of the gravity force and hence to keep the skier on the slope.

Since

$$\mathbf{c} = \pm \sin(\beta) \mathbf{i} + \cos(\beta) \mathbf{j} \quad (24)$$

(see figure 3) and hence $(\mathbf{s} \cdot \mathbf{c}) = -\sin \alpha \cos \beta$, the scalar multiplication of equation (20) with \mathbf{c} yields

$$F_{n,c} = \frac{mv^2}{R} - mg \sin \alpha \cos \beta. \quad (25)$$

Now we can find the angle between \mathbf{F}_n and the normal direction to the slope \mathbf{s}

$$\tan \Phi = \frac{F_{n,c}}{F_{n,k}} = \frac{v^2}{gR} \frac{1}{\cos \alpha} - \tan \alpha \cos \beta. \quad (26)$$

Equation (25) has the appearance of a force balance. However this not the case as mv^2/R is not a force but the skier mass multiplied by its radial acceleration. Strictly speaking (25) is a differential equation, whose time derivative is hidden away through the definition of R .




FIG. 3. The centripetal unit vector \mathbf{c} for left and right turns.

It defines the component of normal reaction along the slope plane which is needed to force a curvilinear motion with the local radius of curvature R .

Formally, equations (23,25) can be written as one vector equation

$$\mathbf{F}_n + \mathbf{F}_c + \mathbf{F}_{g,\text{lat}} = 0, \quad (27)$$

where

$$\mathbf{F}_{g,\text{lat}} = -(mg \cos \alpha) \mathbf{k} + (mg \sin \alpha \cos \beta) \mathbf{c} \quad (28)$$

is the lateral (normal to \mathbf{u}) component of the gravity force and

$$\mathbf{F}_c = -(mv^2/R) \mathbf{c}. \quad (29)$$

Equation (27) also holds in the accelerated (non-inertial) frame of the skier. In this frame, the skier is at rest and equation (27) is a balance law between the normal reaction, gravity and the inertial force \mathbf{F}_c called the centrifugal force. This inertial force has the same properties as the gravity force and can be combined with the lateral gravity into the effective gravity force

$$\mathbf{F}_{g,\text{eff}} = \mathbf{F}_{g,\text{lat}} + \mathbf{F}_c, \quad (30)$$

turning equation (27) into the balance between the normal reaction force and the effective gravity

$$\mathbf{F}_n + \mathbf{F}_{g,\text{eff}} = 0. \quad (31)$$

Now we can find the total strength of the normal reaction force (and hence the effective weight of the skier) as

$$F_n^2 = \left(\frac{mv^2}{R} \right)^2 - 2 \frac{m^2 v^2 g}{R} \sin \alpha \cos \beta + m^2 g^2 (\cos^2 \alpha + \sin^2 \alpha \cos^2 \beta). \quad (32)$$




FIG. 4. Upper-C and Lower-C sections of the right and left turns.

It is easy to see that in the case of a fall-line glide, where $R = \infty$ and $\beta = 90^\circ$, this equation reduces to the familiar $F_n = mg \cos \alpha$, as expected. It is also clear that the normal reaction force is stronger in the Lower-C part of the turn ($90^\circ < \beta < 180^\circ$), where the gravity force and the centrifugal force act in the same direction, and weaker in the Upper-C ($0^\circ < \beta < 90^\circ$), where they act in opposite directions (see figure 4. The terminology is acquired from [2].).

The weight is often measured in the units of the standard weight mg , in which case it is called a g-force. Hence we have got

$$\text{g-force} = \left[\left(\frac{v^2}{gR} \right)^2 - 2 \left(\frac{v^2}{gR} \right) \sin \alpha \cos \beta + (\cos^2 \alpha + \sin^2 \alpha \cos^2 \beta) \right]^{1/2}. \quad (33)$$

The expressions (32,26) for the strength and direction of \mathbf{F}_n do not allow to close the system. Indeed, they involve the turn radius R which remains undetermined yet. It has entered the model via equation (19) which cannot be used to calculate it as it requires to know the trajectory, which is not known until the whole solution to the problem is found. On the other hand, it is known that for caving turns with shaped skis R is largely determined by the ski's inclination to the slope [5]. In order to introduce R into our model via this route, we need to go beyond the CM dynamics, take into account the overall shape of a skier and hence analyse the torques they may experience.

B. Turn radius

In the mechanics of solid bodies, by a balance we understand not only vanishing combined (total) force but also vanishing combined torque [4]. Although skiers are not exactly solid bodies, torques are still important in their dynamics. Indeed, the combined force governs the motion of skier's CM only. The combined torque determines if the skier stays in control or ends up over-rotated and even crashing. When discussing torques it is important to where each of the involved forces are applied. Both the centrifugal and the gravity force (and hence the effective gravity) per unit mass are uniform and hence one may think of these forces as being applied directly at the CM. This is not true for the snow friction and normal reaction forces, which are applied at the skis, and for the aerodynamic force which is distributed over the skiers body in a rather complicated way.

When discussing torques acting on a skier it is convenient to use the skier's frame of reference and introduce the three mutually orthogonal fundamental planes passing through skier's CM : the frontal, sagittal and transverse planes[7]. The frontal plane is normal to \mathbf{u} , the sagittal plane is parallel to \mathbf{u} and \mathbf{F}_n , the transverse plane is defined as the orthogonal to the other two. In the transverse plane, the total torque does not have to vanish as skiers often rotate in this plane in the same sense as the skis (though a bit of counter-rotation can be used as well). In the other two planes, the total torque must be close to zero. If the torque in the sagittal plane did not vanish, the skier would fall forward (or backward) and if the torque in the frontal plane did not not vanish the skier would fall to the inside (or outside) of the turn.

In our model, we will assume that everything is fine with the skier balance in the sagittal and transverse planes and consider in details only the balance in the frontal plane as it is the one which determining the turn radius. How we analyse torques in this plane depends on the nature of skier's contact with the snow. It can be either (1) a sliding contact, in which case we have to deal with the torques about skier's CM or (2) the skis can be regarded as pinned to the slope at the inside edge, in which case we should consider torque about the edge. In carving, it is normally the second case, but in both the cases the total torque vanishes under the same condition, namely both the normal reaction force and the effective gravity have to be applied along the line connecting CM with the inner edge. This is illustrated in figure 5 where the loading of the inside (upper) ski is ignored. If the skier is stacked (not angulated




FIG. 5. Balance of a stacked (un-angulated skier) in the frontal plane. The normal reaction force balances the effective gravity force and introduces zero torque because it points directly at the skier's centre of mass (CM). This configuration corresponds to the Lower-C part of the turn.

at the hip or knee), then this condition requires the skier's body to be aligned with \mathbf{F}_n and hence $\mathbf{F}_{g,eff}$.

By its nature, \mathbf{F}_n is normal to the contact surface with the snow. If this is just a part of the bottom surface of the loaded ski located near the inside edge then the balance condition requires it to be normal to the line connecting the edge with the CM, which in the case of a stacked skier is located about their belle button. This is reflected in the overall design of the skiing equipment (boots, skis, bindings). However, the exact outcome also depends on the personal physiological features of individual skiers and a fine tuning is normally carried out via the so-called canting of ski boots [7]. (The issue appears less important when we take into account that most skiers angle their body in a particular way which prevents such torque and the side-slipping it causes. We consider the angulation later in the paper.) Here we will assume that the ski boots are properly canted and hence in the balanced position the skis are inclined to the slope at the same angle Φ as the effective gravity and normal reaction forces.

Now we consider the geometry of a carving ski and how it changes when placed at the inclination angle Ψ to a flat surface, which we assume to be hard and hence not changed in the process apart from a small cut possibly made in it by the ski's sharp steel edges. We




FIG. 6. *Left panel:* Shaped ski, its sidecut h_{sc} and sidecut radius R_{sc} . The dashed line shows the ski tip which is not important for determining R_{sc} . *Right panel:* The same ski with the edge BDF pressed against a hard flat surface at the inclination angle Ψ as seen in projection on the plane normal to the ski at its waist. In this projection, DGB is a right angle triangle.

start with the case where the ski lays flat as shown in the left panel of figure 6. Its edges ACE and BDF can be approximated as circular arcs of radius R_{sc} , called the sidecut radius. Denote as CD the waist of the ski and as l the distance between B and D (or A and C) along the edge. As seen in this figure, $l = R_{sc}\delta$, where δ is the angular size of the edge DB as seen from its centre of curvature. This angle is normally rather small. For an SL ski of length $l_{ski} \approx 2l = 1.65$ m and $R_{sc} = 12.7$ m we have $\delta \approx 0.065$ (3.7°). For other kinds of racing skis, it is even smaller. The sidecut depth h_{sc} shows by how much the ski edge is receded compared to the straight line BF at the waist. From the right angle triangle shown in the figure

$$h_{sc} = R_{sc}(1 - \cos \delta). \quad (34)$$

Using the first two terms of the Maclaren expansion for $\cos \delta$

$$\cos \delta = 1 - \frac{1}{2}\delta^2 + O(\delta^4)$$

and then substituting $\delta = l/R_{sc}$ one finds the approximation

$$h_{sc} \simeq \frac{l^2}{2R_{sc}}. \quad (35)$$

Now suppose that this ski is kept at the angle Ψ to a firm flat surface and that it is pressed in the middle until its lower edge comes into the contact with the surface along its whole length (excluding the tip). In this position the edge can still be approximated as an arc but a different one. Denote its radius as R and its “sidecut” depth as h (see the right panel of figure 6). Obviously R and h are connected in the same way as R_{sc} and h_{sc}

$$h = R(1 - \cos \delta) . \quad (36)$$

where now $\delta = l/R$. When $\delta \ll 1$, this is approximated as

$$h \simeq \frac{l^2}{2R} \quad (37)$$

and hence

$$\frac{R}{R_{\text{sc}}} = \frac{h_{\text{sc}}}{h} . \quad (38)$$

Analysing the right angle triangle GDB of the right panel in figure 6, one finds that $h_{\text{sc}} = h \cos \Psi$ and hence

$$\frac{R}{R_{\text{sc}}} = \cos \Psi \quad (39)$$

(This is Eq.3.7 in [5].). In ideal carving turns, there is no side-slippage and the ski is transported along its curved loaded edge. This makes the ski curvature radius R the same as the local curvature radius of the turn trajectory.

Combining equations (39) and (26) with the balance condition $\Phi = \Psi$ one finds that

$$\left(\left(\frac{R_{\text{sc}}}{R} \right)^2 - 1 \right)^{1/2} = \frac{v^2}{gR \cos \alpha} - \tan \alpha \cos \beta . \quad (40)$$

This equation, called by Jentschura and Fahrbach the Ideal-Carving Equation [3], defines R as a function of skier’s velocity and therefore allows to close the CM equations of motion. Hence it allows deterministic mathematical modelling of carving runs using ordinary differential equations. In practical terms this means that in a carved turn the skier has no control over its local curvature and hence trajectory! What is up to the skier is the decision when to switch from one turn to another and carve the next arc.

According to equation (39), $R \rightarrow R_{\text{sc}}$ as $\Psi \rightarrow 0$ and hence the turn radius of marginally edged ski is exactly R_{sc} . Moreover, $R < R_{\text{sc}}$ for $\Psi \neq 0$ and it monotonically reduces with Ψ . For $\Psi \rightarrow 90^\circ$ we have $R \rightarrow 0$. Obviously, $R = 0$ corresponds to a ski curled into a point, which does not make any sense. This result reflects the fact that the approximation (37)

breaks down for $\delta = l/R \geq 1$. Although we are not aware of relevant experiments, it feels all but impossible to bend a ski into an arc of the radius as small as the ski length without actually breaking it. For $R = l_{\text{ski}}$ the exact equations (35,36) yield the inclination angle $\Phi_{\text{s,max}} = 80^\circ$, which can be considered as an upper limit on Ψ .

The upper limit on Φ can also be set by the athlete's ability to sustain high g-forces building up via centrifugal acceleration. In the case of a slope with $\alpha = 0^\circ$, equation (32) relates the g-force directly to the inclination angle:

$$\text{g-force} = \sqrt{1 + \tan^2 \Phi}. \quad (41)$$

From this equation we find that $\Phi = 80^\circ$ corresponds to g-force=5.7, which is extremely high. According to LeMaster[7] the best athletes can sustain g-force up to only about 3 corresponding to the less extreme $\Phi = 70^\circ$.

IV. SPEED LIMITS IMPOSED BY THE IDEAL CARVING EQUATION

Equation (40) can be written in the form

$$(\xi^2 - 1)^{1/2} = a\xi + b, \quad (42)$$

where $\xi = R_{\text{sc}}/R$, $a = v^2/gR_{\text{sc}} \cos \alpha$ and $b = -\cos \beta \tan \alpha$. From the definitions it is clear that

$$\xi \geq 1, \quad a > 0 \quad \text{and} \quad -\tan \alpha < b < +\tan \alpha.$$

When squared, equation (42) becomes quadratic, which is easy to analyse, but this operation may introduce an additional root and for this reason we will deal with this equation as it is.

The right-hand side of (42) is the linear function $g(\xi) = a\xi + b$. Since its gradient $g'(\xi) = a > 0$ this function increases with ξ . The left-hand side of (42) is the radical function $f(\xi) = \sqrt{\xi^2 - 1}$. Since $f'(\xi) = \xi/\sqrt{\xi^2 - 1} > 0$ it also grows with ξ . Moreover, $f'(\xi) \rightarrow +\infty$ as $\xi \rightarrow 1$ and $f'(\xi) \rightarrow 1$ as $\xi \rightarrow +\infty$. Finally, $f''(\xi) = -1/\sqrt{\xi^2 - 1} < 0$ and hence $f'(\xi)$ decreases monotonically from $+\infty$ to 1 for $\xi \in [1, +\infty)$.

A. The case $a > 1$

In this case, the gradient of $g(\xi)$ is higher than the asymptotic gradient of $f(\xi)$ and as one can see in the left panel of figure 7 there are three distinguish possibilities with either




FIG. 7. Finding roots of the ideal carving equation (ICE). The solid line shows $f(\xi) = \sqrt{\xi^2 - 1}$ and the dashed lines $g(\xi) = a\xi + b$. The roots of ICE are given by the intersection points of the curves. *Left panel:* The case $a > 1$. Depending on the value of b , one can have none, one or two roots. *Right panel:* The case $0 < a < 1$. Now one can have either one root or none.

none, one or two roots. First, if $g(1) = a + b < 0$ then there exists one and only one root. Indeed, in this case $g(1) < f(1)$ but the faster growth of $g(\xi)$ for large ξ ensures that it will eventually overcome $f(\xi)$ (see the lower dashed line in the left panel of figure 7).

As we increase b above $-a$, initially there are two roots but eventually they merge and disappear. The bifurcation occurs at the point b_c where

$$f(\xi_c) = g(\xi_c) \quad \text{and} \quad f'(\xi_c) = g'(\xi_c).$$

Solving these two simultaneous equations for ξ_c and b_c , we find that

$$b_c = -(a^2 - 1)^{1/2} \tag{43}$$

and

$$\xi_c = \frac{a}{(a^2 - 1)^{1/2}}. \tag{44}$$

There are no solutions when $b > b_c$, which includes all positive values of b and hence all $\beta > 90^\circ$ (Lower-C).




FIG. 8. The parameter space of carving turn. The regions where ICE has solutions are shaded, in grey where the solution is unique and in cyan where there are two solutions.

B. The case $a < 1$

In this case, the gradient of $g(\xi)$ is lower than the asymptotic gradient of $f(\xi)$ and as illustrated in the right panel of figure 7 there is either one solution or none. The corresponding condition for existence is now

$$a + b \geq 0. \quad (45)$$

This includes the case of $b > 0$ and hence there is always a unique solution for the Lower-C.

If $a + b < 0$ then there is no solutions. This condition implies negative b and hence the Upper-C part of the turn.

The results for both cases are collected in figure 8.

C. Interpretation.

In terms of the usual physical parameters, the condition $a < 1$ reads

$$v < V_{sc} \quad \text{where} \quad V_{sc} = \sqrt{g R_{sc} \cos \alpha} \quad (46)$$

This is the same as equation (36) in [3]. For $v > V_{sc}$, there are no solutions for sufficiently large angle of traverse, including the whole of the Lower-C. Thus, the speed should not be too high or the skier will not be able to carve Lower-C.

What are the indicators that a skier is about to hit the speed limit? Consider the entrance

point to the Lower-C where $\beta = 90^\circ$, $b = 0$ and hence equation (42) reads

$$(\xi^2 - 1)^{1/2} = a\xi.$$

Its solution

$$\xi = \frac{1}{\sqrt{1 - a^2}} \rightarrow \infty \quad \text{as} \quad a \rightarrow 1.$$

Thus the turn becomes very tight (formally $R \rightarrow 0$) and the skier's body close to horizontal (formally $\Phi \rightarrow 90^\circ$).

However, when the speed limit is exceeded in the Upper-C there may not be such a clear indicator. In fact, after this the turn can be continued for a while until it reaches the critical traverse angle β_c where

$$\cos \beta_c = \cot \alpha \left(\left(\frac{v}{V_{sc}} \right)^4 - 1 \right)^{1/2}$$

and

$$\tan \Phi_c = \left(\left(\frac{v}{V_{sc}} \right)^4 - 1 \right)^{-1/2}. \quad (47)$$

After this point the lateral balance can no longer be sustained. Thus when v grows slowly at the point of going over the limit and stays only just above V_{sc} , the crash occurs close to the fall line at extreme inclination angle. If however the growth is fast, v may significantly overshoot V_{sc} quite early in the High-C. In this case the crash may also occur early at Φ significantly below 90° . As we shall see in Sec.VI C this may occur on very steep slopes.

When $v < V_{sc}$ and the carving turn is possible in the Lower-C, it may still not be possible in the Upper-C. This is what is governed by the condition (45). In terms of the turn speed and the angle of traverse this condition reads

$$\cos \beta < \frac{v^2}{gR_{sc} \sin \alpha}. \quad (48)$$

It is convenient to rewrite this as a condition on the angle between the skier velocity and the fall line, $\gamma = |\beta - 90^\circ|$,

$$\sin \gamma < \frac{v^2}{gR_{sc} \sin \alpha}. \quad (49)$$

Thus, for sufficiently low speeds, namely $v^2 < gR_{sc} \sin \alpha$, carving is possible only close to the fall line. To allow fully rounded turns, with γ approaching 90° in the transition between the Lower-C and Upper-C parts of the turn, the speed must satisfy the condition

$$v > \sqrt{gR_{sc} \sin \alpha}. \quad (50)$$

For a half-circle turn, which starts and finishes perpendicular to the fall line ($\beta = 0, \pi$) the skier speed has to be within the range

$$\sqrt{gR_{\text{sc}} \sin \alpha} < v < \sqrt{gR_{\text{sc}} \cos \alpha}, \quad (51)$$

which can be quite narrow for sufficiently large α and collapse completely for $\alpha \geq 45^\circ$, making such turns impossible.

It is easy to see that

$$V_{\text{sc}} = V_{\text{g}} \left(\frac{R_{\text{sc}}}{L_{\text{g}} \tan \alpha} \right)^{1/2}. \quad (52)$$

For the typical parameters of slalom competitions, this gives

$$V_{\text{sc}} = 0.3V_{\text{g}} \left(\frac{R_{\text{sc}}}{13 \text{ m}} \right)^{1/2} \left(\frac{L_{\text{g}}}{400 \text{ m}} \right)^{-1/2} \left(\frac{\tan \alpha}{\tan 20^\circ} \right)^{-1/2}. \quad (53)$$

Thus, for carving turns in slalom V_{sc} is significantly smaller than the characteristic speed V_{g} (see eq.8) set by the aerodynamic drag. This suggests that pure ideal carving turns are normally impossible for the typical parameters of slalom competitions and the racers have to shave their speed via skidding on a regular basis. The only exceptions are probably (i) very flat sections of the track where the last factor of equation (53) can be sufficiently large, (ii) the first few turns right after the start, where the speed has not yet approached V_{sc} and (iii) the first turns after transitions from steep to flat. In the last case, V_{sc} significantly increases at the transition, creating the opportunity for accelerating pure carving turns. It is easy to verify that the situation is quite similar in other race disciplines. For example, for the downhill parameters

$$V_{\text{sc}} = 0.52V_{\text{g}} \left(\frac{R_{\text{sc}}}{50 \text{ m}} \right)^{1/2} \left(\frac{L_{\text{g}}}{700 \text{ m}} \right)^{-1/2} \left(\frac{\tan \alpha}{\tan 15^\circ} \right)^{-1/2}. \quad (54)$$

V. DIMENSIONLESS EQUATIONS OF CARVING TURN

It is a common practice of mathematical modelling to operate with dimensionless equations, which are derived using a set of scales characteristic to the problem under consideration, instead of standard of units. This leads to simpler equations which are easier to interpret and to the results which are at least partly scale-independent. The analysis of fall-line gliding given in Section II suggests that V_{g} , T_{g} and L_{g} are good candidates for the characteristic speed, time, and length scales of alpine skiing. However, the analysis of ICE

given in Sec.IV shows that in ideal carving the speed remains significantly below V_g and suggests to use V_{sc} as a speed scale instead. Moreover, the turn radius is bound to remain below R_{sc} which suggest that this is a more suitable length scale. The corresponding time scale is $T_{sc} = R_{sc}/V_{sc}$. These are the scales we adopt here.

In order to derive the dimensionless equations we now write

$$v = V_{sc}\tilde{v}, \quad t = T_{sc}\tilde{t}, \quad R = L_{sc}\tilde{R},$$

substitute these into the dimensional equations and where possible remove common dimensional factors. Finally, we ignore tilde in the notation. In other words, we do the substitutions $v \rightarrow V_{sc}v$, $t \rightarrow T_{sc}t$, $R \rightarrow R_{sc}R$ and then simplify the results. For example, substituting $v \rightarrow V_{sc}v$ into the equation (7) gives the dimensionless equation for the saturation speed of fall-line gliding as

$$v_s = Sr\sqrt{1 - \mu \cot \alpha}, \quad (55)$$

which includes the dimensionless speed parameter $Sr = V_g/V_{sc}$. Similarly we deal with other dimensional variables should they appear in the equations, e.g. $x \rightarrow R_{sc}x$ and $y \rightarrow R_{rc}y$. In particular, the application of this procedure to (21) gives the dimensionless speed equation

$$\frac{dv}{dt} = \sin \beta \tan \alpha - \mu F_n - Kv^2, \quad (56)$$

where

$$K = \frac{R_{sc}}{L_g}, \quad (57)$$

is a dimensionless parameter which we will call the dynamic sidecut parameter and

$$F_n^2 = 1 + \frac{v^4}{R^2} - 2 \tan \alpha \cos \beta \frac{v^2}{R} + \cos^2 \beta \tan^2 \alpha, \quad (58)$$

is the dimensionless effective weight. The g-force, which was defined as a dimensionless quantity in the first place, is related to F_n via

$$\text{g-force} = \cos(\alpha)F_n. \quad (59)$$

The dimensionless ideal-carving equation (40) reads

$$\left(\frac{1}{R^2} - 1 \right)^{1/2} = \frac{v^2}{R} - \cos \beta \tan \alpha. \quad (60)$$

The equation governing the evolution of β follows from equation (19) upon the substitution $dl = vdt$. It reads

$$\frac{d\beta}{dt} = \frac{v}{R} \quad (61)$$

and the dimensionless skier coordinates can be found via integrating

$$\frac{dx}{dt} = \mp v \cos \beta \quad (62)$$

and

$$\frac{dy}{dt} = v \sin \beta. \quad (63)$$

Equations (56-63) determine the arc of a carving turn and the skier motion along the arc. What they do not tell us is when one turn ends and the next one begins. These transitions have to be introduced independently. In this regard the angle of traverse is a more suitable independent variable than the time because its value is a better indicator of how far the turn has progressed. Replacing t with β via equation (61) we finally obtain the complete system of equations which we use to simulate carving runs in this study. It includes three ordinary differential equations

$$\frac{dv}{d\beta} = \frac{R}{v} (\sin \beta \tan \alpha - \mu F_n - Kv^2), \quad (64)$$

$$\frac{dx}{d\beta} = \mp R \cos \beta, \quad (65)$$

$$\frac{dy}{d\beta} = R \sin \beta, \quad (66)$$

where

$$F_n = \left(1 + \frac{v^4}{R^2} - 2 \tan \alpha \cos \beta \frac{v^2}{R} + \cos^2 \beta \tan^2 \alpha \right)^{1/2} \quad (67)$$

and the constitutive equation

$$\left(\frac{1}{R^2} - 1 \right)^{1/2} = \frac{v^2}{R} - \cos \beta \tan \alpha. \quad (68)$$

The definition of β implies that it increases both during the left and the right turns, but not in the transition between turns. In the ideal transition the direction of motion \mathbf{u} remains unchanged and hence the angle of traverse has to change from β to $180^\circ - \beta$. According to equation (68) this implies a jump in the local turn radius and hence the skier inclination. Hence during the whole run, which may consist of many turns, β remains confined between 0° and 180° , provided each turn terminates before going uphill.

Finally, we observe that equations (64) and (68) do not involve x and y and hence can be solved independently from equations (65-66). However, all these equations should be integrated simultaneously when we are interested in skier's trajectory. Equation (61) should be added as well when the time characteristics are important.

VI. IDEAL CARVING SOLUTIONS

Here we present the results of our study of ideal carving runs as described by equations (64-68) with instantaneous ideal transitions between turns. This means the jump $\beta \rightarrow 180^\circ - \beta$ at a single point of skier's trajectory, which is the termination point of the previous turn and at the same time the initiation point of the next one. At the start we specify the initial angle of traverse β_{ini} and speed v_{ini} , achieved during the run-up phase. Each turn is terminated when the traverse angle reaches a given value, denoted as β_{fin} . Hence, beginning from second one, all turns are initiated at $\beta_{\text{ini}} = 180^\circ - \beta_{\text{fin}}$. In all the runs we set the coefficient of friction to $\mu = 0.04$. In this section we focus on the effect of slope steepness and fix the sidecut parameter to $K = 0.0325$. For the aerodynamic length scale $L_g = 400$ m this corresponds to SL skis with the dimensional radius of $R_{\text{sc}} = 13$ m. Later we discuss the effect of sidecut radius as well.

A. Flat slope

First we consider a slope with $\alpha = 5^\circ$. For such a flat slope, the friction plays an important role even in the case of fall-line gliding where it keeps the speed well below V_g . Indeed, for $\beta = 90^\circ$ equation (55) implies that the gliding speed stops growing after reaching $v_s = 0.737V_g$. Secondly, equation (53) gives the speed limit $V_{\text{sc}} = 0.612V_g$, indicating that in carving runs the speed may eventually get over the speed limit.

Here we present the results for the initial data $x_{\text{ini}} = y_{\text{ini}} = 0$, $\beta_{\text{ini}} = 0.3\pi$ (54°), and $v_{\text{ini}} = 0.49V_{\text{sc}}$ ($0.3V_g$). Each turn terminates when the angle of traverse reaches $\beta_{\text{fin}} = 0.9\pi$ (162°). Figure 9 shows the trajectory for the few first turns of the run, which indicates a rather slow evolution of the turn shape. This is confirmed by the data presented in Figure 10, which shows the evolution of R , v , Φ and the g-force for the first 20 turns. One can see that the turn radius does not vary much along the track. Moreover, although each next turn is not an




FIG. 9. The first few turns of the flat-slope run ($\alpha = 5^\circ$).

exact copy of the previous turn, for each of the variables we observe a convergence to some limiting curve, which we will refer to as the asymptotic turn solution. (This is reminiscent of the so-called limit cycle solutions in the theory of dynamical systems.) In practical terms, well down the slope each next turn becomes indistinguishable from the previous one.

Interestingly, the speed of the asymptotic solution remains everywhere below V_{sc} , with its mean value $\langle v \rangle \approx 0.75V_{sc}$. As we mentioned in Section II, this can be down to the longer length of our wiggly trajectory compared to the length of fall line and the higher friction force due to the centrifugally-increased skier weight. In order to test this explanation one can use the equilibrium version of the speed equation (64)

$$\langle \sin \beta \rangle \tan \alpha - \mu \langle F_n \rangle - K \langle v^2 \rangle = 0, \quad (69)$$

where $\langle \rangle$ stands for the mean turn value. One can estimate $\langle \sin \beta \rangle$ via

$$\langle \sin \beta \rangle = \frac{1}{0.8\pi} \int_{0.1\pi}^{0.9\pi} \sin(\beta) d\beta \approx 0.796.$$

According to figure 10, $\langle F_n \rangle \approx \langle \text{g-force} \rangle \approx 1.2$. Substituting the estimates into equation (69) we find $\langle v \rangle \approx 0.8V_{sc}$, which is quite close to what we have actually see in the numerical data. Using equations (12,52), we find the corresponding dimensional value $\langle v \rangle \approx 27 \text{ km/h}$.

B. Moderately steep slope

The setup of this run differs with that of the flat slope run only by the slope gradient, $\alpha = 15^\circ$. In this case, equation (55) gives the saturation speed of a fall-line run $v_s = 0.922V_g$.




FIG. 10. Evolution of $R(\beta)$, $v(\beta)$, $\Phi(\beta)$ and F_n for the flat-slope run ($\alpha = 5^\circ$) during the first 20 turns. In each of these plots, there are 20 curves which show the evolution of these variables during each turn. Each such curve (except the one corresponding to the first turn which originates at $\beta = 0.3\pi$) originates at $\beta = 0.1\pi$ (the turn initiation point) and terminated at $\beta = 0.9\pi$ (the turn completion point). The transition between turns is a jump from the termination point of the previous turn to the initiation point of the next turn. In the R -panel this transition is a jump to the lower curve and in the v -, Φ - and F_n -panels to the upper curve.

The fact that it is very close to V_g tells us that for this gradient the role of the friction force in determining v_s is substantially smaller than in the flat case. On the other hand, equation (53) gives $V_{sc} = 0.35V_g$. This is significantly below v_s , suggesting that the skier speed will eventually exceed the carving upper limit, hence leading to a crash. However, this is not




FIG. 11. The first few turns for the moderately steep slope ($\alpha = 15^\circ$).

what we find in the simulations.

Figure 11 shows the turns of this run on the same section of the slope as in figure 9 for the flat run. Obviously, now the turns are much shorter and not so rounded, with the individual turn shape reminiscent of the letter “J”, rather than “C”. Figure 12 shows the evolution of R , v , Φ and the g-force for the first 20 turns of this run. One can see that like in the flat slope case the solution converges to an asymptotic one. Now this occurs very quickly – in figure 12 the curves become indistinguishable beginning from the forth turn. The top-left panel of figure 12 confirms that on average the turn radius is significantly lower than for the flat run. Moreover it varies dramatically, from $R \approx 0.6R_{sc}$ at the turn initiation down to $R \approx 0.12R_{sc}$ soon after the fall-line. The latter is approximately 1.6m (one SL ski length) when the solution is scaled to $R_{sc} = 13\text{m}$. The corresponding inclination angles and g-force also vary dramatically, with the g-force reaching unbearable heights. Overall, we conclude that while the theoretical solution exists, it is unlikely that such runs will ever be made in real life.

Surprisingly, the speed of the limiting solution remains just below V_{sc} and well below V_g . The latter indicates that contrary to the expectations based on the analysis of fall-line gliding, in this slalom run the aerodynamic drag is not the dominant factor in determining the saturation speed. The reason is the extremely high effective weight and hence the friction force. Indeed, according to the data the aerodynamic term of equation (64) $Kv^2 \approx 0.03$ whereas even for the lowest value of $F_n \approx 1.5$ found at the start of the turn the friction term $\mu F_n \approx 0.06$ is higher than this.

The corresponding dimensional value of the mean skier speed in the limiting solution is $v \approx 35 \text{ km/h}$. This is indeed very close to the typical speed of slalom runs, indicating that




FIG. 12. The same as in figure 10 but for the moderately steep slope ($\alpha = 15^\circ$).




FIG. 13. The trajectory of ideal carving run for the steep slope ($\alpha = 35^\circ$).




FIG. 14. The same as in figure 10 but for the steep slope ($\alpha = 35^\circ$).

racers ski very close to the carving limit.

C. Steep slope

The setup of this run differs from the previous ones by the higher slope gradient, $\alpha = 35^\circ$. For this gradient the difference between the aerodynamic and carving speed limits is even more dramatic than for the moderately steep slope, with $V_{sc} = 0.2V_g$. In this run the initial speed is set to $v_{ini} = 0.46V_{sc}$ ($0.1V_g$). For these parameters, the solution becomes singular even before reaching the fall-line of the second turn and cannot be continued thereafter (see figure 13). Figure (14) reveals the nature of the problem. Already during the first turn the skier is forced into an extreme inclination and experiences overwhelming g-force. In




FIG. 15. The trajectories of ideal carving runs on the $\alpha = 15^\circ$ slope with SL (solid), GS (dashed) and DH (dash-dotted) skis.

real life, skier's legs would collapse at this point but in the model the run continues. Very soon after the transition into the second turn the speed shoots above the carving limit and the solution approaches the point where no equilibrium position for the skier can be found. Hence the run cannot be continued even theoretically. As one can see, this occurs still in the Upper-C at the point where the turn radius remains relatively large and the skier inclination relatively small. We have discussed this possibility in Section IV C.

VII. THE APPROXIMATE SIMILARITY OF CARVING RUNS

How different are the carving turns in different alpine disciplines? The equations of carving turn (64–68) have only three dimensionless parameters which modify these equations and hence their solutions – the coefficient of dynamic friction μ , the slope angle α and the dynamic sidecut parameter $K = R_{sc}/L_g$. Out of the three only K depends on the ski sidecut radius. If we put $K = 0$ then the dependence of dimensionless solutions of the sidecut radius would vanish completely. This means that for the same slope angle and coefficient of friction (the same snow conditions and wax), the slalom and downhill carving runs would be just scale versions of each other, with the trajectory size scaling as $\propto R_{sc}$, skier's speed as $\propto \sqrt{R_{sc}}$ and both the g-force and the inclination angle being exactly the same.

In reality, $K \neq 0$ and it grows with the sidecut radius. However, it remains small ($K \ll 1$) even for DH skis. In the speed equation (64), K is a coefficient of the aerodynamic drag term Kv^2 . Hence, what differentiates carving runs of different alpine disciplines is the relative




FIG. 16. The asymptotic turn parameters for ideal carving runs on the $\alpha = 15^\circ$ slope with SL (solid), GS (dashed) and DH (dash-dotted) skis.

importance of the aerodynamic drag. Because the drag increases with K we expect the speed of the downhill run to be slower than that of the slalom run, when both are measured in the units of their own V_{sc} . The smaller speed implies the larger turn radius (in the units of R_{sc} , see Eq.68) and hence the lower g-force (Eqs.58 and 59) and inclination angle (Eq.39). To check this, we compared the runs on the same slope ($\alpha = 13^\circ$ and $\mu = 0.04$) with SL, GS (giant slalom) and DH skis. Figure 15 shows the dimensionless trajectories of these runs in the asymptotic regime. Although the trajectories are relatively similar they are not exact copies of one another, with larger K yielding longer and smoother turns. Figure 16 shows the evolution of the main run variables during the turns and confirms that the larger sidecut

radius yields turns with the smaller inclination angles and lower g-force.

We extended the study described in the previous section in order to determine the critical slope steepness α_c beyond which pure carving is impossible for SL, GS and DH skis. The result is

$$\alpha_c = \begin{cases} 17^\circ & \text{for } K = 0.0325 \text{ (13m) ,} \\ 19^\circ & \text{for } K = 0.0875 \text{ (30m) ,} \\ 21^\circ & \text{for } K = 0.1250 \text{ (50m) ,} \end{cases} \quad (70)$$

where in the brackets we show the dimensional sidecut radius corresponding to $L_g = 400\text{m}$.

VIII. THE EFFECT OF ANGULATION

In the derivation of the ideal carving equation (40) it is implicitly assumed the the whole skier body is aligned with the direction of the effective gravity and hence the angle Ψ between the ski “plane” and the slope is the same as the angle Φ between the effective gravity force and the normal to the slope. However, in general $\Theta = \Psi - \Phi \neq 0$ as skiers tend to bend (angulate) their body in the frontal plane (see figure 17)). This is normally done by moving hip and to some degree knees to the inside of the turn so that $\Theta > 0$ [7]. There at least two benefits of such angulation.

Firstly, it introduces a good safety margin against accidental side-slippage of the skis. Indeed, the case of $\Theta = 0$, which we have been focusing on so far, is only marginally stable against such a slippage because in this case the effective gravity has no component aligned with the platform that skis make in the snow. A small deviation from this equality towards the $\Theta < 0$ state, e.g. due to the skier’s hip unintentionally moving to the outside of the turn, will create such a component in the direction away from the slope. The snow reaction force will remain normal to the platform and hence unable to prevent ski from side-slipping [7].

If $\Theta > 0$ then the platform-aligned component of effective gravity exists as well but it points towards the slope and presses the skis against the inside wall of the platform, making the side-slippage impossible. The wall reacts with its own reaction force (force $F_{n,w}$ in figure 17) and as the result, the total \mathbf{F}_n is no longer normal to the platform and hence to the bottom surface of the ski. This allows the total normal reaction force to balance the effective gravity force even if the latter is not normal to the platform. Provided Ψ significantly exceeds

Φ , small variations of these angles due to various imperfections (perturbations) will not affect the outcome.

Secondly, it allows to vary the turn radius, thus introducing some control over the carving turn. Using the notation of Section IV

$$\tan \Psi = (\xi^2 - 1)^{1/2},$$

whereas

$$\tan \Phi = a\xi + b.$$

For $0 < \Phi < \Psi < 90^\circ$, we have

$$\tan \Psi = \eta \tan \Phi,$$

where $\eta > 1$. This yields the modified ideal carving equation

$$(\xi^2 - 1)^{1/2} = \eta(a\xi + b). \quad (71)$$

A careful (and a bit tedious) analysis of this equation shows that $d\xi/d\eta > 0$ provided $\eta a < 1$ and hence higher η means lower R . In other words, by increasing the angulation the skier can tighten the arc and the other way around, which is well known to skiing experts [e.g. 2].

Introducing $a' = \eta a > a$ and $b' = \eta b > b$, we can write equation (71) in exactly the same form as the original ICE (equation 40), with a replaced by a' and b by b' . This allows us to make a number of useful conclusions on the carving speed limits for positive angulation. Clearly, if $a + b > 0$ then $a' + b' > 0$ as well and hence the lower speed limit (48,49) remains unchanged. However, a sufficiently large η may yield $a' > 1$ even when $a \leq 1$ thus making the upper speed limit even more restrictive (see Section IV A).

In order to illustrate the effect of angulation, we consider the so-called 360° carving turn. By this we understand a carving turn which continues in the clockwise (or counter-clockwise) direction all the way, first downhill then uphill and finally downhill again. In this example, skier's angulation is a function $\Theta = \Theta(\Phi)$ defined via

$$\tan \Psi = A + \tan \Phi, \quad (72)$$

where A is constant. It is easy to see that $A = \tan \Theta(0)$. It is also easy to verify that $\Theta(\Phi)$ is a monotonically decreasing function which vanishes as $\Phi \rightarrow 90^\circ$, and that it yields $\Psi < 90^\circ$ for all $0 \leq \Phi < 90^\circ$.




FIG. 17. Inclination and angulation. Left panel: Because of the angulation at the hip, the inclination angle of the skis Ψ is higher than the inclination angle of the skier Φ . Right panel: Because $\Psi > \Phi$ the effective gravity has not only the component normal to the platform cut in the snow by the ski (F_{\perp}) but also the component parallel to the platform and pushing the ski into the platform wall (F_{\parallel}). The wall reacts with the force $F_{n,w}$, balancing F_{\parallel} .

Here we present the results for the slope angle $\alpha = 5^\circ$, the initial skier speed $v_{ini} = 0.5V_g$ and the initial angle of traverse $\beta_{ini} = 54^\circ$. Such a high speed could be gained at a steeper uphill section of the slope. Figure 19 shows the results for two runs: one with no angulation ($A = 0$, dashed lines) and one with strong angulation ($A = \tan(30^\circ)$, solid lines). The run without angulation falls a bit short of success. It stops on approach to the summit as the skier's speed drops to zero.

In the run with angulation, the turn is much tighter and the skier reaches the summit retaining a fair fraction of the initial speed. This allows them to continue and complete the 360° -turn. Because of the similar initial speed but the lower turn radius, the centrifugal force and hence the total g-force of this run are higher. This is also reflected in the higher inclination angles, with Φ peaking at $\beta \approx 130^\circ$ where it reaches $\approx 67^\circ$. Significant body angulation is physically demanding also because the skeleton is no longer well stacked and hence bears a smaller fraction of the total skier's weight. This results in a higher risk of injury.




FIG. 18. The trajectory of ideal carving run for the steep slope ($\alpha = 20^\circ$) and steered turn initiation with $\Delta\beta = 30^\circ$.

IX. THE EFFECT OF STEERING

There are two obvious limitations associated with pure carving. First, the carving turn radius cannot be changed at will as it is almost uniquely determined by the speed, traverse angle and slope gradient. If the race course dictates a smaller or larger turn radius some deviation from pure carving has to be introduced. Second, our experiments with ideal carving model show that on steep slopes this type of skiing leads to building up of excessive speed and eventual crash. To enable runs on such slopes skiers must have additional means of speed control.

These limitations can be overcome using the well-known technique of steering. In this technique, the skis are set at some angle relative to the direction of motion. We will denote this angle, called the steering angle, as $\Delta\beta$. In such a position, skis displace a certain amount of snow (snowploughing), or scrape the top layer of ice, which leads to braking. The exact degree of braking depends on how the steering is executed. In the traditional fully steered turns, the steering is continued all the way to the turn completion. In its modern variant, it is used mostly at the initiation phase of the turn. Once the speed is sufficiently reduced and/or the desired direction of the skis is reached, it is terminated and the rest of the turn




FIG. 19. The attempts at 360° turn with (solid lines) and without (dashed lines) angulation.

is continued as carving. In the so-called “power-slide” the return to carving is accompanied by forceful reduction of the steering angle via pivoting of skis.

In our modelling of such hybrid turns, we assume that at the turn initiation skis are rapidly pivoted to the desired steering angle and then they preserve their orientation relative to the fall line until carving is resumed. At the resumption point, the velocity component normal to the skis vanishes completely. Thus if β_{fin} and v_{fin} are the angle of traverse and the speed at the completion of the previous turn respectively, the carving phase of the next turn starts with

$$\beta_{ini} = (180^\circ - \beta_{fin}) + \Delta\beta \quad (73)$$




FIG. 20. The trajectory of ideal carving run for the steep slope ($\alpha = 20^\circ$) and steered turn initiation with $\Delta\beta = 45^\circ$.

and

$$v_{\text{ini}} = v_{\text{fin}} \cos(\Delta\beta). \quad (74)$$

Correspondingly, each time the skier's kinetic energy is reduced by the factor $\cos^2 \Delta\beta \approx 1 - (\Delta\beta)^2$ for small $\Delta\beta$. In the model we also ignore the finite duration of the steering phase and assume that it is negligibly short instead.

To illustrate the effects of steering we made a run for a steep slope with $\alpha = 20^\circ$ run and the steering angle $\Delta\beta = 45^\circ$. For this steering angle, exactly one half of skier's kinetic energy disappears at the transition between turns. This is just enough to avoid the singularity in the carving phase. The run is initiated with $x_{\text{ini}} = y_{\text{ini}} = 0$, $\beta_{\text{ini}} = 0.3\pi$ (54°), and $v_{\text{ini}} = (1/3)V_{\text{sc}} (=0.1V_g)$. Each turn terminates when the angle of traverse reaches $\beta_{\text{fin}} = 0.9\pi$ (162°). $\mu = 0.04$ and $K = 0.0325$.

Figure 20 shows the trajectory of the run and figure 21 the evolution of its key variables. One can see that the trajectory is not smooth any more as each transition between turns introduces a break. It would smooth out in a more accurate model where the transition is not instantaneous. As before, the solution quickly converges to a limiting one where each next turn is a copy of the previous one. The local radius of curvature, inclination angle and g-force still vary dramatically throughout the turn but remain at more-or-less realistic levels. Obviously, a higher steering angle would moderate them furthermore.




FIG. 21. The same as in figure 10 but for the steep slope ($\alpha = 20^\circ$) and steered turn initiation with $\Delta\beta = 45^\circ$.

X. LATERAL STABILITY

A. Stacked position

We first consider the stability of a stacked skier. Suppose that in the equilibrium position the inclinations angles of the ski (skier) and effective gravity are Ψ_0 and $\Phi_0 = \Psi_0$ respectively. Using the notation of Sec.IV,

$$\tan \Psi_0 = (\xi_0^2 - 1)^{1/2}, \quad \tan \Phi_0 = a\xi_0 + b, \quad (75)$$

where ξ_0 stands for the equilibrium turn radius. Consider a perturbation which changes the ski angulation position but keeps the skier velocity unchanged (and hence a and b as well). However, the turn radius changes and so does the effective gravity. We need to determine if the modified effective gravity is a restoring force or it pushes the system further away from the equilibrium. In the perturbed state $\Psi = \Psi_0 + \delta\Psi$, $\Phi = \Phi_0 + \delta\Phi$ and $\xi = \xi_0 + \delta\xi$, where δA stands for the perturbation of A . It is clear that when $\delta\Psi > 0$ the instability condition reads $\Psi > \Phi$ or $\delta\Psi > \delta\Phi$, whereas for $\delta\Psi < 0$ it is $\delta\Psi < \delta\Phi$. Both these cases are captured in the instability condition

$$\frac{\delta\Phi}{\delta\Psi} < 1. \quad (76)$$

Using equation (75) we find

$$\delta(\tan \Psi) = \frac{\xi_0}{\tan \Psi_0} \delta\xi, \quad (77)$$

and

$$\delta(\tan \Phi) = a\delta\xi. \quad (78)$$

Hence

$$\frac{\tan \delta\Phi}{\tan \delta\Psi} = \frac{a \tan \Psi_0}{\xi_0} = a \sin \Psi_0. \quad (79)$$

According to the condition (46), in carving turns $a < 1$ and hence equation (79) immediately yields $\tan \delta\Phi / \tan \delta\Psi < 1$. This implies $\delta\Phi / \delta\Psi < 1$ and therefore we conclude the lateral equilibrium is unstable.

Skiers have ways of controlling this instability. Lind and Sanders [5] discuss the stabilising arm moment, similar to what is used by trapeze artists. A skier can also use ski poles to provide an additional point of support. Finally and most importantly, when both skis are sufficiently wide apart and loaded the skier is more like a solid table than an inverted pendulum.

B. Angulated position

While the balance of an angulated skier is still unstable, in this case there is an additional way of controlling the instability, namely by a suitable change of the angulation. If $\delta\Psi$ is the perturbation of the ski inclination angle and $\delta\Phi$ is the corresponding perturbation of the effective gravity angle that the skier can restore their balance via changing their angulation

by the amount

$$\delta\psi = \delta\Psi - \delta\Phi. \quad (80)$$

XI. CONCLUSIONS

In this paper we described a simple mathematical model of carving turns in alpine skiing, which can be applied to snowboarding as well. The model combines a system of ordinary differential equations governing the CM motion with the algebraic Ideal Carving Equation (ICE), which emerges from the analysis of skier balance in their frontal plane. ICE relates the local radius of the CM trajectory to skier's speed and direction of motion relative to the fall line and hence provides closure to the system.

In the case of fall-line gliding, the skier speed grows until the gravity force is balanced by the aerodynamic drag and snow friction forces. Unless the slope is very flat, the snow friction is a minor factor and can be ignored, whereas the balance between the gravity and the aerodynamic drag yields the saturation speed V_g . Therefore this speed and the distance L_g required to reach it may be used as the characteristic scales of alpine skiing. However, ICE introduces another characteristic speed, the carving speed limit V_{sc} [3]. When the skier speed exceeds V_{sc} , carving becomes impossible in the whole Lower-C part of the turn. In fact, at the fall line the turn radius $R \rightarrow 0$ and the g-force diverges as $v \rightarrow V_{sc}$. Because as a rule $V_{sc} < V_g$, it is a more relevant speed scale in the dynamics of carving turns. Moreover, the radius of carving turns cannot exceed the ski sidecut radius R_{sc} , making the latter a natural length scale of this dynamics. These two scales yield the dimensionless equations of carving turn which have only three dimensionless control parameters: the slope gradient angle α , the dynamic coefficient of friction μ and the dynamic sidecut parameter $K = R_{sc}/L_g$, which determines the relative strength of the aerodynamic drag.

In addition to the upper limit, IDE also sets a lower speed limit, which must be satisfied to enable carving in the upper-C. Below this limit the skier can be balanced in the frontal plane only when the turn radius is above R_{sc} , which is not possible.

We used the model to explore ski runs composed of linked carving turns on an slope with constant gradient. While in reality such turns are linked via a transition phase of finite duration, in our current model the transition is instantaneous. At the transition, skier's speed and direction of motion remain invariant, whereas the inclination angle and hence the

turn radius jump to the values corresponding to the next turn. All these transitions occur as soon as the traverse angle (the angle between the skier trajectory and the fall line) reaches a particular fixed value. This often forces solutions towards a limit cycle, where the turns become indistinguishable one from another, unless the run terminates due to violation of the upper speed limit.

The results show the existence of a critical slope gradient above which the ideal carving runs become impossible. Its value depends on the value of K and hence on the sidecut radius of skis. For the typical value of L_b , we find it to be $\alpha_c \leq 17^\circ$ for SL skis, $\alpha_c \leq 19^\circ$ for GS skis and $\alpha_c \leq 21^\circ$ for DH skis. Slopes with smaller gradient can be roughly divided into the flat and moderately-steep groups. For flat slopes, the aerodynamic drag is not dominant over the snow friction even in the case of fall-line gliding. In our carving runs, the role of the snow friction is even more important and the turn speed saturates well below both V_g and V_{sc} . The carved arcs are nearly circular and their radius is only slightly below the sidecut radius of the skis. The skier inclination angle and the g-force stay relatively small.

On slopes of moderate gradient the saturation speed of fall-line gliding is very close to V_g and well above V_{sc} . However, the speed of carving runs still saturates well below V_g and close only to V_{sc} . The last condition makes the carving turns quite extreme. Their shape begins to deviate from the rounded shape of letter C and remind J instead, with the local turn radius significantly decreasing on the approach to the fall line. This results in increasing centrifugal force increases and hence the total effective gravity (g-force). The inclination angle becomes extremely high too. The high effective gravity leads to high normal reaction from the snow and significantly increased friction, which is the reason why the speed stays well below V_g .

On steep slopes with $\alpha > \alpha_c$ the skier's speed quickly exceeds V_{sc} after which the carving turn cannot be continued. In practice this means a loss of balance and crash.

When given in the units of R_{sc} and V_{sc} , the carving run solutions corresponding to $13\text{m} < R_{sc} < 50\text{m}$ are rather similar. In fact, R_{sc} enters the problem only via the dynamic sidecut parameter $K = R_{sc}/L_g$, which remains small for these radii. Moreover, K appears only in the speed equation where it defines the relative strength of the term describing the aerodynamic drag force. The fact that $K \ll 1$ ensures that the dynamical importance of the drag term is small and hence the solutions are not very sensitive to the variations of R_{sc} for as long as K remains small. Yet the drag term is not entirely negligible and the

solutions show some mild variation with R_{sc} within the studied range. In particular, turns corresponding to a larger sidecut radius are less extreme, with smaller inclination angles and weaker g-forces. This slight variation also explains the weak dependence of the critical gradient α_c on R_{sc} , as described earlier.

In the recent field study of the forces experienced by top GS skiers, it was found that on average the aerodynamic drag was about 7 times below the snow friction [10]. The authors attributed this result to the fact that the participating racers could not perform pure carving and skidded instead. This skidding could be understood as a sign of imperfect turn execution, implying that a better prepared skier could approach the aerodynamic speed limit V_g . Given the fact that this would involve the speed increase by more than twice, this seems highly unlikely. Our study provides an alternative explanation. First of all, on steep slopes the perfect carving is theoretically impossible. On such slopes some efficient measures of speed control, of which skidding seems to most available, must be employed to keep it below V_{sc} and hence well below V_g . Secondly, in carving runs on slopes of subcritical gradient, the snow friction outperforms the aerodynamic drag force. For example in our simulated carving run on a 15° slope (see Sec.VI B), the friction dominates the drag on average by the factor of 3.

Skiing on super-critical slopes must involve speed control via skidding. Watching runs made by the best athletes, including those competing on the World Cup circuit, one can notice that they do pure carving only on the flat sections of the race track whereas on the steep sections they perform hybrid turns which combine pivoting and skidding early in the turn with carving later on. In order to probe the potential of such technique, we made a simple modification of our model via instantaneous ski pivoting and corresponding speed reduction at the transition between carving turns. The solutions obtained with this crude model are in agreement with the observations. They confirm that such hybrid technique allows tskiing on slopes of super-critical gradient, and that skidding phase is crucially important.

Since the radius of a carving turn is not well controlled by the skier but is rather dictated by the slope and speed, ski pivoting is also helpful when the skier has to execute a tighter turn than the one possible with pure carving.

Some degree of control over the radius of carving turns can also be reclaimed via the skier angulation in the frontal plane. We derived a slightly modified version of ICE which takes the angulation into account and analysed its implications. First, the angulation does

not allow to remove the carving speed limit V_{sc} . On the contrary, it becomes even stricter. Secondly, for the same speed and slope, the turn radius is smaller for higher angulation. This shows how the angulation can be used to control the turn radius. In particular, we described a little experiment which demonstrated how the angulation helps to execute the full 360° carving turn. The angulation also helps to prevent side slipping of skis, reducing the importance of ski boot canting. The shorter radius also makes the g-force higher and hence the skiing more physically demanding. This is particularly unwelcome as the angulation reduces the ability of the body's skeletal structure to bear most of skier's weight and puts more stress of the body's muscular system.

We have also studied the stability of the lateral balance set by the original ICE, where the skier is balancing on a single ski. Although we find that it is unstable we do not think this has important implications. In particular, skiers can control the instability using their angulation. Moreover usually the skier weight is distributed over both skis which is obviously a stabilising factor as well.

There is no hiding from the fact that our model is rather simplified and leaves plenty of scope for improvement. One can think of a more realistic model for the transition between pure carving turns than the instantaneous jump of the inclination angle. The role of angulation can be explored more thoroughly and the way it is used in various phases of the turn can be based more closely on the runs performed by actual athletes. The shape of modern skies allows to slightly vary the effective sidecut radius by preferential loading of either the tip or the tail sections of the skis. The hybrid turn requires more realistic treatment of the skidding phase. Real slopes vary in steepness and skiers actively move their CM up and down to handle the terrain etc. All these factors can be taken into account and we expect much more realistic models of skiing, including ski racing, to emerge in foreseeable future.

ACKNOWLEDGMENTS

In order to solve the equations of ideal carving, we used the software package *Maple* (Maple is a trademark of Waterloo Maple Inc.).

- [2] Harb H.R., Essentials of Skiing (Hatherleigh Press, New York, 2006).
- [3] Jentschura, U.D., Fahrbach, Can. J. Phys., **82**, 249 (2004).
- [4] Landau, L.D., Lifshitz E.M., Mechanics, (Pergamon Press, Oxford, 1969)
- [5] Lind D., Sanders, S.P., The physics of skiing: Skiing at the triple point (Springer-Verlag, New York, 1996).
- [6] Nordt, A.A., Springer, G.S. Kollár, L.P., Sports Engineering, **2**, 181 (1999)
- [7] LeMaster R., Ultimate Skiing (Human Kinetics, Champaign, 2010).
- [8] Supej M., J. Appl. Biomech., **24**, 121(2008).
- [9] Supej, M., Kipp, R., Holmberg, H. C., Scand. J. Med. Sci. Sports, **21**, 72 (2011).
- [10] Supej, M. et al., Scand. J. Med. Sci. Sports, **23**, 38 (2013).
- [11] <http://www.speed-ski.com/Schedule/Event/2018/World-Cup-Race11-Grandvalira>.
- [12] Tada N., Hirano Y., Sports Engineering, **5**, 15 (2002)