LATENT COVERAGE FROM NOISY CALIBRATION

Kun Woo Park

Kookmin University

Abstract: Prediction intervals for a latent quantity, such as the true mean of an unsampled small area, are often calibrated on noisy proxies with the usual conformal threshold. We ask what latent coverage such calibration guarantees when the law and the variance of the noise are unknown and differ between units. Unimodality of the latent law gains nothing over no shape assumption. Under bi-log-concavity we give the noisy level that guarantees a latent coverage target, exactly for Gaussian and symmetric unimodal noise. Under Gaussian noise at a target of ninety percent the usual threshold is short by less than one thousandth, and for small noise only residual densities that increase outward at an end of the interval can make it short. A small raise of the conformal rank, which needs no noise variances, restores a high-probability guarantee for every noise configuration in a class, and in the worst case no smaller raise does. When a share of the latent law departs from bi-log-concavity, the exact level grows with that share, which prices a doubtful shape assumption. Simulations and a school-district illustration accompany the theory.

Key words and phrases: Bi-log-concavity, conformal prediction, measurement error, small area estimation, tolerance interval.

1. Introduction

Suppose that a residual W, the latent target minus a centre fitted on independent training data, is observed only through $V = W + e$, with noise e independent of W. A survey statistician who calibrates a prediction interval for the true mean of an unsampled area on the direct estimates of sampled areas is in this position (Fay and Herriot, 1979; Rao and Molina, 2015), as is a conformal analyst whose calibration responses carry measurement error (Einbinder et al., 2024; Cohen et al., 2025). The threshold t at which the noisy scores $|V|$ have coverage p is observable; the quantity of interest is the latent coverage $\mathrm{pr}(|W| \leq t)$. The noise law and its variance are often unknown, or known only through rough estimates, and they differ between units: direct estimates of areas with different sample sizes have different variances.

The interval is for a unit without its own noisy observation, such as an unsampled area, predicted synthetically; intervals for sampled areas that combine a direct estimate with a model are a different problem. The practice is to calibrate on the noisy scores as if they were latent, with the usual conformal rank. We ask when this covers the latent target, and what is the smallest correction that makes it do so when the noise laws and scales are unknown and differ between units.

Two answers are known. If W is symmetric and unimodal about zero, no interval of length $2t$ carries more of its mass than $[-t, t]$, so $\mathrm{pr}(|W + e| \le t) \le \mathrm{pr}(|W| \le t)$ for every independent noise and the noisy threshold is conservative; this is the one-dimensional case of the theorem of Anderson (1955), and Einbinder et al. (2024, Proposition 1) prove a version for conformal intervals. Without a shape assumption, median-zero noise turns a noisy level $1 - \alpha$ into a latent level $1 - 2\alpha$ (Guille-Escuret and Ndiaye, 2024, Proposition 2.3). Symmetry is a strong assumption, since residuals of area means of skewed outcomes are asymmetric, and the shape-free answer is very costly.

We take the latent law to be bi-log-concave (Dümbgen, Kolesnyk, and Wilke, 2017): its distribution function F is continuous and both $\log F$ and $\log(1 - F)$ are concave. The class contains the log-concave laws and some multimodal ones and does not require symmetry. The results determine how close noisy calibration is to valid without symmetry, what exact validity requires, and how fast the guarantee weakens when the shape assumption fails.

1. *A sharp transfer.* Unimodality of the latent law, even with mean zero, gives nothing over no shape assumption (Proposition 1). Under bi-log-concavity we give the noisy level $\Psi_{\mathcal{C}}(q)$ that guarantees latent coverage

q, exactly for Gaussian and symmetric unimodal noise (Theorems 1 and 2). For Gaussian noise at $q = 0.9$ it exceeds q by 8.0×10^{-4}: noisy coverage 0.9 guarantees latent coverage 0.89918 and no more.

2. *The price of a doubtful shape assumption.* If a share ξ of the latent law is arbitrary, the exact level is $(1-\xi)\Psi_{\mathcal{C}}\{q/(1-\xi)\} + \xi/2$ (Proposition 5). At $q = 0.9$ under Gaussian noise it is 0.90081, 0.90322, 0.90564 and 0.92517 for $\xi = 0$, 0.5%, 1% and 5%: each percent of the latent law outside the class raises the level by about 0.005, five times the cost, by a sufficient bound, of a percent of noise outside its class.

3. *A variance-free rank, and when the usual one suffices.* Calibrating at these levels gives a conformal rank that keeps the latent guarantee with probability at least $1 - \delta$ for every noise law and scale in the class, possibly different between units (Theorem 3); in the worst case no smaller rank is valid, and at $K = 10^3$ the raise is one or two ranks for Gaussian and symmetric unimodal noise without contamination, and 24 ranks at $\xi = 5\%$. For small noise the usual rank can lose its guarantee only if $f'(r) > f'(-r)$, so that the residual density increases outward at an end of the interval, as when a biased centre puts the mode beyond the threshold (Propositions 3 and 4, § 5.1).

Related methods target different quantities or use more information (Table 1). LatentCP (Zheng, Zhou, and Zhu, 2026) gives a marginal latent guarantee through a known forward model; Cohen, Goldberger, and Tirer (2025) and deconvolution with known heteroscedastic or unknown symmetric errors (Delaigle and Meister, 2008; Delaigle and Hall, 2016) estimate the latent law or threshold without a coverage guarantee; Ignatiadis and Wager (2022) bound functionals of a nonparametric latent law with high probability when the noise law is known and common to all units, and Soloff, Guntuboyina, and Sen (2025) estimate it under heteroscedastic Gaussian noise. Bounds for noisy labels, contamination or distribution shift (Sesia, Wang, and Tong, 2025; Penso, Goldberger, and Fetaya, 2025; Clarkson, Xu, Cucuringu, and Reinert, 2024; Barber, Candès, Ramdas, and Tibshirani, 2023; Cauchois, Gupta, Ali, and Duchi, 2024; Feldman, Bates, and Romano, 2026) control marginal coverage up to a term that does not shrink with the sample, whereas here the high-probability guarantee is at stake. Our rank is the tolerance limit of Wilks (1941), the training-conditional rank of Vovk (2012) and Bian and Barber (2023), with a raised level, and Guille-Escuret and Ndiaye (2024, Appendix C.3) already carry their shape-free bound to a high-probability guarantee; what is new is the sharp level under a shape assumption, its price when the assumption fails, and when the raise is

needed. Conformal intervals for small areas, survey or hierarchical data cover observable responses or conditional expectations (Bersson and Hoff, 2024; Wieczorek, 2023; Dunn et al., 2023; Zhang and Tuoto, 2026). Intervals for latent small-area means (Yu and Hoff, 2018; Burris and Hoff, 2020; Diao et al., 2014; Yoshimori and Lahiri, 2014; Hall and Maiti, 2006; Chatterjee et al., 2008; Armstrong et al., 2022) use sampling variances, known or estimated with their uncertainty taken into account (Wang and Fuller, 2003; You and Chapman, 2006; Dass et al., 2012), and a model for the latent effects; tolerance intervals for true values (Wang and Iyer, 1994) and for random-effects models with heterogeneous error variances (Liao et al., 2005; Lin et al., 2008) rely on normality. The rank studied here needs neither variance estimates nor a parametric latent model.

Section 2 sets up the problem, Section 3 gives the coverage transfer and Section 4 the calibration rank, when the usual rank is valid, and its robustness. Sections 5 and 6 give numerical evidence and an application. Proofs and further results are in the Supplementary Material.

2. Setting

Let $\mathcal{L}$ be the class of log-concave laws on the real line, including point masses, and $\mathcal{B} \supset \mathcal{L}$ the class of laws that are point masses or have a con-

Table 1: Interval methods for a latent target calibrated on noisy scores, by the noise information used and the guarantee targeted

method	noise information	guarantee
noisy threshold, usual rank	none	none in general; nearly latent under bi-log-concavity (§ 4.2)
noisy threshold, symmetric case (Anderson, 1955; Einbinder et al., 2024)	none (independent noise)	latent, symmetric unimodal latent law
shape-free level (Guille-Escuret and Ndiaye, 2024)	median-zero noise	latent, any latent law
LatentCP (Zheng, Zhou, and Zhu, 2026)	known forward model	marginal latent
deconvolution threshold (Cohen, Goldberger, and Tirer, 2025)	known common noise kernel	none in finite samples
F-localization (Ignatiadis and Wager, 2022)	known common noise law	latent with probability $1-\delta$, nonparametric latent law
this paper, Theorem 3	a class of noise laws only	latent with probability $1-\delta$, bi-log-concave latent law

tinuous distribution function F with $\log F$ and $\log(1-F)$ concave. Both classes are closed under translation and scaling, and convolution with a log-concave law preserves them (Supplementary § S1); $\mathcal{B}$ contains some bimodal mixtures of normal laws (Supplementary Lemma S1).

A class $\mathcal{C}$ of noise laws is closed under scaling and negation. The classes considered are the Gaussian laws $\mathcal{N} = \{\sigma Z : \sigma > 0\}$; symmetric unimodal laws; mean-zero log-concave laws; mean-zero unimodal laws; and, for rules without a latent shape assumption, median-zero laws. For a noise law and a latent law, the noisy coverage of $[-t, t]$ is $\mathrm{pr}(|W + e| \le t)$ and the latent coverage is $\mathrm{pr}(|W| \le t)$.

In calibration, conditionally on the training data, $W_1, \ldots, W_K, W_{\mathrm{new}}$

are independent latent residuals and $e_i = \sigma_i \epsilon_i$ are independent noises, independent of the residuals, with laws of ϵ_i in $\mathcal{C}$ that may differ between units and fixed unknown scales $\sigma_i > 0$. Let $\mathcal{D}$ denote the calibration data and $T = |V|_{(k)}$, the kth smallest of $|V_1|, \ldots, |V_K|$. The interval $[-T, T]$ is placed around the fitted centre of a new unit. We call

$$\mathrm{pr}_{\mathcal{D}}\{\mathrm{pr}(|W_{\mathrm{new}}| \leq T \mid \mathcal{D}) \geq q\}$$

the latent reliability of rank k: the probability, over calibration samples, that the interval attains latent coverage q. A rank is valid at level $1-\delta$ if its latent reliability is at least $1-\delta$ for every latent law and noise configuration in the model. The usual rank is the smallest k with $\mathrm{pr}\{\mathrm{Bin}(K, q) \leq k-1\} \geq 1-\delta$ (Wilks, 1941; Vovk, 2012), valid when the scores are noise free. Q_q denotes the lower q-quantile, Φ and ϕ the standard normal distribution function and density.

3. Coverage transfer

3.1 Unimodality does not suffice

Let $\beta_{\mathcal{C}} = \inf_{\epsilon \in \mathcal{C}} \mathrm{pr}(\epsilon \geq 0)$: $1/2$ for Gaussian, symmetric unimodal and median-zero noise, e^{-1} for mean-zero log-concave noise, attained by the centred exponential law, and 0 for mean-zero unimodal noise.

Proposition 1. *Let* $\beta_{\mathcal{C}} > 0$ *and* $1 - \beta_{\mathcal{C}} < p < 1$. *Then* $\mathrm{pr}(|W| \le 1) \ge 1-(1-p)/\beta_{\mathcal{C}}$ *for every* W *and every noise with law in* $\mathcal{C}$ *and* $\mathrm{pr}(|W+e| \le 1) \ge p$. *If the infimum defining* $\beta_{\mathcal{C}}$ *is approached by laws continuous at* 0, *then for every* $\gamma > 0$ *there are a unimodal* W *with* $E(W) = 0$ *and such a noise with* $\mathrm{pr}(|W| \le 1) \le 1 - (1-p)/\beta_{\mathcal{C}} + \gamma$. *If* $\beta_{\mathcal{C}} = 0$, *approached in the same way, then for every* $p < 1$ *and* $\gamma > 0$ *there are such* W *and noise with* $\mathrm{pr}(|W| \le 1) \le \gamma$.

The lower bound is that of Guille-Escuret and Ndiaye (2024, Proposition 2.3); Proposition 1 adds that unimodal latent laws with mean zero attain it. With median-zero noise, noisy coverage 0.9 thus guarantees only latent coverage 0.8. The extremal laws place a thin layer of mass just outside the threshold, and the noise carries a fraction $1 - \beta_{\mathcal{C}}$ of it inside. Bi-log-concavity excludes such a layer: concavity of $\log F$ makes f/F non-increasing, so the density cannot jump up at the threshold.

3.2 One end of the interval

For a class $\mathcal{C}$ let

$$\Psi_{\mathcal{C}}(q) = \sup_{\epsilon \in \mathcal{C}} E\{\min(1, q\mathrm{e}^{-\epsilon})\}. \tag{3.1}$$

The extremal latent laws are exponential tails, because concavity of $\log F_Y$ limits how fast mass can accumulate below a quantile: with X_u exponential

with rate $u > 0$, the law of $b - X_u$ has distribution function $\min\{1, \mathrm{e}^{u(y-b)}\}$.

Proposition 2. *Let ϵ have a law in $\mathcal{C}$ and be independent of Y, with $\log F_Y$ concave. If $\mathrm{pr}(Y + \epsilon \le 0) \ge p > \Psi_{\mathcal{C}}(q)$, then $F_Y(0) \ge q$. If $p < \Psi_{\mathcal{C}}(q)$, some exponential tail Y with $F_Y(0) < q$ and some ϵ with law in $\mathcal{C}$ have $\mathrm{pr}(Y + \epsilon \le 0) > p$.*

Proof. Suppose that $v = F_Y^{-1}(q) > 0$, let $a = \log(1/q)$ and let u be a supergradient of $\log F_Y$ at v. Then $F_Y(y) \le \min\{1, q\mathrm{e}^{u(y-v)}\}$, so the exponential tail $v + a/u - X_u$ lies stochastically below Y. Adding ϵ preserves this, and as $v > 0$ and $u\epsilon$ has a law in $\mathcal{C}$,

$$p \le \mathrm{pr}(Y + \epsilon \le 0) \le \mathrm{pr}(a/u - X_u + \epsilon \le 0) = E\{\min(1, q\mathrm{e}^{-u\epsilon})\} \le \Psi_{\mathcal{C}}(q).$$

This uses $F_Y(v) = q$. Since a concave function is continuous on the interior of its domain, the only other case is an atom of mass at least q at the left end v of the support of Y; then $Y + \epsilon \le 0$ requires $\epsilon < 0$, and $p \le \mathrm{pr}(\epsilon < 0) \le \lim_{u\to\infty} E\{\min(1, q\mathrm{e}^{-u\epsilon})\} \le \Psi_{\mathcal{C}}(q)$. Conversely, $a - X_1$ with noise ϵ attains $E\{\min(1, q\mathrm{e}^{-\epsilon})\}$; shift it slightly to the right. □

Applied to $Y = (W - t)/s$ for any scale $s > 0$, Proposition 2 controls the right end of $[-t, t]$; applied to $-W$, it controls the left end. For a one-sided interval $(-\infty, t]$ it is the complete answer under concavity of $\log F_W$ alone.

3.3 Both ends

A noisy failure probability $1-p$ may be split in any way between the two ends, and each end transfers separately. Let $\lambda_{\mathcal{C}}(p) = \sup\{q' \in (0,1) : \Psi_{\mathcal{C}}(q') < p\}$, with $\sup\emptyset = 0$, and $\vartheta_{\mathcal{C}}(a) = 1 - \lambda_{\mathcal{C}}(1-a)$, the largest latent failure probability at one end compatible with noisy failure probability a there; $\vartheta_{\mathcal{C}}(0) = 0$ when $\beta_{\mathcal{C}} > 0$. Define the sufficient noisy level

$$p_{\mathcal{C}}(q) = \inf\Big\{p : \sup_{a_L,a_R\geq 0,\ a_L+a_R\leq 1-p} \{\vartheta_{\mathcal{C}}(a_L) + \vartheta_{\mathcal{C}}(a_R)\} \leq 1-q\Big\}. \tag{3.2}$$

Noisy coverage above $p_{\mathcal{C}}(q)$ at threshold t gives latent coverage q for every $W \in \mathcal{B}$ and every noise with law in $\mathcal{C}$ (Supplementary Lemma S2). Putting all the failure at one end shows that $\Psi_{\mathcal{C}}(q) \leq p_{\mathcal{C}}(q)$, and if $\Psi_{\mathcal{C}}$ is convex the worst split puts the whole failure at one end, so that $p_{\mathcal{C}}(q) = \Psi_{\mathcal{C}}(q)$ (Supplementary Lemma S3).

Theorem 1. *Let $q > \mathrm{e}^{-1}$.*

(i) For symmetric unimodal noise, $\Psi_{\mathcal{C}}(q) = (1+q\mathrm{e}^{-r})/2$, where $r > 0$ solves $(1+r)\mathrm{e}^{-r} = (1+\log q)/q$, attained by uniform noise; $\Psi_{\mathcal{C}}$ is strictly convex, so $p_{\mathcal{C}}(q) = \Psi_{\mathcal{C}}(q)$.

(ii) (Computer-assisted.) For mean-zero log-concave noise, $\Psi_{\mathcal{C}}(q)$ is the supremum over centred log-affine laws on segments; at $q = 0.9$ it lies in $(0.90517, 0.9052)$, the lower end attained by the centred exponential law,

and certified splits of the failure probability give $p_{\mathcal{C}}(0.9) \leq 0.906$.

(iii) For mean-zero unimodal noise, $\Psi_{\mathcal{C}}(q) = 1$.

Part (i) holds because symmetric unimodal laws are scale mixtures of uniform laws (Supplementary Proposition S1). For part (ii), the localization theorem of Fradelizi and Guédon (2004) reduces the supremum to log-affine laws on segments, which a branch and bound in ball arithmetic bounds (Supplementary §§ S2.4 and S5). For part (iii), a noise spread uniformly over $[-a, 0]$, balanced to mean zero by a small mass far above zero, makes $E\{\min(1, q\mathrm{e}^{-\epsilon})\}$ arbitrarily close to 1.

3.4 Gaussian noise

For the Gaussian class $\mathcal{N}$ the supremum in (3.1) is over the scale only. With $a = \log(1/q)$,

$$\Psi_{\mathcal{N}}(q) = \sup_{s>0} \Gamma(q, s), \qquad \Gamma(q, s) = \Phi(-a/s) + q\mathrm{e}^{s^2/2}\Phi(a/s - s). \tag{3.3}$$

Let $m(b) = \phi(b)/\Phi(b)$ and $A(b) = m(b)\{b + m(b)\} = 1 - \mathrm{var}(Z \mid Z \leq b)$, which decreases strictly from 1 to 0 as b runs over the real line (Sampford, 1953).

Theorem 2. *Consider the Gaussian class* $\mathcal{N}$.

(i) If $q \leq \mathrm{e}^{-1}$, *then* $\Psi_{\mathcal{N}}(q) = 1/2$, *approached as* $s \to \infty$ *and not attained.*

(ii) If $q > \mathrm{e}^{-1}$, then $\Gamma(q, \cdot)$ has a unique maximiser $s^* = m(b^*)$, where b^* is the unique solution of $A(b^*) = \log(1/q)$, and

$$\Psi_{\mathcal{N}}(q) = \Phi(-c) + \phi(c)/m(b^*), \qquad c = b^* + m(b^*).$$

(iii) $\Psi_{\mathcal{N}}$ is continuously differentiable, with $\Psi'_{\mathcal{N}}(q) = \mathrm{e}^{m(b^*)^2/2}\,\Phi(b^*)$ for $q > \mathrm{e}^{-1}$, and strictly convex on $(\mathrm{e}^{-1}, 1)$. Hence $p_{\mathcal{N}}(q) = \Psi_{\mathcal{N}}(q)$ for every $q \in (\mathrm{e}^{-1}, 1)$.

Outline of the proof. Let $b = a/s - s$. The identity $q\mathrm{e}^{s^2/2}\phi(b) = \phi(a/s)$ gives $\partial_s\Gamma = \phi(a/s)\{s/m(b) - 1\}$, so the sign of $\partial_s\Gamma$ is that of $s - m(b)$, and $s > m(b)$ if and only if $a > A(b)$. Since A increases through $(0, 1)$ as b decreases, $\Gamma(q, \cdot)$ increases throughout when $a \geq 1$, which gives (i), and has a unique interior maximum at $A(b^*) = a$ when $a < 1$, which gives (ii). By the envelope theorem $\Psi'_{\mathcal{N}}(q) = D(b^*)$ with $D(b) = \mathrm{e}^{m(b)^2/2}\Phi(b)$; b^* increases with q and $\{\log D(b)\}' = m(b)\mathrm{var}(Z \mid Z \leq b) > 0$, so $\Psi'_{\mathcal{N}}$ increases strictly. Details are in Supplementary § S2.6. □

At $q = 0.8$, 0.9 and 0.95, $\Psi_{\mathcal{N}}(q) = 0.8044106$, 0.9008035 and 0.9501590 to seven decimals; noisy coverage 0.9 guarantees latent coverage 0.89918 and no more. For Gaussian and symmetric unimodal noise the transfer is nontrivial exactly when $q > \mathrm{e}^{-1}$; for mean-zero log-concave noise $\Psi_{\mathcal{C}}(q) \geq 1 - \mathrm{e}^{-1}$ for every q, from scaled centred exponential noise. Table 2 collects

the levels.

Table 2: Levels and ranks at $q = 0.9$ and $\delta = 0.05$: what is assumed, the latent coverage guaranteed by noisy coverage 0.9, the noisy level that guarantees latent coverage 0.9, the smallest valid rank, and the result that gives it

assumption	latent cov.	level	$K = 110$	$K = 10^3$	$K = 10^4$	result
symm. unimodal residuals, any noise	≥ 0.9	0.9	105	916	9050	Anderson
$\mathcal{B}$, Gaussian noise	0.8991	0.90081	105	917	9058	Thm 3
$\mathcal{B}$, symm. unimodal noise	0.8981	0.90179	105	918	9068	Thm 3
$\mathcal{B}$, mean-zero log-conc. noise	≥ 0.8935	$(0.90517, 0.906]$	105	921–922	9101–9109	Thm 3
$\mathcal{B}$, mean-zero unimodal noise	none	1	–	–	–	Thm 1
$\mathcal{B}_\xi$, $\xi = 1\%$, Gaussian noise	0.8942	0.90564	105	922	9105	Prop. 5
$\mathcal{B}_\xi$, $\xi = 5\%$, Gaussian noise	0.8745	0.92517	107	940	9296	Prop. 5
any residuals, median-zero noise	0.8	0.95	109	962	9537	Prop. 1
any residuals, mean-zero log-conc. noise	$1 - \mathrm{e}/10$	$1 - 0.1/\mathrm{e}$	110	974	9664	Prop. 1
any residuals, $\mathrm{pr}(e \geq 0), \mathrm{pr}(e \leq 0) \geq 0.3$	0.6666	0.97	110	980	9729	Prop. 1

Levels are rounded up and latent coverages down, so both are safe to use; the Gaussian and symmetric unimodal levels are enclosed in ball arithmetic. For log-concave noise the levels run from a certified lower bound on $\Psi_{\mathcal{C}}(0.9)$ to a certified sufficient level, certified only at $q = 0.9$, and of a range of ranks the lower is necessary and the upper sufficient. Proposition 5 needs $q \geq 0.53$. For strongly skewed or discrete noise outside the log-concave class only the last row applies, with 0.3 replaced by a lower bound on the probabilities that the noise is nonnegative and nonpositive (§ 5.1). Noise whose scale depends on the latent value is covered only through the conditional ordering of § 4.3.

4. Calibration with unknown, heterogeneous noise

4.1 A valid rank

Theorem 3. *In the setting of § 2, let the latent residuals have a common law $G \in \mathcal{B}$. Then*

$$\mathrm{pr}_{\mathcal{D}}\{\mathrm{pr}(|W_{\mathrm{new}}| \leq T \mid \mathcal{D}) \geq q\} \geq \mathrm{pr}\{\mathrm{Bin}(K, p_{\mathcal{C}}(q)) \leq k - 1\}.$$

Conversely, if $\mathrm{pr}\{\mathrm{Bin}(K, \Psi_{\mathcal{C}}(q)) \le k-1\} < 1-\delta$*, there are a log-concave* G *and a noise law in* $\mathcal{C}$*, the same in every unit, for which the latent reliability is below* $1-\delta$*. Hence the smallest rank valid for every* $G \in \mathcal{B}$ *lies between the smallest* k *with* $\mathrm{pr}\{\mathrm{Bin}(K, \ell) \le k-1\} \ge 1-\delta$ *at* $\ell = \Psi_{\mathcal{C}}(q)$ *and at* $\ell = p_{\mathcal{C}}(q)$*, which coincide for Gaussian and symmetric unimodal noise.*

Proof. Let $r = Q_q(|W|)$ for $W \sim G$ and $N = \#\{i : |V_i| < r\}$. Then

$$\mathrm{pr}(|W_{\mathrm{new}}| \le T \mid \mathcal{D}) \ge q \iff T \ge r \iff N \le k-1.$$

For $s < r$, $\mathrm{pr}(|W| \le s) < q$, so by the transfer of § 3 at threshold s, $\mathrm{pr}(|V_i| \le s) \le p$ for every $p > p_{\mathcal{C}}(q)$, whatever the law and scale of e_i; letting $s \uparrow r$, $\mathrm{pr}(|V_i| < r) \le p_{\mathcal{C}}(q)$. N is a sum of independent indicators with these success probabilities, so it is stochastically smaller than $\mathrm{Bin}\{K, p_{\mathcal{C}}(q)\}$. Conversely, for $p < \Psi_{\mathcal{C}}(q)$ the one-ended laws of § 3 give a log-concave G and a noise with $\mathrm{pr}(|V_i| < r) > p$, so the latent reliability is at most $\mathrm{pr}\{\mathrm{Bin}(K, p) \le k-1\}$, which tends to $\mathrm{pr}\{\mathrm{Bin}(K, \Psi_{\mathcal{C}}(q)) \le k-1\}$ as $p \uparrow \Psi_{\mathcal{C}}(q)$. □

The bound holds unit by unit, so the noise laws and scales need not be known or shared; it uses independence of the units, not only exchangeability. For Gaussian and symmetric unimodal noise the rank is sharp: along the laws of the converse, the latent reliability of the smallest valid rank tends

to exactly $1-\delta$ as $K \to \infty$. These laws are degenerate, an exponential tail whose scale tends to zero relative to the threshold, with noise of the same order, so sharpness is an infimum that is approached but not attained; with a known lower bound on the noise relative to the threshold the needed level can only be lower, a question taken up for Gaussian noise in Supplementary § S4. The procedure needs only K, q, δ and a noise class.

1. Take the level of the noise class from Table 2, whose levels are rounded up and therefore sufficient, or from the accompanying software, which returns the upper end of a certified enclosure, for other $q > \mathrm{e}^{-1}$ (Gaussian, symmetric unimodal); a level rounded to the nearest value may be too low and lower the rank by one. For mean-zero log-concave noise a sufficient level is certified only at $q = 0.9$.

2. Take k as the smallest rank with $\mathrm{pr}\{\mathrm{Bin}(K, \mathrm{level}) \leq k-1\} \geq 1-\delta$; if even $k = K$ fails, no valid interval exists at this K.

3. Report $[\hat{\mu}_{\mathrm{new}} - T, \hat{\mu}_{\mathrm{new}} + T]$ with $T = |V|_{(k)}$.

With known positive scales s_i, fixed before calibration, the normalized scores $|V_i|/s_i$ and the interval $[-s_{\mathrm{new}}T, s_{\mathrm{new}}T]$ have the same guarantee if the normalized residuals W_i/s_i and $W_{\mathrm{new}}/s_{\mathrm{new}}$ have a common law in $\mathcal{B}$, or more generally under the ordering of Corollary 1 for the normalized

residuals, because both classes are closed under scaling; bi-log-concavity of each W_i/s_i alone is not enough, since scales that do not match the residuals rescale the target. At $K = 110$ the three bi-log-concave rows of Table 2 give the usual rank, $k = 105$, against 109 and 110 without a shape assumption.

Table 2 summarizes the choice of level. The class levels cost one to six ranks at $K = 10^3$, the levels without a latent shape assumption about fifty. Skewed or discrete noise, as for means of a few units of a skewed population, calls for the log-concave level and, if even log-concavity of the noise is doubtful, for a level without a shape assumption on the noise (§ 5.1).

4.2 When the usual rank is valid

The rank of Theorem 3 exceeds the usual rank because the level $\Psi_{\mathcal{C}}(q)$ exceeds q by a fixed amount, while consecutive ranks differ in level by about $1/K$. At the extremal laws this raise cannot be dropped.

Proposition 3. *Let $q \in (0,1)$. There are a log-concave law G and $D > 0$ such that, with $e_i \sim N(0, D)$ for all i, $\mathrm{pr}_{\mathcal{D}}\{\mathrm{pr}(|W_{\mathrm{new}}| \le |V|_{(k_K)} \mid \mathcal{D}) \ge q\} \to 0$ as $K \to \infty$ for every sequence of ranks with $k_K/K \to q$, such as the marginal and, for fixed δ, the usual high-probability rank.*

Proof. By Theorem 2, $\Psi_{\mathcal{N}}(q) > q$ for every $q \in (0,1)$. Take p with $q < p < \Psi_{\mathcal{N}}(q)$. The converse construction in the proof of Theorem 3 gives a

log-concave latent law and $D > 0$ with $H = \mathrm{pr}(|V_i| < r) > p$, $r = Q_q(|W|)$; the latent reliability of rank k is $\mathrm{pr}\{\mathrm{Bin}(K, H) \leq k - 1\}$, which tends to zero because $k_K/K \to q < H$. □

The latent coverage itself barely moves: at the extremal Gaussian-noise law, the usual rank's latent coverage tends to 0.89918 at $q = 0.9$. What fails is the guarantee, once K is of order $q(1-q)/\{\Psi_{\mathcal{N}}(q) - q\}^2$: the reliability falls below 0.5 at $K \approx 2.2 \times 10^4$ and 3.8×10^5 for $q = 0.8$ and 0.9 (§ 5.1). The extremal law, however, is far from a typical residual. It is an exponential tail $W = t + s\{\log(1/q) - X_1\}$ with s small: almost all of its mass lies just inside the threshold, on one side, so that its mean is close to t. A residual with a correctly fitted centre does not look like this.

How far from typical the extremal law must be is measured by a dial on centring. For $\zeta \geq 0$ let

$$p_{\mathcal{C}}^{\zeta}(q) = \sup\{\mathrm{pr}(|W + e| < 1) : W \in \mathcal{L},\ |E(W)| \leq \zeta,\ \mathrm{pr}(|W| < 1) \leq q\},$$

the supremum being also over noise laws in $\mathcal{C}$ and their scales. With the threshold scaled to 1, ζ bounds the bias of the fitted centre relative to the half-width of the interval; the extremal law above has ζ close to 1.

Proposition 4. *Let $\mathcal{C}$ be the Gaussian or the symmetric unimodal class, $\mathrm{e}^{-1} < q < 1$ and $\zeta \geq 0$.*

(i) If the latent residuals have a common log-concave law with $|E(W)| \leq \zeta\, Q_q(|W|)$*, the bound of Theorem 3 holds with* $p_{\mathcal{C}}^{\zeta}(q)$ *in place of* $p_{\mathcal{C}}(q)$*, for noise laws and scales in the class that differ between units.*

(ii) $p_{\mathcal{C}}^{\zeta}(q) = \Psi_{\mathcal{C}}(q)$ *for* $\zeta \geq 1$*, and* $q < p_{\mathcal{C}}^{\zeta}(q) \leq \Psi_{\mathcal{C}}(q)$ *for every* $\zeta \geq 0$*.*

(iii) In the supremum, W *may be restricted to point masses and to laws with density proportional to* $\mathrm{e}^{-\psi}$ *on a segment, with* ψ *the maximum of at most three affine functions, and the noise to Gaussian or to uniform laws.*

Part (i) is the proof of Theorem 3, applied at $r = Q_q(|W|)$ itself. In part (ii), the extremal law above has its mean inside the interval, and a slightly tilted centred law whose density rises through the right end gains coverage from small noise, at the order of the noise variance; so the usual rank is not exactly valid even for centred residuals. Part (iii) follows from the localization theorem of Fradelizi and Guédon (2006), with the coverage and the two bounds on the mean as linear constraints, and reduces $p_{\mathcal{C}}^{\zeta}(q)$ to a search over six parameters (Supplementary § S3.4). Table 3 gives the values. They are numerical: each is attained by an explicit law, checked in 30-digit arithmetic, so no lower level is valid; that it is the supremum rests on a global search, which no other log-concave law among 3700 tried exceeded.

For centred residuals the worst law found has excess 2.8×10^{-6} at

Table 3: Centred log-concave residuals: excess $p_{\mathcal{C}}^{\zeta}(q) - q$ of the level over q, and the calibration size K from which the usual rank's latent reliability is below 0.5 at the worst law, $\delta = 0.05$

	Gaussian, $q = 0.8$		Gaussian, $q = 0.9$		symm. unimodal, $q = 0.9$	
ζ	excess	K	excess	K	excess	K
0	1.7×10^{-4}	1.5×10^{7}	2.8×10^{-6}	3.1×10^{10}	1.3×10^{-5}	1.4×10^{9}
0.1	3.9×10^{-4}	2.8×10^{6}	3.1×10^{-5}	2.5×10^{8}	1.1×10^{-4}	1.8×10^{7}
0.3	1.2×10^{-3}	2.8×10^{5}	4.1×10^{-4}	1.4×10^{6}	1.1×10^{-3}	1.9×10^{5}
0.5	3.3×10^{-3}	4.1×10^{4}	7.4×10^{-4}	4.4×10^{5}	1.7×10^{-3}	8.6×10^{4}
≥ 1	4.4×10^{-3}	2.2×10^{4}	8.0×10^{-4}	3.8×10^{5}	1.8×10^{-3}	7.6×10^{4}

Rows with $\zeta \geq 1$ are $\Psi_{\mathcal{C}}(q) - q$ (proved); the other rows are numerical. K from an exact scan of the binomial reliability.

$q = 0.9$ under Gaussian noise, nearly three hundred times smaller than $\Psi_{\mathcal{N}}(0.9) - 0.9$; at that law the usual rank's reliability is first below 0.95 at $K = 966$, by 2×10^{-5}, and stays above 0.948 up to $K = 3 \times 10^{6}$, a negligible loss though not zero, and a statement about the laws found rather than a certified bound. The excess grows quickly with the bias: at $\zeta = 0.1$ and 0.3 the usual rank's reliability is below 0.5 from $K \approx 2.5 \times 10^{8}$ and 1.4×10^{6}, and the whole excess is reached at $\zeta = 1$. At $q = 0.8$ the centred margin is larger, 1.7×10^{-4}, attained by a centred exponential law. The worst laws have a hard edge just outside the threshold. The raise of Theorem 3 is thus insurance against a biased centre, at eight ranks at $K = 10^{4}$ and $q = 0.9$. The proposition is for log-concave residuals; the localization argument does not cover $\mathcal{B}$.

4.3 Robustness of the latent assumptions

The common latent law in Theorem 3 is the assumption a user is least able to check. Two relaxations keep an exact guarantee.

Corollary 1. *In the setting of § 2, let* $W_1, \dots, W_K, W_{\text{new}}$ *be independent with laws* $G_1, \dots, G_K, G_0$*, and let* $r_0 = Q_q(|W_{\text{new}}|)$*. If* $G_i \in \mathcal{B}$ *and* $Q_q(|W_i|) \geq r_0$ *for every* $i = 1, \dots, K$*, then the bound of Theorem 3 holds. The law* G_0 *need not be bi-log-concave.*

Proof. The first display in the proof of Theorem 3 uses only the law of W_{new}, with r_0 in place of r. For $s < r_0 \leq Q_q(|W_i|)$, $\text{pr}(|W_i| \leq s) < q$, and the transfer applied to $G_i \in \mathcal{B}$ and the noise of unit i gives $\text{pr}(|V_i| < r_0) \leq p_{\mathcal{C}}(q)$. □

The condition says that the new unit is no harder to cover than any calibration unit at the q-quantile of $|W|$. It is sufficient, not necessary: a few easier calibration units are absorbed by slack in the others (Supplementary § S3.2). It cannot simply be dropped: at the extremal law lowering every $Q_q(|W_i|)$ by a factor $1 - 5 \times 10^{-6}$ lowers the reliability to 0.094 at $K = 10^6$, and if the new unit is a random draw from a population of units with different laws, a shape assumption on the components gives nothing, because mixtures of point masses are arbitrary laws. If the noise scale is random

and the latent residual depends on it, the corollary applies conditionally on the scale; without that, a large noise scale exactly for units whose residual lies just outside the threshold carries them inside, as can happen when the noise variance is a function of the latent quantity, such as $\theta_i(1-\theta_i)/n_i$ for estimated proportions.

The second relaxation lets a fraction of the latent law be arbitrary. For $0 \le \xi < 1$ let $\mathcal{B}_\xi = \{(1-\xi)G + \xi H : G \in \mathcal{B},\ H \text{ any law}\}$.

Proposition 5. *Let $\mathcal{C}$ be the Gaussian or the symmetric unimodal class, $q \ge 0.53$ and $0 \le \xi < 1-q$, and let*

$$p_\xi(q) = (1-\xi)\,\Psi_{\mathcal{C}}\{q/(1-\xi)\} + \xi/2,$$

and $p_\xi(q) = (1+q)/2$ for $\xi \ge 1-q$. Noisy coverage above $p_\xi(q)$ at a threshold guarantees latent coverage q at that threshold for every $W \in \mathcal{B}_\xi$ and every noise with law in $\mathcal{C}$, and no lower level does, even with G log-concave and H a point mass. The level increases continuously from $\Psi_{\mathcal{C}}(q)$ at $\xi = 0$ to the shape-free level $(1+q)/2$ at $\xi = 1-q$, and Theorem 3 and Corollary 1 hold with $\mathcal{B}_\xi$ and $p_\xi(q)$ in place of $\mathcal{B}$ and $p_{\mathcal{C}}(q)$, the rank again being sharp.

The bound $q \ge 0.53$ is what the proof needs and covers every q of practical interest. The worst contamination is a layer just outside the threshold, half of which the noise carries inside, as in Proposition 1; the rest of the

latent law is then extremal for the reduced target $q/(1-\xi)$. The proof is in Supplementary § S3.3. At $q = 0.9$ under Gaussian noise the level rises by about $0.48\,\xi$: it is 0.90322, 0.90564 and 0.92517 for $\xi = 0.5\%$, 1% and 5%, rounded up, and the ranks at $K = 10^3$ are 919, 922 and 940, against 917 for $\xi = 0$ and 962 without a shape assumption. Conversely, a given rank k is valid over $\mathcal{B}_\xi$ exactly when $p_\xi(q)$ is at most the δ-quantile of the $\mathrm{Beta}(k, K+1-k)$ law, which gives the contamination it tolerates: the rank 105 at $K = 110$ tolerates $\xi = 1.2\%$, but the rank of Theorem 3 at $K = 10^3$ tolerates only 0.09%, and at $K = 10^4$ only 0.01%. A user who doubts the shape assumption should calibrate at $p_\xi(q)$ for a plausible ξ. If calibration and new units are independent draws from a population whose latent law is in $\mathcal{B}_\xi$, the guarantee holds for a random new unit.

Noise outside the class can be priced in the same way, though not sharply. If the noise of each unit is drawn, independently of its latent residual, from a law in $\mathcal{C}$ with probability $1-\eta$ and from an arbitrary law otherwise, the noisy coverage of the second part is at most one, so noisy coverage above $p_{\xi,\eta}(q) = (1-\eta)p_\xi(q) + \eta$ guarantees latent coverage q for every $W \in \mathcal{B}_\xi$, and Theorem 3 holds at this level. At $q = 0.9$ under Gaussian noise with $\xi = 0$, $\eta = 1\%$ and 5% give 0.90180 and 0.90577, and ranks 918 and 922 at $K = 10^3$. A share of noise outside the class costs at

most $1 - p_\xi(q)$, about 0.1, per unit share, against about 0.48 for the latent law.

5. Numerical studies

All reliabilities below are latent, with $\delta = 0.05$. Where a law is specified exactly, the reliability is computed exactly from a binomial distribution; otherwise latent coverage given each data set is exact and the reliability is a share of data sets.

5.1 When the correction is needed

At the extremal law for Gaussian noise, scaled so that the noisy coverage at the latent q-quantile of $|W|$ equals $\Psi_{\mathcal{N}}(q)$, Figure 1 compares the marginal rank, the usual rank and the rank of Theorem 3; Supplementary Table S4 gives the exact values. At $q = 0.9$ the usual rank has reliability 0.918 at $K = 10^4$ and 0.150 at $K = 10^6$, against 0.952 and 0.950 for the rank of Theorem 3. The loss is in the guarantee, not in the coverage: at $K = 10^6$ the mean latent coverage of the usual rank's interval is 0.89968, and that of the rank of Theorem 3 is 0.90050. A reliability of 0.150 means that 15% of calibration samples give an interval with latent coverage at least 0.9, not a coverage of 15%.




Figure 1: Exact latent reliability (top) and mean latent coverage (bottom) at the extremal law for Gaussian noise, $q = 0.8$, 0.9 and 0.95, $\delta = 0.05$, against K on a logarithmic axis: rank of Theorem 3 (solid), usual rank (dashes), marginal rank (dots, top only); grey lines at 0.95 and at q. Reliability is the share of calibration samples whose interval has latent coverage at least q; the mean coverage of the usual rank never falls more than 0.001 below q for $q = 0.9$.

The usual rank loses the guarantee as K grows, while the rank of Theorem 3 stays above 0.95 and tends to it. Noise laws drawn per unit within a class delay the failure of the usual rank but do not remove it; the class rank kept reliability at least 0.9501 in all 20 such cases (Supplementary § S6.2). Under contamination of the extremal law the rank of Theorem 3 fails as well, with reliability 0.51 at $K = 10^4$ for $\xi = 1\%$ and $q = 0.9$, while the rank at level $p_\xi(q)$ keeps at least 0.950 (Supplementary § S6.6).

Between the extremal and the benign laws, the reliability was computed exactly for 8028 designs (Supplementary § S6.9): log-concave residuals that are symmetric, smoothly skewed or J-shaped, with the mode at an end of the support, biased up to $\zeta = 0.75$; Gaussian or uniform noise with variance 0.001 to 3 times the latent one, equal or heterogeneous; small K; and estimated proportions, whose noise variance depends on the latent value. For the symmetric and smoothly skewed residuals of the study the usual rank was valid at every K and noise level. It lost reliability only for J-shaped residuals under small noise: for a reflected exponential with $\zeta = 0.5$ and noise variance 0.01 at $q = 0.8$, 0.934, 0.841 and 0.319 at $K = 10^3$, 10^4 and 10^5, against 0.967, 0.982 and 0.999 for the rank of Theorem 3, which kept at least 0.950 in every design. These are findings for the designs: to first order in the noise variance v, the excess $H - q$ is $\frac{1}{2}v\{f'(r) - f'(-r)\}$,

so a smooth density gains coverage from small noise when $f'(r) > f'(-r)$ and loses it when $f'(r) < f'(-r)$; a density that increases outward at one end is necessary for a gain but not sufficient, since a steeper increase at the other end can outweigh it.

A rank computed for one noise class can fail under noise outside it (Supplementary § S6.8). At $q = 0.9$ the Gaussian rank survives Laplace, logistic and t_5 noise at the extremal law, but not uniform or skewed noise: under centred exponential noise its reliability is 0.58 at $K = 10^4$. Strongly skewed noise outside the log-concave class, such as the sampling error of a mean of a few units of a skewed population, exceeds even the log-concave level; the noise class must be chosen with the skewness of the noise in mind.

5.2 The cost with the same information

No rule in this comparison is told the noise law within its class or the variances. Six standardized latent laws were used: normal, Laplace, centred gamma with shape 2, truncated exponential, a bimodal bi-log-concave mixture that is not log-concave, and t_3, outside $\mathcal{B}$, as a control. Eight noise designs, with heterogeneous variances $D_i = 0.577 L_i / E(L_i)$ for L_i lognormal with log-scale 0.7, comprise five single laws (Gaussian, uniform, Laplace, t_3, centred exponential) and laws drawn per unit within the symmetric uni-

modal class, within the mean-zero log-concave class, and across the two classes; 2000 data sets per setting.

Table 4: Rules with the same information at $K = 1000$: rank and width relative to the usual rank and to the rule for the same noise class without a latent shape assumption, ranges over the settings a rule covers (medians in parentheses)

	$q = 0.8$			$q = 0.9$		
rule	k	+usual	−no shape	k	+usual	−no shape
usual	822	0	–	916	0	–
Gaussian	826	0.9–1.2	20–26 (24)	917	0.3–0.4	16–21 (21)
symm. unimodal	829	1.5–2.2	19–28 (24)	918	0.6–1.0	15–24 (20)
mean-zero log-conc.	–	–	–	922	1.8–2.9	19–33 (26)
no shape, median 0	916	25–42	–	962	18–36	–
no shape, log-conc.	941	36–63	–	974	25–54	–

Widths in percent: +usual, above the usual rank; −no shape, below the rule for the same noise class without a shape assumption (median-zero for the Gaussian and symmetric unimodal rules). The log-concave level is certified only at $q = 0.9$.

Every rule, the usual rank included, had estimated reliability at least 0.998 in every setting, covered or not, with 2000 data sets per setting: these laws are far from the extremal one, so they measure the cost of the guarantee, not its need. The cost over the usual rank is at most 3% of width at $K = 1000$ and nothing at $K = 110$, where the bi-log-concave ranks equal the usual rank; the saving over the rules without a shape assumption is 15–33% at $K = 1000$ and 18–48% at $K = 110$ for $q = 0.8$ and 0.9, where at $q = 0.95$ only the bi-log-concave rules have a valid rank at all. Against the oracle radius $Q_q(|W|)$, the usual rank was 44–86% wider at $K = 110$ and 25–55% wider at $K = 1000$, with mean latent coverage 0.95–0.99:

the dominant cost is that of calibrating on noisy scores at all, with noise variance about half the latent one.

5.3 Comparison with existing methods

Methods that use more information were run on the same latent laws at $q = 0.9$, with 300 data sets at $K = 110$ and 200 at $K = 1000$, under Gaussian, Laplace and mixed symmetric unimodal noise with the same variances. Each rule is judged against its own target (Table 5).

Table 5: Comparison with existing methods at $q = 0.9$, grouped by target: information used, whether the target was met over the bi-log-concave latent laws and three noise scenarios, and width relative to the usual rank

rule	information	target met	width vs usual (%)	
			$K = 110$	$K = 1000$
target: reliability ≥ 0.95				
Theorem 3, Gaussian	none	yes, 1.000	0	0.3–0.5
Theorem 3, symm. unimodal	none	yes, 1.000	0	0.6–0.9
Markov rule*	known D_i	yes, 1.000	19–25	20–26
Fay–Herriot tolerance*	known D_i	no, down to 0.900	−24 to −15	−20 to −16
target: mean latent coverage ≥ 0.9				
LatentCP	known D_i	yes, ≥ 0.97	4–7	13–21
deconvolution	known D_i	no, 0.88–0.92	−41 to −30	−32 to −19

*Constructed here (Supplementary § S4.7); the Fay–Herriot tolerance multiplier uses the parametric bootstrap of Chatterjee, Lahiri, and Li (2008). Monte Carlo standard error of a reliability near 0.95: about 0.013 (300 data sets) and 0.015 (200). Fay–Herriot met its target for normal latent laws with Gaussian noise; *down to* is a minimum over 30 settings, each with that standard error. Deconvolution: our implementation of Algorithm 1 of Cohen, Goldberger, and Tirer (2025) with bin width 0.2.

The rank of Theorem 3, which uses no noise information, met its target in every setting. Among rules with the same target, the one that is shorter, the Fay–Herriot tolerance interval, relies on the normal model and lost the

guarantee outside it. LatentCP met its weaker, marginal target throughout, also where its Gaussian forward model is wrong. The deconvolution threshold, which has no finite-sample guarantee, fell below nominal coverage on average in our implementation, which averages the Gaussian kernels of heterogeneous units and fixes the bin width; this reflects our adaptation to heterogeneous noise and is not evidence against the method in its own setting of a known common kernel. Rules that are given the variances and assume Gaussian noise can be shorter while keeping the guarantee, by about 10% in one setting (Supplementary § S4); that is what the variances buy.

6. Application: school districts

The California Academic Performance Index data (Lumley, 2004) give the 2000 index of every school; districts with at least five schools form the population. Each of 480 replications, in four conditions, splits the districts into 110 training, 110 calibration and the remaining evaluation districts, samples $n = 2$ or 3 schools per training and calibration district, fits a ridge predictor and scores each calibration district by its sample-mean residual; the target of an evaluation district is its true mean residual. At $K = 110$ every class rank equals the usual rank, $k = 105$, so the application checks

the usual interval rather than a new one. It covered at least 90% of the evaluation districts in all 480 replications, with mean coverage 0.97 to 0.99. The Fay–Herriot tolerance interval with plugged-in variance estimates was 19–26% shorter but covered fewer than 90% of the districts in up to 9.2% of the replications of a condition; LatentCP, drawing the variance of a new district from the calibration estimates, was 4.6–5.7% wider. The assumptions cannot be confirmed here: the districts form a finite population, the latent law is a mixture over districts, and the noise of a mean of two or three schools is discrete (Supplementary § S7).

7. Discussion

Without symmetry, noisy calibration is close to valid but not exactly so: under bi-log-concavity and Gaussian noise at $q = 0.9$ the sharp transfer puts the needed level less than a thousandth above q, and 0.0044 above at $q = 0.8$; a share ξ of the latent law outside the class raises it by about $0.48\,\xi$ at $q = 0.9$, and for small noise the usual threshold can lose its guarantee only if the residual density satisfies $f'(r) > f'(-r)$. The paper is about validity. Calibrating on noisy scores makes the interval much wider than the oracle radius, 25–86% in § 5.2 with noise variance about half the latent one, and only knowledge of the noise variances can recover part of

that width (Supplementary § S4); the correction studied here adds at most a few percent. At the worst centred log-concave laws found numerically the loss of reliability is negligible at any realistic calibration size; the exact correction, a raise of the level from q to $\Psi_{\mathcal{C}}(q)$, needs no noise variances, costs at most a few ranks and keeps a high-probability guarantee for every noise configuration in the class. The ordering of Corollary 1 and the share ξ of Proposition 5 are the assumptions that matter most and are hardest to check. Bi-log-concavity cannot be checked on the latent scale, but convolution with log-concave noise preserves it: with a common log-concave noise law, which the heterogeneous setting does not provide, a confidence band for the signed noisy residuals that contains no bi-log-concave distribution function (Dümbgen, Kolesnyk, and Wilke, 2017) rules it out; the bands of Laha, Miao, and Wellner (2021) could serve the same purpose. A known Gaussian noise variance allows the threshold to be widened instead of its rank raised, by an amount of the order of the noise standard deviation, or of its variance when the latent mean is away from the threshold (Supplementary § S4).

The centred levels of Proposition 4 are numerical below $\zeta = 1$ and are for log-concave residuals only; a certificate, and an extension to $\mathcal{B}$, would complete them. Two further questions remain open. For mean-zero log-

concave noise the two-sided level is known only within $(0.90517, 0.906]$ at $q = 0.9$, and the extremality of the centred exponential law is supported by certified computation but not proved. Whether bi-log-concavity is necessary for a nontrivial transfer, or some weaker condition excluding a layer of mass at the threshold suffices, is also open.

Supplementary Material

The Supplementary Material contains the proofs of all results; the certificates in ball arithmetic; the theory for a known Gaussian noise variance; the full tables of the numerical studies, including mixed noise, latent laws that differ between units, contamination and miscentring; details of the application; and a map from every number to the code. Code and data access are at `https://github.com/kunhailP/noisy-calibration-latent-coverage`.

References

Anderson, T. W. (1955). The integral of a symmetric unimodal function over a symmetric convex set and some probability inequalities. *Proc. Amer. Math. Soc. 6*, 170–176.

Armstrong, T. B., M. Kolesár, and M. Plagborg-Møller (2022). Robust empirical Bayes confidence intervals. *Econometrica 90*, 2567–2602.

Barber, R. F., E. J. Candès, A. Ramdas, and R. J. Tibshirani (2023). Conformal prediction

beyond exchangeability. *Ann. Statist. 51*, 816–845.

Bersson, E. and P. D. Hoff (2024). Optimal conformal prediction for small areas. *J. Surv. Statist. Methodol. 12*, 1464–1488.

Bian, M. and R. F. Barber (2023). Training-conditional coverage for distribution-free predictive inference. *Electron. J. Statist. 17*, 2044–2066.

Burris, K. C. and P. D. Hoff (2020). Exact adaptive confidence intervals for small areas. *J. Surv. Statist. Methodol. 8*, 206–230.

Cauchois, M., S. Gupta, A. Ali, and J. C. Duchi (2024). Robust validation: confident predictions even when distributions shift. *J. Amer. Statist. Assoc. 119*, 3033–3044.

Chatterjee, S., P. Lahiri, and H. Li (2008). Parametric bootstrap approximation to the distribution of EBLUP and related prediction intervals in linear mixed models. *Ann. Statist. 36*, 1221–1245.

Clarkson, J., W. Xu, M. Cucuringu, and G. Reinert (2024). Split conformal prediction under data contamination. In *Proceedings of the 13th Symposium on Conformal and Probabilistic Prediction with Applications*, Volume 230 of *Proceedings of Machine Learning Research*, pp. 5–27.

Cohen, Y., J. Goldberger, and T. Tirer (2025). Efficient conformal prediction for regression models under label noise. *arXiv preprint arXiv:2509.15120*.

Dass, S. C., T. Maiti, H. Ren, and S. Sinha (2012). Confidence interval estimation of small area

parameters shrinking both means and variances. *Survey Methodology 38*, 173–187.

Delaigle, A. and P. Hall (2016). Methodology for non-parametric deconvolution when the error distribution is unknown. *J. R. Statist. Soc. B 78*, 231–252.

Delaigle, A. and A. Meister (2008). Density estimation with heteroscedastic error. *Bernoulli 14*, 562–579.

Diao, L., D. D. Smith, G. S. Datta, T. Maiti, and J. D. Opsomer (2014). Accurate confidence interval estimation of small area parameters under the Fay–Herriot model. *Scand. J. Statist. 41*, 497–515.

Dümbgen, L., P. Kolesnyk, and R. A. Wilke (2017). Bi-log-concave distribution functions. *J. Statist. Plann. Inference 184*, 1–17.

Dunn, R., L. Wasserman, and A. Ramdas (2023). Distribution-free prediction sets for two-layer hierarchical models. *J. Amer. Statist. Assoc. 118*, 2491–2502.

Einbinder, B.-S., S. Feldman, S. Bates, A. N. Angelopoulos, A. Gendler, and Y. Romano (2024). Label noise robustness of conformal prediction. *J. Mach. Learn. Res. 25*(328), 1–66.

Fay, R. E. and R. A. Herriot (1979). Estimates of income for small places: an application of James–Stein procedures to census data. *J. Amer. Statist. Assoc. 74*, 269–277.

Feldman, S., S. Bates, and Y. Romano (2026). Conformal prediction with corrupted labels: uncertain imputation and robust re-weighting. In *International Conference on Learning Representations*.

Fradelizi, M. and O. Guédon (2004). The extreme points of subsets of s-concave probabilities and a geometric localization theorem. *Discrete Comput. Geom. 31*, 327–335.

Fradelizi, M. and O. Guédon (2006). A generalized localization theorem and geometric inequalities for convex bodies. *Adv. Math. 204*, 509–529.

Guille-Escuret, C. and E. Ndiaye (2024). From conformal predictions to confidence regions. *arXiv preprint arXiv:2405.18601*.

Hall, P. and T. Maiti (2006). On parametric bootstrap methods for small area prediction. *J. R. Statist. Soc. B 68*, 221–238.

Ignatiadis, N. and S. Wager (2022). Confidence intervals for nonparametric empirical Bayes analysis. *J. Amer. Statist. Assoc. 117*, 1149–1166.

Laha, N., Z. Miao, and J. A. Wellner (2021). Bi-s^*-concave distributions. *J. Statist. Plann. Inference 215*, 127–157.

Liao, C.-T., T.-Y. Lin, and H. K. Iyer (2005). One- and two-sided tolerance intervals for general balanced mixed models and unbalanced one-way random models. *Technometrics 47*, 323–335.

Lin, T.-Y., C.-T. Liao, and H. K. Iyer (2008). Tolerance intervals for unbalanced one-way random effects models with covariates and heterogeneous variances. *J. Agric. Biol. Environ. Stat. 13*, 221–241.

Lumley, T. (2004). Analysis of complex survey samples. *J. Statist. Software 9*(8), 1–19.

Penso, C., J. Goldberger, and E. Fetaya (2025). Conformal prediction of classifiers with many classes based on noisy labels. In *Proceedings of the 14th Symposium on Conformal and Probabilistic Prediction with Applications*, Volume 266 of *Proceedings of Machine Learning Research*, pp. 82–95.

Rao, J. N. K. and I. Molina (2015). *Small Area Estimation* (2nd ed.). Hoboken, NJ: Wiley.

Sampford, M. R. (1953). Some inequalities on Mill's ratio and related functions. *Ann. Math. Statist. 24*, 130–132.

Sesia, M., Y. X. R. Wang, and X. Tong (2025). Adaptive conformal classification with noisy labels. *J. R. Statist. Soc. B 87*, 796–815.

Soloff, J. A., A. Guntuboyina, and B. Sen (2025). Multivariate, heteroscedastic empirical Bayes via nonparametric maximum likelihood. *J. R. Statist. Soc. B 87*, 1–32.

Vovk, V. (2012). Conditional validity of inductive conformal predictors. In *Proceedings of the Asian Conference on Machine Learning*, Volume 25 of *Proceedings of Machine Learning Research*, pp. 475–490.

Wang, C. M. and H. K. Iyer (1994). Tolerance intervals for the distribution of true values in the presence of measurement errors. *Technometrics 36*, 162–170.

Wang, J. and W. A. Fuller (2003). The mean squared error of small area predictors constructed with estimated area variances. *J. Amer. Statist. Assoc. 98*, 716–723.

Wieczorek, J. (2023). Design-based conformal prediction. *Survey Methodology 49*, 443–473.

Wilks, S. S. (1941). Determination of sample sizes for setting tolerance limits. *Ann. Math. Statist. 12*, 91–96.

Yoshimori, M. and P. Lahiri (2014). A second-order efficient empirical Bayes confidence interval. *Ann. Statist. 42*, 1233–1261.

You, Y. and B. Chapman (2006). Small area estimation using area level models and estimated sampling variances. *Survey Methodology 32*, 97–103.

Yu, C. and P. D. Hoff (2018). Adaptive multigroup confidence intervals with constant coverage. *Biometrika 105*, 319–335.

Zhang, L.-C. and T. Tuoto (2026). Conformal confidence intervals with an application to small area estimation. *arXiv preprint arXiv:2608.02766*.

Zheng, M., W. Zhou, and S. Zhu (2026). Beyond prediction: conformal inference for latent distributional parameters. *arXiv preprint arXiv:2608.03607*.

Kun Woo Park, Department of Political Science and International Relations, Kookmin University, 77 Jeongneung-ro, Seongbuk-gu, Seoul 02707, Republic of Korea

E-mail: pkw6094@kookmin.ac.kr