

The Tale of a Boundary-Layer Problem

Martin Kunz

Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University
in Prague

April 24, 2023

Abstract

I will demonstrate the solution with all its steps (and missteps) to the boundary layer problem

$$\varepsilon y'' = y \cdot y' - y, \quad (1)$$

such that

$$y(0) = 1 \quad \text{and} \quad y(1) = -1 \quad \text{and where} \quad \varepsilon \ll 1. \quad (2)$$

The journey towards the solution will take us on an excursion through some basics of *perturbation theory* and *asymptotic approximation matching*. I will demonstrate visualizations that will give you insight into the reasoning behind the steps taken towards the solution serving as an exhibition for the leverage in problem solving accessible to us through the computational power at our fingertips.

Outline

- 1 Motivation
- 2 Theory
- 3 Solution of the DE
- 4 Interactive Solution

Classifying the problem

$$\varepsilon y'' = y \cdot y' - y$$

such that

$$y(0) = 1 \quad \text{and} \quad y(1) = -1 \quad \text{and where} \quad \varepsilon \ll 1.$$

Classifying the problem

$$\varepsilon y'' = y \cdot y' - y$$

such that

$$y(0) = 1 \quad \text{and} \quad y(1) = -1 \quad \text{and where} \quad \varepsilon \ll 1.$$

ordinary single unknown function

Classifying the problem

$$\varepsilon y'' = y \cdot y' - y$$

such that

$$y(0) = 1 \quad \text{and} \quad y(1) = -1 \quad \text{and where} \quad \varepsilon \ll 1.$$

ordinary single unknown function (as opposed to a PDE)

Classifying the problem

$$\varepsilon y'' = y \cdot y' - y$$

such that

$$y(0) = 1 \quad \text{and} \quad y(1) = -1 \quad \text{and where} \quad \varepsilon \ll 1.$$

ordinary single unknown function

second-order contains a second order derivative

Classifying the problem

$$\varepsilon y'' = y \cdot y' - y$$

such that

$$y(0) = 1 \quad \text{and} \quad y(1) = -1 \quad \text{and where} \quad \varepsilon \ll 1.$$

ordinary single unknown function

second-order contains a second order derivative

first-degree highest degree of the unknown function is 1

Classifying the problem

$$\varepsilon y''(x) = y(x) \cdot y'(x) - y(x)$$

such that

$$y(0) = 1 \quad \text{and} \quad y(1) = -1 \quad \text{and where} \quad \varepsilon \ll 1.$$

ordinary single unknown function

second-order contains a second order derivative

first-degree highest degree of the unknown function is 1

homogeneous does not contain a free term in the x variable

Classifying the problem

$$\varepsilon y'' = y \cdot y' - y$$

such that

$$y(0) = 1 \quad \text{and} \quad y(1) = -1 \quad \text{and where} \quad \varepsilon \ll 1.$$

ordinary single unknown function

second-order contains a second order derivative

first-degree highest degree of the unknown function is 1

homogeneous does not contain a free term in the x variable

non-linear contains a mixed term

Classifying the problem

$$\varepsilon y'' = y \cdot y' - y$$

such that

$$y(0) = 1 \quad \text{and} \quad y(1) = -1 \quad \text{and where} \quad \varepsilon \ll 1.$$

ordinary single unknown function

second-order contains a second order derivative

first-degree highest degree of the unknown function is 1

homogeneous does not contain a free term in the x variable

non-linear contains a mixed term

(boundary-value) constrained by values on the boundaries

Similar Differential Equations

What is the biggest problem in solving this DE?

Similar Differential Equations

What is the biggest problem in solving this DE?

It is *non-linear* !

Similar Differential Equations

Have we encountered other non-linear DEs?

Similar Differential Equations

Have we encountered other non-linear DEs?

- Newtons gravitational equation

Similar Differential Equations

Have we encountered other non-linear DEs?

- Newtons gravitational equation (ordinary, second-order, second-degree, homogeneous, non-linear)

Similar Differential Equations

Have we encountered other non-linear DEs?

- Newtons gravitational equation (ordinary, second-order, second-degree, homogeneous, non-linear)
- Simple pendulum

Similar Differential Equations

Have we encountered other non-linear DEs?

- Newtons gravitational equation (ordinary, second-order, second-degree, homogeneous, non-linear)
- Simple pendulum (ordinary, second-order, homogeneous, non-linear)

Similar Differential Equations

Have we encountered other non-linear DEs?

- Newtons gravitational equation (ordinary, second-order, second-degree, homogeneous, non-linear)
- Simple pendulum (ordinary, second-order, homogeneous, non-linear)

How did we manage to solve these equations?




Figure: Comparison of the numerical solution and the small angle approximation of the simple pendulum DE for the initial angle $\varphi(0) = \pi/4$ and $\dot{\varphi}(0) = 0$.




Figure: Comparison of the numerical solution and the small angle approximation of the simple pendulum DE for the initial angle $\varphi(0) = \pi/4$ and $\dot{\varphi}(0) = 0$.




Figure: Comparison of the numerical solution and the small angle approximation of the simple pendulum DE for the initial angle $\varphi(0) = \pi/4$ and $\dot{\varphi}(0) = 0$.

Classifying the problem

$$\varepsilon y'' = y \cdot y' - y$$

such that

$$y(0) = 1 \quad \text{and} \quad y(1) = -1 \quad \text{and where} \quad \varepsilon \ll 1.$$

ordinary single unknown function

second-order contains a second order derivative

first-degree highest degree of the unknown function is 1

homogeneous does not contain a free term in the x variable

non-linear contains a mixed term

(boundary-value) constrained by values on the boundaries

Classifying the problem

$$\varepsilon y'' = y \cdot y' - y$$

such that

$$y(0) = 1 \quad \text{and} \quad y(1) = -1 \quad \text{and where} \quad \varepsilon \ll 1.$$

ordinary single unknown function

second-order contains a second order derivative

first-degree highest degree of the unknown function is 1

homogeneous does not contain a free term in the x variable

non-linear contains a mixed term

(boundary-value) constrained by values on the boundaries

singularly-perturbed contains one small term



How good is our numerical solution?

Note (Analytical solution - Salas et al. 2014)

There is an analytical solution to the pendulum DE in the form

$$\varphi(t) = 2 \cdot \arcsin\left(\sin(\varphi_0/2) \cdot \operatorname{cd}(t \cdot \sqrt{g/L}, \sin^2(\varphi_0/2))\right), \text{ where}$$

$$\operatorname{cd}(u, m) = \operatorname{cn}(u, m) / \operatorname{dn}(u, m)$$

$$\operatorname{cn}(u, m) = \cos(\operatorname{am}(u, m)) \quad \operatorname{dn}(u, m) = \frac{d}{du}(\operatorname{am}(u, m))$$

and $\operatorname{am}(u, m) = \varphi$ is the Jacobi amplitude which is the upper-bound of the integral such that

$$u = \int_0^\varphi \frac{d\vartheta}{\sqrt{1 - m \sin^2(\vartheta)}}.$$



Numerical Methods




Figure: Comparison of the analytical solution and the numerical solution of the simple pendulum DE for the initial angle $\varphi(0) = \pi/4$ and $\dot{\varphi}(0) = 0$.



Why would we want to bother with perturbation methods?



Why would we want to bother with perturbation methods, when we can make the numerical solution arbitrarily more precise?



Why would we want to bother with perturbation methods, when we can make the numerical solution arbitrarily more precise?

- Perturbation methods approximation is interpretable



Why would we want to bother with perturbation methods, when we can make the numerical solution arbitrarily more precise?

- Perturbation methods approximation is interpretable
- We can prove some good qualities about the solution



Why would we want to bother with perturbation methods, when we can make the numerical solution arbitrarily more precise?

- Perturbation methods approximation is interpretable
- We can prove some good qualities about the solution



Why would we want to bother with perturbation methods, when we can make the numerical solution arbitrarily more precise?

- Perturbation methods approximation is interpretable
- We can prove some good qualities about the solution (uniform convergence, etc.)



Why would we want to bother with perturbation methods, when we can make the numerical solution arbitrarily more precise?

- Perturbation methods approximation is interpretable
- We can prove some good qualities about the solution (uniform convergence, etc.)
- Numerical solution is not guaranteed to be correct

Why would we want to bother with perturbation methods, when we can make the numerical solution arbitrarily more precise?

- Perturbation methods approximation is interpretable
- We can prove some good qualities about the solution (uniform convergence, etc.)
- Numerical solution is not guaranteed to be correct
- *Size of the latent space for the solution!*

Why would we want to bother with perturbation methods, when we can make the numerical solution arbitrarily more precise?

- Perturbation methods approximation is interpretable
- We can prove some good qualities about the solution (uniform convergence, etc.)
- Numerical solution is not guaranteed to be correct
- *Size of the latent space for the solution!*

Why would we want to bother with perturbation methods, when we can make the numerical solution arbitrarily more precise?

- Perturbation methods approximation is interpretable
- We can prove some good qualities about the solution (uniform convergence, etc.)
- Numerical solution is not guaranteed to be correct
- *Size of the latent space for the solution! (we cannot search the whole space and find all the solutions numerically)*



How precise is a perturbation solution?

Note (Perturbation solution - Beléndez et al. 2006)

There is an approximation to the simple pendulum problem based on a homotopy perturbation method. It is uniformly convergent and is precise from the first few terms of the expansion.



Numerical Methods




Figure: Comparison of the perturbation solution and the analytical solution of the simple pendulum DE for the initial angle $\varphi(0) = \pi/4$ and $\dot{\varphi}(0) = 0$.

2 Theory

- Mathematical Basics
- Perturbation Theory
- Boundary Layers

Theorem (Taylor's theorem)

Given a function $f(\varepsilon)$, suppose its $(n+1)$ st derivative $f^{(n+1)}$ is continuous for $\varepsilon_a < \varepsilon < \varepsilon_b$. Suppose that the points $\varepsilon, \varepsilon_0 \in (\varepsilon_a, \varepsilon_b)$. Then

$$f(\varepsilon) = f(\varepsilon_0) + (\varepsilon - \varepsilon_0)f'(\varepsilon_0) + \dots + \frac{1}{n!}(\varepsilon - \varepsilon_0)^n f^{(n)}(\varepsilon_0) + R_{n+1},$$

where the error term R_{n+1} is given by

$$R_{n+1} = \frac{1}{(n+1)!}(\varepsilon - \varepsilon_0)^{n+1} f^{(n+1)}(\xi),$$

and $\xi \in]\varepsilon, \varepsilon_0[$

Definition (Bachmann-Landau Symbols - Holmes 2012)

1 $f = O(\varphi)$ as $\varepsilon \rightarrow \varepsilon_0^+$ if there is a constant k and $\tilde{\varepsilon}$ so that

$$|f(\varepsilon)| \leq k |\varphi(\varepsilon)| \quad \text{for } \varepsilon_0 < \varepsilon < \tilde{\varepsilon}$$

Definition (Bachmann-Landau Symbols - Holmes 2012)

1 $f = O(\varphi)$ as $\varepsilon \rightarrow \varepsilon_0^+$ if there is a constant k and $\tilde{\varepsilon}$ so that

$$|f(\varepsilon)| \leq k |\varphi(\varepsilon)| \quad \text{for } \varepsilon_0 < \varepsilon < \tilde{\varepsilon}$$

2 $f = o(\varphi)$ as $\varepsilon \rightarrow \varepsilon_0^+$ if $\forall \delta > 0$ there is a $\tilde{\varepsilon}$ so that

$$|f(\varepsilon)| \leq \delta |\varphi(\varepsilon)| \quad \text{for } \varepsilon_0 < \varepsilon < \tilde{\varepsilon}$$

Definition (Asymptotic approximation - Holmes 2012)

Given a function $f(\varepsilon)$ and $\varphi(\varepsilon)$ we say that $\varphi(\varepsilon)$ is an *asymptotic approximation* to $f(\varepsilon)$ as $\varepsilon \rightarrow \varepsilon_0^+$ whenever $f = \varphi + o(\varphi)$ as $\varepsilon \rightarrow \varepsilon_0^+$. We write $f \sim \varphi$ as $\varepsilon \rightarrow \varepsilon_0^+$.

Definition (Asymptotic expansion - Holmes 2012)

- 1 The sequence of functions $\varphi_1(\varepsilon), \varphi_2(\varepsilon), \dots$ form an *asymptotic sequence (are well ordered)* as $\varepsilon \rightarrow \varepsilon_0^+$ if $\varphi_{m+1} = o(\varphi_m)$ as $\varepsilon \rightarrow \varepsilon_0^+$ for $\forall n$.

Definition (Asymptotic expansion - Holmes 2012)

- 1 The sequence of functions $\varphi_1(\varepsilon), \varphi_2(\varepsilon), \dots$ form an *asymptotic sequence (are well ordered)* as $\varepsilon \rightarrow \varepsilon_0^+$ if $\varphi_{m+1} = o(\varphi_m)$ as $\varepsilon \rightarrow \varepsilon_0^+$ for $\forall n$. φ_k are called the *basis functions*.

Definition (Asymptotic expansion - Holmes 2012)

- 1 The sequence of functions $\varphi_1(\varepsilon), \varphi_2(\varepsilon), \dots$ form an *asymptotic sequence (are well ordered)* as $\varepsilon \rightarrow \varepsilon_0^+$ if $\varphi_{m+1} = o(\varphi_m)$ as $\varepsilon \rightarrow \varepsilon_0^+$ for $\forall n$. φ_k are called the *basis functions*.
- 2 Let $\varphi_1(\varepsilon), \varphi_2(\varepsilon), \dots$ be basis functions. Then $f(\varepsilon)$ has an *asymptotic expansion* to n terms for this sequence if

$$f = \sum_{k=1}^m a_k \varphi_k + o(\varphi_m) \quad \text{for } m \in \hat{n} \quad \text{as } \varepsilon \rightarrow \varepsilon_0^+,$$

where a_k are independent of ε . We can write

$$f \sim a_1 \varphi_1(\varepsilon) + a_2 \varphi_2(\varepsilon) + \dots + a_n \varphi_n(\varepsilon) \quad \text{as } \varepsilon \rightarrow \varepsilon_0^+.$$

Asymptotic Solution of the Projectile Problem

$$\frac{d^2x}{dt^2} = -\frac{g \cdot R^2}{(x + R)^2} \quad \text{for } 0 < t.$$

Asymptotic Solution of the Projectile Problem

$$\frac{d^2x}{dt^2} = -\frac{g \cdot R^2}{(x + R)^2} \quad \text{for } 0 < t.$$

First we scale the equation to make the approximation easier

Asymptotic Solution of the Projectile Problem

$$\frac{d^2x}{dt^2} = -\frac{g \cdot R^2}{(x + R)^2} \quad \text{for } 0 < t.$$

First we scale the equation to make the approximation easier

$$\tau = t \cdot g/v_0$$

$$y(\tau) = x(t) \cdot g/v_0^2$$

Asymptotic Solution of the Projectile Problem

$$\frac{d^2x}{dt^2} = -\frac{g \cdot R^2}{(x + R)^2} \quad \text{for } 0 < t.$$

First we scale the equation to make the approximation easier

$$\tau = t \cdot g/v_0$$

$$y(\tau) = x(t) \cdot g/v_0^2$$

Doing so transforms the problem into

$$\frac{d^2y}{d\tau^2} = -\frac{1}{(1 + \varepsilon y)^2} \quad \text{for } 0 < \tau,$$

where

$$y(0) = 0 \quad \text{and} \quad y'(0) = 1 \quad \text{and} \quad \varepsilon = v_0^2/Rg.$$

Perturbation Theory




Figure: Comparison of the two term approximation, the linearised problem (one-term approximation) and the numerical solution for $\varepsilon = 1/10$

Boundary Layer

A boundary layer in lay terms is an interval of the solution of a DE where the derivative is large compared with surrounding times.




Figure: Shlieren image of the air waves of a supersonic airflow produced by a fighter jet [Source: NASA (2015)]

3 Solution of the DE

- Locating the Boundary Layers
- Symmetry of the Unknown
- Phase Portrait
- Perturbation Theory
 - Outer Solution
- Inner Solution

Plausibility Argument of no Border Boundary Layers - Holmes 2012

If we are looking for a boundary layer at $x = 0$, then $y' < 0$ and if we assume that $y'' > 0$, then $\varepsilon y'' > 0$, but $y \cdot (y' - 1)$ on the left hand side is both > 0 on some interval and < 0 . Therefore we can rule out a boundary layers at the interval borders for $y'' > 0$.

Is there any symmetry in the DE?

Is there any symmetry in the DE?

$$(x, y) \rightarrow (1 - x, -y)$$

Phase Portrait of the DE

$$\varepsilon y'' = y \cdot y' - y$$

Phase Portrait of the DE

$$y'(t) = z(t)$$

$$z'(t) = \frac{1}{\varepsilon} \cdot y(t) \cdot (z(t) - 1)$$

Phase Portrait




Figure: Arrows representing the vector field of our system of differential equations

Phase Portrait




Figure: Trajectories of imaginary particles in the phase portrait of our system of differential equations for $\varepsilon = 0.1$

Phase Portrait




Figure: Trajectories of imaginary particles in the phase portrait of our system of differential equations for $\varepsilon = 1$ with the boundary condition hyperplanes

Phase Portrait




Figure: Solutions B_0 , M and B_1 in the phase plane of the DE

Outer Solution

Outer Solution

$$y_0(x) = x + a \quad \text{for some } a \in \mathbb{R}$$

Outer Solution

$$y_0(x) = x + a \quad \text{for some } a \in \mathbb{R}$$

$$y_0^L(x) = 1 + x \quad \text{and} \quad y_0^R(x) = x - 2$$

Scaling

For the boundaries, we have to scale the solutions as

$$X \equiv \frac{x - x_0}{\delta},$$

where δ is the thickness of the layer and $\delta = \delta(\varepsilon)$.

Scaling

For the boundaries, we have to scale the solutions as

$$X \equiv \frac{x - x_0}{\delta},$$

where δ is the thickness of the layer and $\delta = \delta(\varepsilon)$.

$$Y(X) \equiv y(x) = y(x_0 + \delta X)$$

Scaling

For the boundaries, we have to scale the solutions as

$$X \equiv \frac{x - x_0}{\delta},$$

where δ is the thickness of the layer and $\delta = \delta(\varepsilon)$.

$$Y(X) \equiv y(x) = y(x_0 + \delta X)$$

If δ is chosen correctly, then Y and all its derivatives should be $O(1)$ as $\varepsilon \rightarrow 0^+$ and we can find the limit when $\varepsilon/\delta^2 = 1/\delta$ or $\delta = \varepsilon$.

Scaling

For the boundaries, we have to scale the solutions as

$$X \equiv \frac{x - x_0}{\delta},$$

where δ is the thickness of the layer and $\delta = \delta(\varepsilon)$.

$$Y(X) \equiv y(x) = y(x_0 + \delta X)$$

If δ is chosen correctly, then Y and all its derivatives should be $O(1)$ as $\varepsilon \rightarrow 0^+$ and we can find the limit when $\varepsilon/\delta^2 = 1/\delta$ or $\delta = \varepsilon$. The inner equation layer is

$$\frac{d^2 Y}{dX^2} = Y \frac{dY}{dX} - \varepsilon Y.$$

Inner Solutions

Solving the equation using the same expansion get us

$$Y_0(X) = b \cdot \tanh\left(c - \frac{b}{2} \cdot X\right)$$

Inner Solutions

Solving the equation using the same expansion get us

$$Y_0(X) = b \cdot \tanh\left(c - \frac{b}{2} \cdot X\right)$$

Composing the solutions

$$\tilde{y} = y_{\text{outer}} + Y_{\text{inner}} - y_{\text{match}}$$

Inner Solutions

Solving the equation using the same expansion get us

$$Y_0(X) = b \cdot \tanh\left(c - \frac{b}{2} \cdot X\right)$$

Composing the solutions

$$\tilde{y} = y_{\text{outer}} + Y_{\text{inner}} - y_{\text{match}}$$

When we create these composite solutions, we get




$$\tilde{y}^M = x - \frac{1}{2} - \frac{3}{2} \cdot \tanh\left[\frac{3}{4\varepsilon}\left(x - \frac{1}{2}\right)\right] + O(\varepsilon)$$

$$\tilde{y}^{B0} = x - 2 \cdot \tanh\left[\frac{x}{\varepsilon} - \tanh^{-1}(1/2)\right] + O(\varepsilon)$$

$$\tilde{y}^{B1} = x - 1 - 2 \cdot \tanh\left[\frac{x - 1}{\varepsilon} + \tanh^{-1}(1/2)\right] + O(\varepsilon)$$

4 Interactive Solution

References

-  Beléndez, Augusto et al. (2006). “Application of the homotopy perturbation method to the non-linear pendulum”. In: *European Journal of Physics* 28.1, p. 93.
-  Holmes, Mark H. (2012). *Introduction to perturbation methods*. Vol. 20. Springer Science & Business Media.
-  Salas, Alvaro H et al. (2014). “Exact solution to Duffing equation and the pendulum equation”. In: *Applied Mathematical Sciences* 8.176, pp. 8781–8789.