

A quantitative Hilbert's basis theorem and the constructive Krull dimension

Ryota Kuroki

The University of Tokyo

17 October 2025

These slides are available at



About me

Ryota Kuroki

Graduate student in mathematics at the University of Tokyo.

- My supervisor: Ryu Hasegawa
- Research interest: constructive algebra

These slides are available at



What is constructive algebra?

Constructive algebra: Algebra without nonconstructive principles (e.g., excluded middle, Zorn's lemma, ...).

Constructive proofs have computational content. They can be regarded as programs for proof assistants.

Proof of $\exists n \in \mathbb{N}. \varphi(n) \rightsquigarrow$ Algorithm to compute n s.t. $\varphi(n)$

- 1 α -Noetherian rings
- 2 Quantitative Hilbert's basis theorem and Krull dimension
- 3 Proof
- 4 Summary

Noetherian rings (non-constructive)

Definition 1

A ring A is *Noetherian* if

$$\forall I_0 \leq I_1 \leq \dots. \quad \exists n. \quad I_n = I_{n+1} = \dots.$$

Example 1

- ❶ All fields are Noetherian.
- ❷ \mathbb{Z} is Noetherian.
- ❸ $\mathbb{Z}[X_0, X_1, \dots]$ is not Noetherian.

$$0_{\mathbb{Z}} < \langle 12 \rangle < \langle 6 \rangle < \langle 3 \rangle < \langle 1 \rangle = \mathbb{Z}$$

$$0_{\mathbb{Z}[X_0, X_1, \dots]} < \langle X_0 \rangle < \langle X_0, X_1 \rangle < \langle X_0, X_1, X_2 \rangle < \dots$$

Noetherian rings (Richman–Seidenberg)

Problems:

- There are mysterious ideals like $\{x \in \mathbb{Z} : (x = 0) \vee \varphi\}$.
- $\langle 2 \rangle \leq \langle 2 \rangle \leq \dots$
(is it $\langle 2 \rangle$ forever? or will it be \mathbb{Z} at somewhere?).

There are several constructive definition of Noetherianity (Buriola, Schuster, and Blechschmidt [2023]).

Definition by Richman [1974] and Seidenberg [1974]:

Definition 2

A ring A is *Noetherian* if

$$\forall I_0 \leq I_1 \leq \dots \text{ (f.g.) } \exists n. \quad I_n = I_{n+1}.$$

- If $I \leq \mathbb{Z}$ is f.g., we can compute $a \in \mathbb{Z}$ s.t. $I = \langle a \rangle$.
- We don't have to wait until I_n stabilizes.

Noetherian rings (Jacobsson–Löfwall)

Generalized inductive definition by Jacobsson and Löfwall [1991]:


Definition 3

An ideal $I \leq A$ is *blocked* if

$$\forall x \in A. (x \notin I) \rightarrow (I + \langle x \rangle \text{ is blocked}).$$

A ring A is *Noetherian* if $0 \leq A$ is blocked.

(I prefer $\forall x \in A. (x \in I) \vee (I + \langle x \rangle \text{ is blocked}).$)



Noetherian rings (Coquand–Persson)

Generalized inductive definition by Coquand and Persson [1999]:


Definition 4

A list $[x_0, \dots, x_{n-1}] \in \text{List } A$ is *good* if $\exists k. x_k \in \langle x_0, \dots, x_{k-1} \rangle$.

A list $\sigma \in \text{List } A$ is *barred by good* if

$$(\sigma \text{ is good}) \vee (\forall x \in A. \sigma.x \text{ is barred by good}).$$

A ring is *Noetherian* if $[]$ is barred by good.



α -Noetherian rings


Definition 5

A list $[x_0, \dots, x_{n-1}]$ is (-1) -good (or simply *good*) if
 $(n \geq 1) \wedge x_{n-1} \in \langle x_0, \dots, x_{n-2} \rangle$.

A list $\sigma \in \text{List } A$ is α -good ($\alpha \in \text{Ord}$) if

$$\forall x \in A. \exists \beta \in [-1, \alpha). \sigma.x \text{ is } \beta\text{-good}.$$

A ring is α -Noetherian if $[]$ is α -good.



Classically, the notion of α -Noetherian ring is introduced by Gulliksen [1973] as the length of Noetherian modules.

Examples of α -Noetherian rings

Example 2

- 1 Discrete fields $(\forall x. (x = 0) \vee (x \in K^\times))$ are 1-Noetherian
- 2 \mathbb{Z} is ω -Noetherian.

More generally, we can define α -Euclidean rings and prove that they are α -Noetherian. (Classically, the notion of α -Euclidean ring is essentially introduced by Motzkin [1949].)

Definition 6

- 1 $x \in A$ is called (-1) -Euclidean if $x = 0$.
- 2 $x \in A$ is called α -Euclidean if for every $y \in A$, there exist $\beta \in [-1, \alpha)$ and β -Euclidean element $z \in A$ s.t. $z - y \in \langle x \rangle$.
- 3 A ring A is called α -Euclidean if for every $x \in A$, there exists $\beta \in [-1, \alpha)$ s.t. x is β -Euclidean.

Hilbert's basis theorem (HBT)

Theorem 7 (Classical HBT)

In classical mathematics, if A is Noetherian, then so is $A[X]$.

Theorem 8 (Coquand and Persson [1999])

If A is Coquand–Persson Noetherian, then so is $A[X]$.

There are also Richman–Seidenberg HBT and Jacobsson–Löfwall HBT, but those are theorems about

Noetherianity + (some conditions like coherence),
which are classically equivalent to Noetherianity.

Quantitative Hilbert's basis theorem (QHBT)

Theorem 9 (Kuroki [2025])

If A is α -Noetherian, then $A[X]$ is $(\omega \otimes \alpha)$ -Noetherian.

Classically, this is proved by Brookfield [2003].

Corollary 1

- 1 *If K is a discrete field, $K[X_0, \dots, X_{n-1}]$ is ω^n -Noetherian.*
- 2 *$\mathbb{Z}[X_0, \dots, X_{n-1}]$ is ω^{1+n} -Noetherian.*

Krull dimension (non-constructive)

Definition 10

We write $\text{Kdim } A < n$ if

$$\forall \mathfrak{p}_0 \leq \dots \leq \mathfrak{p}_n. \exists k. \mathfrak{p}_k = \mathfrak{p}_{k+1}$$

Krull dimension (constructive)

Lombardi [2002] has found the following characterization:

Definition 11

We write $\text{Kdim } A < n$ if for every $x_0, \dots, x_{n-1} \in A$, there exists $e_0, \dots, e_{n-1} \geq 0$ such that

$$x_0^{e_0} \cdots x_{n-1}^{e_{n-1}} \in \langle x_0^{e_0+1}, x_0^{e_0} x_1^{e_1+1}, \dots, x_0^{e_0} \cdots x_{n-2}^{e_{n-2}} x_{n-1}^{e_{n-1}+1} \rangle.$$

As noted by [Lombardi, 2002, Proposition 5.2], this definition is closely related to the lexicographic order. For more results in this direction, see Kemper and Trung [2014], Kemper and Yengui [2020].

Example 3

- 1 $\text{Kdim } K < 1$ for a discrete field K .
- 2 $\text{Kdim } \mathbb{Z} < 2$.

α -Noetherianity and Krull dimension (1/2)

Theorem 12 (Kuroki [2025])

Let $f : [0, \alpha) \rightarrow A$ be a function. If A is β -Noetherian for some $\beta < \alpha$, there exist $m \in \mathbb{N}$ and a strictly decreasing sequence $\alpha_0, \dots, \alpha_{m-1} \in [0, \beta]$ s.t. $[f(\alpha_0), \dots, f(\alpha_{m-1})]$ is good.

Proof.

Let $\alpha_0 := \beta$. Then $[f(\alpha_0)]$ is α_1 -good for some $\alpha_1 \in [-1, \alpha_0)$.

- ① If $\alpha_1 = -1$, then $[f(\alpha_0)]$ is good.
- ② If $\alpha_1 \in [0, \alpha_0)$, then $[f(\alpha_0), f(\alpha_1)]$ is α_2 -good for some $\alpha_2 \in [-1, \alpha_1) \dots$



α -Noetherianity and Krull dimension (2/2)

Theorem 13 (Kuroki [2025], Classically proved by Gulliksen [1973])

If A is α -Noetherian for some $\alpha < \omega^n$, then $\text{Kdim } A < n$.

Proof.

Define $f : \omega^n \rightarrow A$ by $f(e_{n-1}, \dots, e_1, e_0) := x_0^{e_0} \cdots x_{n-1}^{e_{n-1}}$. □


Corollary 2 (Lombardi [2002], Lombardi and Quitté [2015])

- ① *If K is a discrete field, $\text{Kdim } K[X_0, \dots, X_{n-1}] < 1 + n$.*
- ② $\text{Kdim } \mathbb{Z}[X_0, \dots, X_{n-1}] < 2 + n$.


Transfinite Chomp (1/5)

To prove QHBT, we use a game called (*transfinite*) *chomp* (Huddleston and Shurman [2002]).


$(\alpha \times \beta)$ -chomp:




Transfinite Chomp (2/5)



Transfinite Chomp (3/5)



Transfinite Chomp (4/5)



Transfinite Chomp (5/5)

The game ends in a finite number of steps. (Dickson's lemma)
Sketch of a proof (Huddleston and Shurman [2002]): We can assign an ordinal *size* P to each position P of the game. Every time you remove circles, the size decreases. □

The size of the initial position is $\alpha \otimes \beta$ (Hessenberg natural product).

Summary and future work

The notion of α -Noetherian ring works well with Krull dimension.

Future work: Constructive dimension theory of Noetherian rings

References I

- Gary Brookfield. The length of Noetherian polynomial rings. *Comm. Algebra*, 31(11): 5591–5607, 2003. ISSN 0092-7872,1532-4125. doi: 10.1081/AGB-120023976. URL <https://doi.org/10.1081/AGB-120023976>.
- Gabriele Buriola, Peter Schuster, and Ingo Blechschmidt. A constructive picture of Noetherian conditions and well quasi-orders. In *Unity of logic and computation*, volume 13967 of *Lecture Notes in Comput. Sci.*, pages 50–62. Springer, Cham, 2023. ISBN 978-3-031-36977-3; 978-3-031-36978-0. doi: 10.1007/978-3-031-36978-0_5. URL https://doi.org/10.1007/978-3-031-36978-0_5.
- Thierry Coquand and Henrik Persson. Gröbner bases in type theory. In *Types for proofs and programs (Irvine, 1998)*, volume 1657 of *Lecture Notes in Comput. Sci.*, pages 33–46. Springer, Berlin, 1999. ISBN 3-540-66537-4. doi: 10.1007/3-540-48167-2_3. URL https://doi.org/10.1007/3-540-48167-2_3.
- Tor H. Gulliksen. A theory of length for Noetherian modules. *J. Pure Appl. Algebra*, 3: 159–170, 1973. ISSN 0022-4049,1873-1376. doi: 10.1016/0022-4049(73)90030-3. URL [https://doi.org/10.1016/0022-4049\(73\)90030-3](https://doi.org/10.1016/0022-4049(73)90030-3).
- Scott Huddleston and Jerry Shurman. Transfinite Chomp. In *More games of no chance (Berkeley, CA, 2000)*, volume 42 of *Math. Sci. Res. Inst. Publ.*, pages 183–212. Cambridge Univ. Press, Cambridge, 2002. ISBN 0-521-80832-4.

References II

- Carl Jacobsson and Clas Löfwall. Standard bases for general coefficient rings and a new constructive proof of Hilbert's basis theorem. *J. Symbolic Comput.*, 12(3):337–371, 1991. ISSN 0747-7171,1095-855X. doi: 10.1016/S0747-7171(08)80154-X. URL [https://doi.org/10.1016/S0747-7171\(08\)80154-X](https://doi.org/10.1016/S0747-7171(08)80154-X).
- Gregor Kemper and Ngo Viet Trung. Krull dimension and monomial orders. *J. Algebra*, 399:782–800, 2014. ISSN 0021-8693,1090-266X. doi: 10.1016/j.jalgebra.2013.10.005. URL <https://doi.org/10.1016/j.jalgebra.2013.10.005>.
- Gregor Kemper and Ihsen Yengui. Valuative dimension and monomial orders. *J. Algebra*, 557:278–288, 2020. ISSN 0021-8693,1090-266X. doi: 10.1016/j.jalgebra.2020.04.017. URL <https://doi.org/10.1016/j.jalgebra.2020.04.017>.
- Ryota Kuroki. A quantitative Hilbert's basis theorem and the constructive Krull dimension, 2025. URL <https://arxiv.org/abs/2509.00363>.
- Henri Lombardi. Dimension de Krull, Nullstellensätze et évaluation dynamique. *Math. Z.*, 242(1):23–46, 2002. ISSN 0025-5874,1432-1823. doi: 10.1007/s002090100305. URL <https://doi.org/10.1007/s002090100305>.
- Henri Lombardi and Claude Quitté. *Commutative algebra: constructive methods*, volume 20 of *Algebra and Applications*. Springer, Dordrecht, revised edition, 2015. ISBN 978-94-017-9944-7. doi: 10.1007/978-94-017-9944-7. Finite projective modules, Translated from the French by Tania K. Roblot.

References III

- Theodore Motzkin. The Euclidean algorithm. *Bull. Amer. Math. Soc.*, 55:1142–1146, 1949. ISSN 0002-9904. doi: 10.1090/S0002-9904-1949-09344-8. URL <https://doi.org/10.1090/S0002-9904-1949-09344-8>.
- Fred Richman. Constructive aspects of Noetherian rings. *Proc. Amer. Math. Soc.*, 44: 436–441, 1974. ISSN 0002-9939,1088-6826. doi: 10.2307/2040452. URL <https://doi.org/10.2307/2040452>.
- Abraham Seidenberg. What is Noetherian? *Rend. Sem. Mat. Fis. Milano*, 44:55–61, 1974. ISSN 0370-7377. doi: 10.1007/BF02925651. URL <https://doi.org/10.1007/BF02925651>.