Maplewood General Hospital

400 Orchard Street, New Haven, CT 06511 | Main line (203) 555-0100

Hospital Discharge Summary and Patient Instructions

Patient	Maria Garcia	**MRN**	MGH-0051877
Date of birth	07/02/1969	**Unit**	Medicine 4 West
Admitted	10/13/2026	**Discharged**	10/17/2026
Attending	Priya Natarajan, MD (Hospital Medicine)	**Preferred language**	Spanish (interpreter used)

Why you were in the hospital

Diagnosis: Community-acquired pneumonia (lung infection) of the right lower lobe. Type 2 diabetes with high blood sugar (hemoglobin A1c 10.4%).

You were treated with IV antibiotics and your breathing improved. Your blood sugar was high, so we started a long-acting insulin.

Hospital course (for your care team)

57F with T2DM (A1c 10.4%), HTN, presenting with 4 days of productive cough, fever to 102.6 F and pleuritic chest pain. CXR with RLL consolidation. Initially on 3 L NC, weaned to room air by HD3. Treated with IV ceftriaxone and azithromycin, transitioned to PO amoxicillin-clavulanate to complete a 7-day total course. Blood cultures no growth at 48 hours. Hyperglycemia to the 300s managed with basal-bolus insulin; discharged on basal insulin plus metformin. Spanish interpreter used for all teaching.

Your medications

START taking these medications

Amoxicillin-clavulanate (Augmentin) 875 mg-125 mg tablet. Take 1 tablet by mouth twice daily with food for 5 days. Last dose: evening of 10/21/2026. Finish all of it even if you feel better.

Insulin glargine (Lantus SoloStar) 100 units/mL pen. Inject 18 units under the skin once daily at bedtime. Purpose: lowers blood sugar overnight and through the day.

Acetaminophen (Tylenol) 500 mg tablet. Take 2 tablets (1,000 mg) by mouth every 8 hours as needed for fever or pain. Do not take more than 3,000 mg in 24 hours.

Benzonatate 100 mg capsule. Take 1 capsule by mouth three times daily as needed for cough. Swallow whole; do not chew.

CONTINUE taking these medications

Metformin 1,000 mg tablet. Take 1 tablet by mouth twice daily with breakfast and dinner. This was paused in the hospital; restart it at home.

Blood glucose meter and test strips. Check your blood sugar before breakfast and before dinner every day and write the numbers down.

Diet and activity

Rest, and slowly increase your activity. Drink plenty of water unless another doctor has told you to limit fluids.

Low blood sugar: if your sugar is below 70, eat or drink 15 grams of fast sugar (4 ounces of juice or 4 glucose tablets), wait 15 minutes, and check again.

Follow-up appointments and tests

Primary care, Dr. Laura Chen: Friday 10/23/2026 at 9:15 AM, Maplewood Primary Care, Suite 110. Bring your blood sugar log.

Chest X-ray: in 6 weeks to make sure the pneumonia has cleared. The order is placed; Radiology will call you to schedule.

Diabetes educator: call (203) 555-0119 to schedule an insulin teaching visit this week.

When to get help

Call Dr. Chen's office at (203) 555-0110 if: you still have a fever above 101 F after 3 days of antibiotics; your blood sugar is above 300 two times in a row; you get a rash, or watery diarrhea more than 3 times a day.

Call 911 or go to the emergency room if: you have trouble breathing; your lips or face turn blue or gray; you become confused or very sleepy; your blood sugar is below 54 and does not come up after treatment, or you pass out.