

SUMMARY OF PRODUCT CHARACTERISTICS

SPUTNIK LIGHT

Gam-COVID-Vac

Vaccine for the prevention of coronavirus infection caused by the SARS-CoV-2 virus

Trade Name: Sputnik Light vaccine

International nonproprietary or generic name: Sputnik Light vaccine against COVID-19

Dosage form: solution for intramuscular injection

Composition per 1 dose (0.5 ml):

Active substance: recombinant adenovirus serotype 26, containing the SARS-CoV-2 protein S gene, in the amount of $1.0 \pm 0.5 \times 10^{11}$ particles.

Excipients: tris(hydroxymethyl)aminomethane – 1.21 mg, sodium chloride – 2.19 mg, sucrose – 25.0 mg, magnesium chloride hexahydrate – 102.0 µg, EDTA disodium salt dihydrate – 19.0 µg, polysorbate 80 – 250 µg, ethanol 95 % – 2.5 µl, water for injection Q.s. to 0.5 ml.

Description

Frozen preparation

Frozen solution – a dense solidified whitish mass. After thawing: a homogeneous colorless or with a yellowish shade slightly opalescent solution.

Liquid preparation

Homogeneous colorless or with a yellowish shade slightly opalescent solution.

Characteristics

The vaccine was obtained by a biotechnological method, which doesn't imply the SARS-CoV-2 virus pathogenic to humans. The composition of the product includes a recombinant adenovirus vector based on human adenovirus serotype 26, carrying the SARS-CoV-2 virus

S protein gene.

Pharmacotherapeutic group:

Immunobiological medicinal products-vaccine (MIBP)

ATC Code: J07B

Pharmacological properties

The vaccine induces the development of humoral and cellular immunity against coronavirus infection caused by the SARS-CoV-2 virus.

Immunological efficacy

The immunological properties and safety of the vaccine were studied in various clinical studies in adult volunteers of both sexes aged over 18 years.

An interim analysis of immunogenicity showed that the vaccine develops a humoral immune response in volunteers. On day 28 the seroconversion was detected in 96.88 % of healthy volunteers.

Patients with pre-immunity to coronavirus experienced a pronounced increase in antibodies titer on day 10 after vaccination, which may indicate the possibility of using the product for vaccination of the previously exposed to COVID-19 after a decrease in antibodies titer to prevent recurrence of the disease.

Following single immunization, 100 % of healthy volunteers (all 30 people whose cellular immunity indicators were studied) experienced a cellular immune response against the S-protein of the SARS-CoV-2.

The protective titer of the antibodies is currently unknown. The duration of the protection is unknown.

Therapeutic indications

Prevention of a new coronavirus infection (COVID-19) in adults aged 18–60.

Contraindications

hypersensitivity to any component of the vaccine or a vaccine containing similar components;
preexisting severe allergic reactions;

acute infectious and non-infectious diseases, chronic diseases aggravated – the vaccination is carried out 2–4 weeks after recovery or remission.

In case of mild acute respiratory viral infections, acute infectious diseases of the gastrointestinal tract, the vaccination is carried out after establishing of average body temperature;

pregnancy and breastfeeding;

people under age 18 (due to a lack of data on efficacy and safety).

With caution

Use the vaccine with caution in chronic diseases of the liver and kidneys, endocrine diseases (severe thyroid disorders and diabetes mellitus in the decompensation stage), severe diseases of the hematopoietic system, epilepsy, and other diseases of the central nervous system, acute coronary syndrome and acute cerebral circulation disorders, myocarditis, endocarditis, pericarditis.

Due to a lack of information, vaccination may increase the risk for the following groups of patients: with autoimmune diseases (stimulation of the immune system may lead to aggravation of the disease, especially patients with autoimmune pathology, which tends to develop severe and life-threatening conditions, should be treated with caution); with malignant neoplasms.

The decision about getting a vaccine should be based on an assessment of the benefit-risk ratio in each specific situation.

Use during pregnancy and breastfeeding

The product is contraindicated during pregnancy and breastfeeding since its effectiveness and safety during this period have not been studied.

Posology and administration

The vaccine is intended for intramuscular administration only. Intravenous administration of the product is strictly prohibited. The vaccine is injected into the deltoid muscle (the upper third of the outer surface of the arm). If it is impossible to inject into the deltoid muscle, the product is injected into the lateral broad muscle of the thigh.

After the vaccine administration, the patient should be under the supervision of medical personnel for 30 minutes.

Frozen preparation

Before vaccination, the vial with the product is removed from the freezer and kept at room temperature until it is completely thawed. The remaining ice inclusions are not allowed! Wipe the vial, ampoule, or syringe from the outside with an alcohol wipe to remove moisture. Mix the contents by tilting carefully. Do not shake the vial.

Remove the protective plastic cap from the vial and treat the rubber stopper with an alcohol wipe. The ampoule is opened along the ring and/or the break point. Using a disposable syringe with a needle, a dose of 0.5 ml is selected for administration to the patient from an ampoule or vial.

Attention

The dose administered to patient should not exceed 0.5 ml.

The product in syringes is ready for use without additional manipulations.

It is not allowed to store the thawed drug for more than 30 minutes! Repeated freezing is not allowed.

Refreezing of the vial or ampoule with the solution is not allowed! The total storage of the diluted solution at room temperature should not exceed 2 hours. An unused vaccine must be destroyed.

If subsequent injections are postponed for any reason, it is allowed to store an opened or closed vial for no more than 2 hours at room temperature.

If subsequent injections are postponed for any reason, it is allowed to store a 3 ml opened vial for no more than 2 hours at room temperature.

Storage of the thawed product in 0.5 ml vials for more than 30 minutes is not allowed.

Refreezing is not allowed.

The drug with defects in the closure system and/or defective labeling of the vial or ampoule, with changes in the physical properties of the solution (turbidity, staining), improper storage and/or expired shelf life is unsuitable for use.

Side effect

Adverse events characteristic of the use of the vaccine, identified in clinical studies, as well as studies of other vaccines based on a similar technological platform, are mainly mild or moderate, may develop in the first or second day after vaccination and resolve within 3 following days. Short-term general (short-term flu syndrome characterized by chills, fever, arthralgia, myalgia, asthenia, general malaise, headache) and local (injection site tenderness, hyperemia, swelling) reactions may be common and very common. It is recommended to prescribe non-steroidal anti-inflammatory drugs (NSAIDs) in the event of increase in temperature after vaccination and antihistamines with a pronounced local reaction.

The following adverse events have been reported according to the System Organ Class and frequency. The frequency is defined as follows: very common ($\geq 1/10$), common ($\geq 1/100$ and $<1/10$), uncommon ($\geq 1/1000$ and $<1/100$), rare ($\geq 1/10\ 000$ and $<1/1000$), very rare ($<1/10\ 000$, including individual cases). The frequency categories were established basing on clinical studies of the product.

Common: nausea, dyspepsia, decreased appetite, very rare: regional lymph nodes enlargement. Some patients may develop allergic reactions, a short-term increase in serum hepatic transaminases, creatinine, and creatine phosphokinase.

As a part of the conducted clinical studies on the safety, tolerability, and immunogenicity of

the medicinal product Sputnik Light after vaccination, the following AE were registered:

“General disorders and administration site condition”: hyperthermia, pain, oedema, vaccination site pruritus, asthenia, pain, malaise, pyrexia, vaccination site warmth, decreased appetite. Frequency – very common and common.

“Respiratory, thoracic, and mediastinal disorders”: oropharyngeal pain, nasal congestion, throat clearing, rhinorrhea. Frequency – common.

“Nervous system disorders”: headache, asthenia – common; dizziness, faint – rare.

“Gastrointestinal disorders”: nausea, vomiting, dyspepsia – common.

“Investigations”: multidirectional deviations of immune status indicators: T-lymphocyte count increased, lymphocyte percentage increased, natural killer cell count decreased, CD4 lymphocytes increased, CD4 lymphocytes decreased, B lymphocyte count increased, B lymphocyte count decreased, natural killer cell count increased, CD8 lymphocytes increased, blood immunoglobulin E (IgE) increased, CD4/CD8 ratio increased, CD4/CD8 ratio decreased, blood immunoglobulin A (IgA) increased, CD8 lymphocytes percentage decreased. Complete blood count abnormal: lymphocytes percentage increased, hematocrit decreased, lymphocyte count increased, red blood cell sedimentation rate increased, white blood cell counts increased, monocyte count increased, platelet count increased, neutrophil count decreased, platelet count decreased. Urine analysis abnormal: red blood cells urine.

Most AE were recovered/resolved. Laboratory abnormalities were not of clinical significance (they did not require additional diagnostic procedures and administration of therapy).

Overdose

No overdose cases have been reported.

Given that the prescription of the product is allowed only for medical and preventive treatment institutions, and the vaccination itself is carried out only by qualified medical personnel, the risk of overdose is extremely low.

However, it can be assumed that with an accidental overdose, the development of the above mentioned toxic and toxic-allergic reactions to a more severe degree is possible. There are no specific antidotes to the medicinal product.

In this case therapeutic measures will include symptomatic therapy in accordance with the indications (corticosteroids, antipyretics/NSAIDs, and desensitizers). Dosing regimen of medicinal products should be selected according to the recommendations for the use and strengths of this product.

Interaction with other medicinal products

No studies have been performed.

Special warnings

Patients receiving immunosuppressive therapy and patients with immunodeficiency may not develop a sufficient immune response. Therefore, administration of medicinal products that inhibit the function of the immune system is contraindicated for at least 1 month before and after vaccination due to the risk of reducing immunogenicity. Only the attending physician can decide on the possibility of immunosuppressive therapy discontinuation!

Like other vaccines, not all vaccinated people may have a protective immune response.

Effects on ability to drive and use machines

No studies on the effects of the vaccine on ability to drive and use potentially dangerous machines have been performed.

Pharmaceutical form

Solution for intramuscular injection, 0.5 ml/dose.

3,0 ml (5 doses) of the medicinal product – in vials, capped with rubber stoppers, crimped with aluminum-plastic tamper evident caps.

1 vial of the medicinal product is placed into a carton pack with a foam holder made of polyurethane foam along with the package insert.

Storage conditions

Store in a light-proof place at a temperature of -18 °C or below. Refreezing is not allowed.

Keep out of reach of children.

Transportation conditions

Transportation of the medicinal product at temperature not exceeding - 18 °C.

Shelf life

6 months. Do not use after the expiry date.

Prescription category: For medical and preventive treatment institutions. For use in emergency only.

7. MARKETING AUTHORIZATION

M/s Dr Reddy's Laboratories Ltd., Global Distribution Centre, Survey No. 41, Bachupally (V),
Bachupally (M), Medchal - Malkajgiri(Dist.), Hyderabad – 500090, Telangana, INDIA™

Trademark under registration

Manufactured by:

M/s Hetero Biopharma Limited, Sy. No. 458 (part), TSIIC-Formulation SEZ, Polepally Village,
Jadcherla Mandal, Mahaboobnagar District-509 301, Telangana State, India.

8. MARKETING AUTHORISATION NUMBER (S)

MF/BIO/22/000024

9. DATE OF FIRST AUTHORISATION/ RENEWAL OF THE AUTHORISATION

20-04-2022