


Photochemistry

- ▶ initiated by absorption of electromagnetic radiation
- ▶ most important : processes that capture solar energy
 - ▶ Some reactions lead to the heating of the atmosphere during the daytime by absorption of uv radiation.
 - ▶ Others include absorption of visible radiation during photosynthesis

Jablonski diagram



ground vibrational states correctly located vertically but other vibrational states are shown schematically
(IC: internal conversion; ISC: intersystem crossing)

primary process : products formed directly from excited state of reactant







Ex. : fluorescence and cis–trans photoisomerization of retinal

secondary process : products originate from intermediates formed directly from excited state of reactant,

Ex. : oxidative processes initiated by oxygen atom formed by ozone photodissociation

Competing with formation of photochemical products are numerous primary photophysical processes that can deactivate excited state

∴ it is important to consider timescales of formation and decay of excited states before describing mechanisms



Primary absorption	$S + h\nu \rightarrow S^*$
Excited-state absorption	$S^* + h\nu \rightarrow S^{**}$
	$T^* + h\nu \rightarrow T^{**}$
Fluorescence	$S^* \rightarrow S + h\nu$
Stimulated emission	$S^* + h\nu \rightarrow S + 2 h\nu$
Intersystem crossing (ISC)	$S^* \rightarrow T^*$
Phosphorescence	$T^* \rightarrow S + h\nu$
Internal conversion (IC)	$S^* \rightarrow S$
Collision-induced emission	$S^* + M \rightarrow S + M + h\nu$
Collisional deactivation	$S^* + M \rightarrow S + M$
	$T^* + M \rightarrow S + M$
Electronic energy transfer:	
Singlet-singlet	$S^* + S \rightarrow S + S^*$
Triplet-triplet	$T^* + T \rightarrow T + T^*$
Excimer formation	$S^* + S \rightarrow (SS)^*$
Energy pooling	
Singlet-singlet	$S^* + S^* \rightarrow S^{**} + S$
Triplet-triplet	$T^* + T^* \rightarrow S^{**} + S$

Interplay of reaction rates \leftrightarrow excited state lifetime is important in determining kinetic feasibility of photochemical process

Interplay of reaction rates \leftrightarrow excited state lifetime is important in determining kinetic feasibility of photochemical process

Electronic transitions: $10^{-15}\text{s} \implies k_{\text{1st-order}} < 10^{16}\text{s}^{-1}$

Interplay of reaction rates \leftrightarrow excited state lifetime is important in determining kinetic feasibility of photochemical process

Electronic transitions: $10^{-15}\text{s} \Rightarrow k_{\text{1st-order}} < 10^{16}\text{s}^{-1}$

$\tau_{\text{Fluorescence}} \sim 10^{-12} - 10^{-6}\text{s} > \tau_{\text{absorption}}$

Interplay of reaction rates \leftrightarrow excited state lifetime is important in determining kinetic feasibility of photochemical process

Electronic transitions: $10^{-15}\text{s} \implies k_{\text{1st-order}} < 10^{16}\text{s}^{-1}$

$\tau_{\text{Fluorescence}} \sim 10^{-12} - 10^{-6}\text{s} > \tau_{\text{absorption}}$

excited singlet state can initiate fast photochemical reactions in $10^{-15} - 10^{-12}\text{s}$

Interplay of reaction rates \leftrightarrow excited state lifetime is important in determining kinetic feasibility of photochemical process

Electronic transitions: $10^{-15}\text{s} \implies k_{\text{1st-order}} < 10^{16}\text{s}^{-1}$

$\tau_{\text{Fluorescence}} \sim 10^{-12} - 10^{-6}\text{s} > \tau_{\text{absorption}}$

excited singlet state can initiate fast photochemical reactions in $10^{-15} - 10^{-12}\text{s}$
primary process of vision is complete in 200fs

Interplay of reaction rates \leftrightarrow excited state lifetime is important in determining kinetic feasibility of photochemical process

Electronic transitions: $10^{-15}\text{s} \Rightarrow k_{\text{1st-order}} < 10^{16}\text{s}^{-1}$

$\tau_{\text{Fluorescence}} \sim 10^{-12} - 10^{-6}\text{s} > \tau_{\text{absorption}}$

excited singlet state can initiate fast photochemical reactions in $10^{-15} - 10^{-12}\text{s}$

primary process of vision is complete in 200fs

for large organic molecules: ISC $\sim 10^{-12} - 10^{-4}\text{s}$ and phosphorescence $\sim 10^{-6} - 10^{-1}\text{s}$

Interplay of reaction rates \leftrightarrow excited state lifetime is important in determining kinetic feasibility of photochemical process

Electronic transitions: $10^{-15}\text{s} \Rightarrow k_{\text{1st-order}} < 10^{16}\text{s}^{-1}$

$\tau_{\text{Fluorescence}} \sim 10^{-12} - 10^{-6}\text{s} > \tau_{\text{absorption}}$

excited singlet state can initiate fast photochemical reactions in $10^{-15} - 10^{-12}\text{s}$

primary process of vision is complete in 200fs

for large organic molecules: ISC $\sim 10^{-12} - 10^{-4}\text{s}$ and phosphorescence $\sim 10^{-6} - 10^{-1}\text{s}$

\Rightarrow excited triplet states are important photochemically; such species can undergo large number of collisions with other reactants before deactivation

Consider

molecular photochemical reaction, $k = 1.7 \times 10^4 \text{ s}^{-1}$

Consider

molecular photochemical reaction, $k = 1.7 \times 10^4 \text{ s}^{-1}$

relaxation time $\tau = \frac{1}{1.7 \times 10^4} \text{ s}^{-1} = 59 \mu\text{s}$

Consider

molecular photochemical reaction, $k = 1.7 \times 10^4 \text{ s}^{-1}$

relaxation time $\tau = \frac{1}{1.7 \times 10^4} \text{ s}^{-1} = 59 \mu\text{s}$

reactant with observed fluorescence lifetime = 1 ns

Consider

molecular photochemical reaction, $k = 1.7 \times 10^4 \text{ s}^{-1}$

relaxation time $\tau = \frac{1}{1.7 \times 10^4} \text{ s}^{-1} = 59 \mu\text{s}$

reactant with observed fluorescence lifetime = 1 ns

observed phosphorescence lifetime = 1 ms

Consider

molecular photochemical reaction, $k = 1.7 \times 10^4 \text{ s}^{-1}$

relaxation time $\tau = \frac{1}{1.7 \times 10^4} \text{ s}^{-1} = 59 \text{ } \mu\text{s}$

reactant with observed fluorescence lifetime = 1 ns

observed phosphorescence lifetime = 1 ms

Excited singlet state is too short-lived and not expected to be major source of product in this reaction.

Consider

molecular photochemical reaction, $k = 1.7 \times 10^4 \text{ s}^{-1}$

relaxation time $\tau = \frac{1}{1.7 \times 10^4} \text{ s}^{-1} = 59 \text{ } \mu\text{s}$

reactant with observed fluorescence lifetime = 1 ns

observed phosphorescence lifetime = 1 ms

Excited singlet state is too short-lived and not expected to be major source of product in this reaction.

Excited triplet state is a good candidate for a precursor

Rates of deactivation of excited state : $\left\{ \begin{array}{l} \text{radiative} \\ \text{non-radiative} \\ \text{chemical} \end{array} \right.$

Rates of deactivation of excited state : $\left\{ \begin{array}{l} \text{radiative} \\ \text{non-radiative} \\ \text{chemical} \end{array} \right.$
determine yield of product in a photochemical reaction

Rates of deactivation of excited state : $\left\{ \begin{array}{l} \text{radiative} \\ \text{non-radiative} \\ \text{chemical} \end{array} \right.$

determine yield of product in a photochemical reaction

Primary quantum yield,

$$\phi = \frac{\# \text{photo-events (physical/chemical)}}{\# \text{photons absorbed in same interval}} =$$

Rates of deactivation of excited state : $\left\{ \begin{array}{l} \text{radiative} \\ \text{non-radiative} \\ \text{chemical} \end{array} \right.$

determine yield of product in a photochemical reaction

Primary quantum yield,

$$\phi = \frac{\# \text{photo-events (physical/chemical)}}{\# \text{photons absorbed in same interval}} = \frac{\text{rate of process(es)}}{\text{intensity of light absorbed (rate of photon absorption)}} = \frac{\nu}{I_{\text{abs}}}$$

Rates of deactivation of excited state : $\left\{ \begin{array}{l} \text{radiative} \\ \text{non-radiative} \\ \text{chemical} \end{array} \right.$

determine yield of product in a photochemical reaction

Primary quantum yield,

$$\phi = \frac{\text{\#photo-events (physical/chemical)}}{\text{\#photons absorbed in same interval}} =$$

$$= \frac{\text{rate of process(es)}}{\text{intensity of light absorbed (rate of photon absorption)}} = \frac{\nu}{I_{\text{abs}}}$$

total #molecules deactivated (radiative/non-radiative/photo-reactive)

Rates of deactivation of excited state : $\left\{ \begin{array}{l} \text{radiative} \\ \text{non-radiative} \\ \text{chemical} \end{array} \right.$

determine yield of product in a photochemical reaction

Primary quantum yield,

$$\phi = \frac{\text{\#photo-events (physical/chemical)}}{\text{\#photons absorbed in same interval}} =$$

$$= \frac{\text{rate of process(es)}}{\text{intensity of light absorbed (rate of photon absorption)}} = \frac{\nu}{I_{\text{abs}}}$$

total #molecules deactivated (radiative/non-radiative/photo-reactive)

= #excited species produced by absorption of light

Rates of deactivation of excited state : $\left\{ \begin{array}{l} \text{radiative} \\ \text{non-radiative} \\ \text{chemical} \end{array} \right.$

determine yield of product in a photochemical reaction

Primary quantum yield,

$$\phi = \frac{\# \text{photo-events (physical/chemical)}}{\# \text{photons absorbed in same interval}} =$$

$$= \frac{\text{rate of process(es)}}{\text{intensity of light absorbed (rate of photon absorption)}} = \frac{\nu}{I_{\text{abs}}}$$

total #molecules deactivated (radiative/non-radiative/photo-reactive)
= #excited species produced by absorption of light

$$\Rightarrow \sum_i \phi_i = \sum_i \frac{\nu_i}{I_{\text{abs}}} = 1$$

Rates of deactivation of excited state : $\left\{ \begin{array}{l} \text{radiative} \\ \text{non-radiative} \\ \text{chemical} \end{array} \right.$

determine yield of product in a photochemical reaction

Primary quantum yield,

$$\phi = \frac{\# \text{photo-events (physical/chemical)}}{\# \text{photons absorbed in same interval}} =$$

$$= \frac{\text{rate of process(es)}}{\text{intensity of light absorbed (rate of photon absorption)}} = \frac{\nu}{I_{\text{abs}}}$$

total #molecules deactivated (radiative/non-radiative/photo-reactive)
= #excited species produced by absorption of light

$$\Rightarrow \sum_i \phi_i = \sum_i \frac{\nu_i}{I_{\text{abs}}} = 1$$

$$\Rightarrow \phi_f + \phi_{IC} + \phi_{ISC} + \phi_p + \phi_R = 1; \quad \phi_i = \frac{\nu_i}{\sum_i \nu_i}$$

ex. : absorbing substance exposed to 490 nm light

from a $P = 100 \text{ W}$ source

for 2700 s

with $f = 60\%$ of incident light absorbed

0.344mol of absorbing substance decomposed

$$\phi = ?$$

ex. : absorbing substance exposed to 490 nm light

from a $P = 100 \text{ W}$ source

for 2700 s

with $f = 60\%$ of incident light absorbed

0.344mol of absorbing substance decomposed

$$\phi = ?$$

$$E_{abs} = fPt = N_{abs} \frac{hc}{\lambda}, \text{ or, } N_{abs} = \frac{fPt\lambda}{hc}$$

ex. : absorbing substance exposed to 490 nm light

from a $P = 100 \text{ W}$ source

for 2700 s

with $f = 60\%$ of incident light absorbed

0.344mol of absorbing substance decomposed

$$\phi = ?$$

$$E_{abs} = fPt = N_{abs} \frac{hc}{\lambda}, \text{ or, } N_{abs} = \frac{fPt\lambda}{hc}$$

$$\phi = \frac{N_{decomposed}}{N_{absorbed}} = \frac{(0.344 \times 6.023 \times 10^{23}) \times 6.626 \times 10^{-34} \times 2.998 \times 10^8}{0.6 \times 100 \times 2700 \times 490 \times 10^{-9}} = 0.52$$

Mechanism of decay of excited singlet states

Mechanism of decay of excited singlet states

Absorption: $S + h\nu_i \rightarrow S^*$; $\nu_{\text{abs}} = I_{\text{abs}}$

Mechanism of decay of excited singlet states

Absorption: $S + h\nu_i \rightarrow S^*$; $v_{\text{abs}} = I_{\text{abs}}$

Fluorescence: $S^* \rightarrow S + h\nu_f$; $v_f = k_f[S^*]$

Mechanism of decay of excited singlet states

Absorption: $S + h\nu_i \rightarrow S^*$; $v_{\text{abs}} = I_{\text{abs}}$

Fluorescence: $S^* \rightarrow S + h\nu_f$; $v_f = k_f[S^*]$

Internal conversion: $S^* \rightarrow S$; $v_{IC} = k_{IC}[S^*]$

Mechanism of decay of excited singlet states

Absorption: $S + h\nu_i \rightarrow S^*$; $v_{\text{abs}} = I_{\text{abs}}$

Fluorescence: $S^* \rightarrow S + h\nu_f$; $v_f = k_f[S^*]$

Internal conversion: $S^* \rightarrow S$; $v_{IC} = k_{IC}[S^*]$

Intersystem crossing: $S^* \rightarrow T^*$; $v_{ISC} = k_{ISC}[S^*]$

Mechanism of decay of excited singlet states

Absorption: $S + h\nu_i \rightarrow S^*$; $v_{\text{abs}} = I_{\text{abs}}$

Fluorescence: $S^* \rightarrow S + h\nu_f$; $v_f = k_f[S^*]$

Internal conversion: $S^* \rightarrow S$; $v_{IC} = k_{IC}[S^*]$

Intersystem crossing: $S^* \rightarrow T^*$; $v_{ISC} = k_{ISC}[S^*]$

rate of formation of $S^* = I_{\text{abs}}$

Mechanism of decay of excited singlet states

Absorption: $S + h\nu_i \rightarrow S^*$; $v_{\text{abs}} = I_{\text{abs}}$

Fluorescence: $S^* \rightarrow S + h\nu_f$; $v_f = k_f[S^*]$

Internal conversion: $S^* \rightarrow S$; $v_{IC} = k_{IC}[S^*]$

Intersystem crossing: $S^* \rightarrow T^*$; $v_{ISC} = k_{ISC}[S^*]$

rate of formation of $S^* = I_{\text{abs}}$

rate of decay of $S^* = -k_f[S^*] - k_{ISC}[S^*] - k_{IC}[S^*]$
 $= -(k_f + k_{ISC} + k_{IC})[S^*]$

Mechanism of decay of excited singlet states

Absorption: $S + h\nu_i \rightarrow S^*$; $v_{\text{abs}} = I_{\text{abs}}$

Fluorescence: $S^* \rightarrow S + h\nu_f$; $v_f = k_f[S^*]$

Internal conversion: $S^* \rightarrow S$; $v_{IC} = k_{IC}[S^*]$

Intersystem crossing: $S^* \rightarrow T^*$; $v_{ISC} = k_{ISC}[S^*]$

rate of formation of $S^* = I_{\text{abs}}$

rate of decay of $S^* = -k_f[S^*] - k_{ISC}[S^*] - k_{IC}[S^*]$
 $= -(k_f + k_{ISC} + k_{IC})[S^*]$

Assuming first order process:

$$[S^*]_t = [S^*]_0 e^{-\frac{t}{\tau_0}}; \quad \tau_0 = \frac{1}{k_f + k_{ISC} + k_{IC}}$$

rate of formation of $S^* = I_{\text{abs}}$

rate of formation of $S^* = I_{\text{abs}}$

rate of decay of $S^* = -k_f[S^*] - k_{\text{ISC}}[S^*] - k_{\text{IC}}[S^*]$

rate of formation of $S^* = I_{\text{abs}}$

rate of decay of $S^* = -k_f[S^*] - k_{\text{ISC}}[S^*] - k_{\text{IC}}[S^*]$

$$\frac{d[S^*]}{dt} = I_{\text{abs}} - (k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*] = 0 \text{ (S.S.)}$$

rate of formation of $S^* = I_{\text{abs}}$

rate of decay of $S^* = -k_f[S^*] - k_{\text{ISC}}[S^*] - k_{\text{IC}}[S^*]$

$$\frac{d[S^*]}{dt} = I_{\text{abs}} - (k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*] = 0 \text{ (S.S.)}$$

$$\text{or, } I_{\text{abs}} = (k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*]$$

rate of formation of $S^* = I_{\text{abs}}$

rate of decay of $S^* = -k_f[S^*] - k_{\text{ISC}}[S^*] - k_{\text{IC}}[S^*]$

$$\frac{d[S^*]}{dt} = I_{\text{abs}} - (k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*] = 0 \quad (\text{S.S.})$$

$$\text{or, } I_{\text{abs}} = (k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*]$$

$$\text{Quantum yield of fluorescence, } \phi_f = \frac{\nu_f}{I_{\text{abs}}} = \frac{k_f[S^*]}{(k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*]} = \frac{k_f}{k_f + k_{\text{ISC}} + k_{\text{IC}}}$$

rate of formation of $S^* = I_{\text{abs}}$

rate of decay of $S^* = -k_f[S^*] - k_{\text{ISC}}[S^*] - k_{\text{IC}}[S^*]$

$$\frac{d[S^*]}{dt} = I_{\text{abs}} - (k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*] = 0 \text{ (S.S.)}$$

$$\text{or, } I_{\text{abs}} = (k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*]$$

$$\text{Quantum yield of fluorescence, } \phi_f = \frac{\nu_f}{I_{\text{abs}}} = \frac{k_f[S^*]}{(k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*]} = \frac{k_f}{k_f + k_{\text{ISC}} + k_{\text{IC}}}$$

Fluorescence lifetime can be measured with pulsed laser technique.

Sample is excited with a short light pulse from a laser using wavelength at which S absorbs strongly

rate of formation of $S^* = I_{\text{abs}}$

rate of decay of $S^* = -k_f[S^*] - k_{\text{ISC}}[S^*] - k_{\text{IC}}[S^*]$

$$\frac{d[S^*]}{dt} = I_{\text{abs}} - (k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*] = 0 \text{ (S.S.)}$$

$$\text{or, } I_{\text{abs}} = (k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*]$$

$$\text{Quantum yield of fluorescence, } \phi_f = \frac{\nu_f}{I_{\text{abs}}} = \frac{k_f[S^*]}{(k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*]} = \frac{k_f}{k_f + k_{\text{ISC}} + k_{\text{IC}}}$$

Fluorescence lifetime can be measured with pulsed laser technique.

Sample is excited with a short light pulse from a laser using wavelength at which

S absorbs strongly

exponential decay of fluorescence intensity after pulse is monitored

$$\tau_0 = \frac{1}{k_f + k_{\text{ISC}} + k_{\text{IC}}} = \frac{k_f}{k_f + k_{\text{ISC}} + k_{\text{IC}}} \times \frac{1}{k_f} = \frac{\phi_f}{k_f}$$

rate of formation of $S^* = I_{\text{abs}}$

rate of decay of $S^* = -k_f[S^*] - k_{\text{ISC}}[S^*] - k_{\text{IC}}[S^*]$

$$\frac{d[S^*]}{dt} = I_{\text{abs}} - (k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*] = 0 \text{ (S.S.)}$$

$$\text{or, } I_{\text{abs}} = (k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*]$$

$$\text{Quantum yield of fluorescence, } \phi_f = \frac{\nu_f}{I_{\text{abs}}} = \frac{k_f[S^*]}{(k_f + k_{\text{ISC}} + k_{\text{IC}})[S^*]} = \frac{k_f}{k_f + k_{\text{ISC}} + k_{\text{IC}}}$$

Fluorescence lifetime can be measured with pulsed laser technique.

Sample is excited with a short light pulse from a laser using wavelength at which

S absorbs strongly

exponential decay of fluorescence intensity after pulse is monitored

$$\tau_0 = \frac{1}{k_f + k_{\text{ISC}} + k_{\text{IC}}} = \frac{k_f}{k_f + k_{\text{ISC}} + k_{\text{IC}}} \times \frac{1}{k_f} = \frac{\phi_f}{k_f}$$

tryptophan in water: $\phi_f = 0.20$; $\tau_0 = 2.6 \text{ ns}$

$$\therefore k_f = \frac{\phi_f}{\tau_0} = \frac{0.20}{2.6 \times 10^{-9} \text{ s}} = 7.7 \times 10^7 \text{ s}^{-1}$$

Quenching: shortening of lifetime of excited state

Quenching: shortening of lifetime of excited state

- either a desired process, e.g., in energy or electron transfer,

Quenching: shortening of lifetime of excited state

- either a desired process, e.g., in energy or electron transfer, or an undesired side reaction decreasing quantum yield of a desired photochemical process.

Quenching: shortening of lifetime of excited state

- either a desired process, e.g., in energy or electron transfer,
or an undesired side reaction decreasing quantum yield of a desired
photochemical process.


Quenching effects are studied by monitoring emission from excited state involved
in photochemical reaction

Quenching: shortening of lifetime of excited state

- either a desired process, e.g., in energy or electron transfer, or an undesired side reaction decreasing quantum yield of a desired photochemical process.

Quenching effects are studied by monitoring emission from excited state involved in photochemical reaction

Addition of quencher, Q, opens additional channel for deactivation:




Quenching: shortening of lifetime of excited state

- either a desired process, e.g., in energy or electron transfer, or an undesired side reaction decreasing quantum yield of a desired photochemical process.

Quenching effects are studied by monitoring emission from excited state involved in photochemical reaction

Addition of quencher, Q, opens additional channel for deactivation:




$$\frac{d[S^*]}{dt} = I_{\text{abs}} - (k_f + k_{ISC} + k_{IC} + k_Q[Q])[S^*] = 0$$

Quenching: shortening of lifetime of excited state

- either a desired process, e.g., in energy or electron transfer, or an undesired side reaction decreasing quantum yield of a desired photochemical process.

Quenching effects are studied by monitoring emission from excited state involved in photochemical reaction

Addition of quencher, Q, opens additional channel for deactivation:



$$\frac{d[S^*]}{dt} = I_{\text{abs}} - (k_f + k_{ISC} + k_{IC} + k_Q[Q])[S^*] = 0$$


$$\phi_f = \frac{k_f}{k_f + k_{ISC} + k_{IC} + k_Q[Q]}$$

Quenching: shortening of lifetime of excited state

- either a desired process, e.g., in energy or electron transfer, or an undesired side reaction decreasing quantum yield of a desired photochemical process.

Quenching effects are studied by monitoring emission from excited state involved in photochemical reaction

Addition of quencher, Q, opens additional channel for deactivation:



$$\frac{d[S^*]}{dt} = I_{\text{abs}} - (k_f + k_{ISC} + k_{IC} + k_Q[Q])[S^*] = 0$$

$$\phi_f = \frac{k_f}{k_f + k_{ISC} + k_{IC} + k_Q[Q]}$$


$$\text{for } [Q]=0, \phi_{f,0} = \frac{k_f}{k_f + k_{ISC} + k_{IC}}$$

Quenching: shortening of lifetime of excited state

- either a desired process, e.g., in energy or electron transfer, or an undesired side reaction decreasing quantum yield of a desired photochemical process.

Quenching effects are studied by monitoring emission from excited state involved in photochemical reaction

Addition of quencher, Q, opens additional channel for deactivation:



$$\frac{d[S^*]}{dt} = I_{\text{abs}} - (k_f + k_{ISC} + k_{IC} + k_Q[Q])[S^*] = 0$$

$$\phi_f = \frac{k_f}{k_f + k_{ISC} + k_{IC} + k_Q[Q]}$$

$$\text{for } [Q]=0, \phi_{f,0} = \frac{k_f}{k_f + k_{ISC} + k_{IC}}$$


$$\frac{\phi_{f,0}}{\phi_f} = \frac{k_f + k_{ISC} + k_{IC} + k_Q[Q]}{k_f + k_{ISC} + k_{IC}} = 1 + \frac{k_Q}{k_f + k_{ISC} + k_{IC}}[Q]$$

Quenching: shortening of lifetime of excited state

- either a desired process, e.g., in energy or electron transfer, or an undesired side reaction decreasing quantum yield of a desired photochemical process.

Quenching effects are studied by monitoring emission from excited state involved in photochemical reaction

Addition of quencher, Q, opens additional channel for deactivation:




$$\frac{d[S^*]}{dt} = I_{\text{abs}} - (k_f + k_{ISC} + k_{IC} + k_Q[Q])[S^*] = 0$$


$$\phi_f = \frac{k_f}{k_f + k_{ISC} + k_{IC} + k_Q[Q]}$$

$$\text{for } [Q]=0, \phi_{f,0} = \frac{k_f}{k_f + k_{ISC} + k_{IC}}$$

$$\begin{aligned} \frac{\phi_{f,0}}{\phi_f} &= \frac{k_f + k_{ISC} + k_{IC} + k_Q[Q]}{k_f + k_{ISC} + k_{IC}} = 1 + \frac{k_Q}{k_f + k_{ISC} + k_{IC}}[Q] \\ &= 1 + \tau_0 k_Q[Q] : \text{Stern-Volmer equation} \end{aligned}$$




The molecule 2,2'-bipyridine forms a complex with the Ru^{2+}



The molecule 2,2'-bipyridine forms a complex with the Ru^{2+}


$\text{Ru}(\text{bipy})_3^{2+}$ has strong metal-to-ligand charge transfer (MLCT) transition at 450 nm.



The molecule 2,2'-bipyridine forms a complex with the Ru^{2+}


$\text{Ru}(\text{bipy})_3^{2+}$ has strong metal-to-ligand charge transfer (MLCT) transition at 450 nm.

Quenching of the $^*\text{Ru}(\text{bipy})_3^{2+}$ excited state by $\text{Fe}(\text{H}_2\text{O})_6^{3+}$ in acidic solution monitored by measuring emission lifetimes at 600 nm.



$[\text{Fe}(\text{H}_2\text{O})_6^{3+}]/(10^{-4} \text{ mol dm}^{-3})$	0	1.6	4.7	7	9.4
$\tau/(\tau_0/10^{-7} \text{ s})$	6	4.05	3.37	2.96	2.17

data:




S

The molecule 2,2'-bipyridine forms a complex with the Ru^{2+}


$\text{Ru}(\text{bipy})_3^{2+}$ has strong metal-to-ligand charge transfer (MLCT) transition at 450 nm.

Quenching of the $^*\text{Ru}(\text{bipy})_3^{2+}$ excited state by $\text{Fe}(\text{H}_2\text{O})_6^{3+}$ in acidic solution monitored by measuring emission lifetimes at 600 nm.



data:

$[\text{Fe}(\text{H}_2\text{O})_6^{3+}]/(10^{-4} \text{ mol dm}^{-3})$	0	1.6	4.7	7	9.4
$\tau/(\tau_0/10^{-7} \text{ s})$	6	4.05	3.37	2.96	2.17




$$s \frac{\tau_0}{\tau} = 1 + \tau_0 k_Q [\text{Q}];$$

The molecule 2,2'-bipyridine forms a complex with the Ru^{2+}


$\text{Ru}(\text{bipy})_3^{2+}$ has strong metal-to-ligand charge transfer (MLCT) transition at 450 nm.

Quenching of the $^*\text{Ru}(\text{bipy})_3^{2+}$ excited state by $\text{Fe}(\text{H}_2\text{O})_6^{3+}$ in acidic solution monitored by measuring emission lifetimes at 600 nm.



data:

$[\text{Fe}(\text{H}_2\text{O})_6^{3+}]/(10^{-4} \text{ mol dm}^{-3})$	0	1.6	4.7	7	9.4
τ/τ_0	6	4.05	3.37	2.96	2.17



$$s \frac{\tau_0}{\tau} = 1 + \tau_0 k_Q [\text{Q}];$$

$$\text{slope} = 2.8 \times 10^9 = k_Q (\text{in dm}^3 \text{mol}^{-1} \text{s}^{-1})$$

Common mechanisms for bimolecular quenching :

Common mechanisms for bimolecular quenching :

Collisional deactivation: $S^* + Q \rightarrow S + Q$

Common mechanisms for bimolecular quenching :

Collisional deactivation: $S^* + Q \rightarrow S + Q$

Resonance energy transfer: $S^* + Q \rightarrow S + Q^*$

Common mechanisms for bimolecular quenching :

Collisional deactivation: $S^* + Q \rightarrow S + Q$

Resonance energy transfer: $S^* + Q \rightarrow S + Q^*$

Electron transfer: $S^* + Q \rightarrow S^+ + Q^-$ or $S^- + Q^+$

Common mechanisms for bimolecular quenching :

Collisional deactivation: $S^* + Q \rightarrow S + Q$

Resonance energy transfer: $S^* + Q \rightarrow S + Q^*$

Electron transfer: $S^* + Q \rightarrow S^+ + Q^-$ or $S^- + Q^+$

Collisional quenching : particularly efficient when Q is a heavy species, such as I^- , which receives energy from S^* and decays primarily by internal conversion to ground state

Common mechanisms for bimolecular quenching :

Collisional deactivation: $S^* + Q \rightarrow S + Q$

Resonance energy transfer: $S^* + Q \rightarrow S + Q^*$

Electron transfer: $S^* + Q \rightarrow S^+ + Q^-$ or $S^- + Q^+$

Collisional quenching : particularly efficient when Q is a heavy species, such as I^- , which receives energy from S^* and decays primarily by internal conversion to ground state

This fact is used to determine the accessibility of amino acid residues of a folded protein to solvent

- fluorescence from tryptophan residue ($\lambda_{abs} = 290 \text{ nm}$, $\lambda_{fluor} = 350 \text{ nm}$) is quenched by I^- ion when residue is on surface of protein and hence accessible to solvent.

Common mechanisms for bimolecular quenching :

Collisional deactivation: $S^* + Q \rightarrow S + Q$

Resonance energy transfer: $S^* + Q \rightarrow S + Q^*$

Electron transfer: $S^* + Q \rightarrow S^+ + Q^-$ or $S^- + Q^+$

Collisional quenching : particularly efficient when Q is a heavy species, such as I^- , which receives energy from S^* and decays primarily by internal conversion to ground state

This fact is used to determine the accessibility of amino acid residues of a folded protein to solvent

- fluorescence from tryptophan residue ($\lambda_{abs} = 290 \text{ nm}$, $\lambda_{fluor} = 350 \text{ nm}$) is quenched by I^- ion when residue is on surface of protein and hence accessible to solvent.

Conversely, residues in hydrophobic interior of protein are not quenched effectively by I^-

Common mechanisms for bimolecular quenching :

Collisional deactivation: $S^* + Q \rightarrow S + Q$

Resonance energy transfer: $S^* + Q \rightarrow S + Q^*$

Electron transfer: $S^* + Q \rightarrow S^+ + Q^-$ or $S^- + Q^+$

Collisional quenching : particularly efficient when Q is a heavy species, such as I^- , which receives energy from S^* and decays primarily by internal conversion to ground state

This fact is used to determine the accessibility of amino acid residues of a folded protein to solvent

- fluorescence from tryptophan residue ($\lambda_{abs} = 290 \text{ nm}$, $\lambda_{fluor} = 350 \text{ nm}$) is quenched by I^- ion when residue is on surface of protein and hence accessible to solvent.

Conversely, residues in hydrophobic interior of protein are not quenched effectively by I^-

Quenching rate constant itself does not give much insight into mechanism of quenching

Common mechanisms for bimolecular quenching :

Collisional deactivation: $S^* + Q \rightarrow S + Q$

Resonance energy transfer: $S^* + Q \rightarrow S + Q^*$

Electron transfer: $S^* + Q \rightarrow S^+ + Q^-$ or $S^- + Q^+$

Collisional quenching : particularly efficient when Q is a heavy species, such as I^- , which receives energy from S^* and decays primarily by internal conversion to ground state

This fact is used to determine the accessibility of amino acid residues of a folded protein to solvent

- fluorescence from tryptophan residue ($\lambda_{abs} = 290$ nm, $\lambda_{fluor} = 350$ nm) is quenched by I^- ion when residue is on surface of protein and hence accessible to solvent.

Conversely, residues in hydrophobic interior of protein are not quenched effectively by I^-

Quenching rate constant itself does not give much insight into mechanism of quenching

Quenching of excited state of $Ru(bipy)_3^{2+}$ is a result of light-induced electron transfer to Fe^{3+} , but quenching data do not prove the mechanism

resonance energy transfer (RET): $S^* + Q \rightarrow S + Q^*$

resonance energy transfer (RET): $S^* + Q \rightarrow S + Q^*$

Oscillating electric field of
incoming radiation induces
oscillating electric dipole moment
in S

resonance energy transfer (RET): $S^* + Q \rightarrow S + Q^*$

Oscillating electric field of incoming radiation induces oscillating electric dipole moment in S

Energy is absorbed by S if frequency of incident radiation $\nu = \frac{\Delta E_s}{h}$, where ΔE_s = energy separation between ground and excited electronic states of S :
'resonance'

resonance energy transfer (RET): $S^* + Q \rightarrow S + Q^*$

Oscillating electric field of incoming radiation induces oscillating electric dipole moment in S

Energy is absorbed by S if frequency of incident radiation $\nu = \frac{\Delta E_s}{h}$, where ΔE_s = energy separation between ground and excited electronic states of S :
'resonance'

Oscillating dipole on S affects electrons bound to nearby Q molecule by inducing oscillating dipole moment in Q.

resonance energy transfer (RET): $S^* + Q \rightarrow S + Q^*$

Oscillating electric field of incoming radiation induces oscillating electric dipole moment in S

Energy is absorbed by S if frequency of incident radiation $\nu = \frac{\Delta E_s}{h}$, where ΔE_s = energy separation between ground and excited electronic states of S :
'resonance'

Oscillating dipole on S affects electrons bound to nearby Q molecule by inducing oscillating dipole moment in Q.

If frequency of oscillation of dipole moment in S is $\nu = \frac{\Delta E_Q}{h}$, then Q absorbs energy from S

resonance energy transfer (RET): $S^* + Q \rightarrow S + Q^*$

Oscillating electric field of incoming radiation induces oscillating electric dipole moment in S


Energy is absorbed by S if frequency of incident radiation $\nu = \frac{\Delta E_s}{h}$, where ΔE_s = energy separation between ground and excited electronic states of S : 'resonance'

Oscillating dipole on S affects electrons bound to nearby Q molecule by inducing oscillating dipole moment in Q.

If frequency of oscillation of dipole moment in S is $\nu = \frac{\Delta E_Q}{h}$, then Q absorbs energy from S

RET Efficiency : $E_T = 1 - \frac{\phi_f}{\phi_{f,0}}$

Fluorescence Resonance Energy Transfer (FRET)




Förster theory (1959)

Förster theory (1959)



energy transfer is efficient when:



- ▶ Energy donor and acceptor are separated by short distance, \mathcal{O} (1 nm)
- ▶ Photons emitted by excited state of donor can be absorbed directly by acceptor.
- ▶ $E_T = \frac{1}{1 + \left(\frac{R}{R_0}\right)^6}$, R in nm and R_0 is a parameter

Donor [†]	Acceptor	R_0 / nm
Naphthalene	Dansyl	2.2
Dansyl	ODR	4.3
Pyrene	Coumarin	3.9
IEDANS	FITC	4.9
Tryptophan	IEDANS	2.2
Tryptophan	Haem (heme)	2.9





FRET:





FRET:


When an amino acid on surface of rhodopsin was labelled covalently with energy donor 1,5-I AEDANS,



FRET:

When an amino acid on surface of rhodopsin was labelled covalently with energy donor 1,5-I AEDANS,

fluorescence quantum yield of label decreased from 0.75 to 0.68 due to quenching by visual pigment 11-cis-retinal.




FRET:

When an amino acid on surface of rhodopsin was labelled covalently with energy donor 1,5-I AEDANS,

fluorescence quantum yield of label decreased from 0.75 to 0.68 due to quenching by visual pigment 11-cis-retinal.

$$E_T = 1 - \frac{0.68}{0.75} = 0.093$$



FRET:

When an amino acid on surface of rhodopsin was labelled covalently with energy donor 1,5-I AEDANS,

fluorescence quantum yield of label decreased from 0.75 to 0.68 due to quenching by visual pigment 11-cis-retinal.

$$E_T = 1 - \frac{0.68}{0.75} = 0.093$$

using known value of $R_0 = 5.4$ nm for 1,5-I AEDANS/11-cis-retinal pair,
we get $R = 7.9$ nm

criteria governing relative efficiencies of collisional quenching, resonance energy transfer, and electron transfer:

criteria governing relative efficiencies of collisional quenching, resonance energy transfer, and electron transfer:

Collisional quenching : particularly efficient when Q, e.g., I^- , receives energy from S^* and decays to ground state primarily by releasing heat

criteria governing relative efficiencies of collisional quenching, resonance energy transfer, and electron transfer:

Collisional quenching : particularly efficient when Q, e.g., I^- , receives energy from S^* and decays to ground state primarily by releasing heat

Electron transfer reactions : Marcus theory (1965) :

criteria governing relative efficiencies of collisional quenching, resonance energy transfer, and electron transfer:

Collisional quenching : particularly efficient when Q, e.g., I^- , receives energy from S^* and decays to ground state primarily by releasing heat

Electron transfer reactions : Marcus theory (1965) :
rates of electron transfer (from ground or excited states) depend on:

criteria governing relative efficiencies of collisional quenching, resonance energy transfer, and electron transfer:

Collisional quenching : particularly efficient when Q, e.g., I^- , receives energy from S^* and decays to ground state primarily by releasing heat

Electron transfer reactions : Marcus theory (1965) :

rates of electron transfer (from ground or excited states) depend on:

- ▶ distance between donor (D) and acceptor (A) ; becoming more efficient as D-A distance decreases

criteria governing relative efficiencies of collisional quenching, resonance energy transfer, and electron transfer:

Collisional quenching : particularly efficient when Q, e.g., I^- , receives energy from S^* and decays to ground state primarily by releasing heat

Electron transfer reactions : Marcus theory (1965) :

rates of electron transfer (from ground or excited states) depend on:

- ▶ distance between donor (D) and acceptor (A) ; becoming more efficient as D-A distance decreases
- ▶ $\Delta_r G$; e-transfer is more efficient as reaction becomes more exergonic, e.g., for efficient photo oxidation of S, reduction potential of S^* must be lower than that of Q

criteria governing relative efficiencies of collisional quenching, resonance energy transfer, and electron transfer:

Collisional quenching : particularly efficient when Q, e.g., I^- , receives energy from S^* and decays to ground state primarily by releasing heat

Electron transfer reactions : Marcus theory (1965) :

rates of electron transfer (from ground or excited states) depend on:

- ▶ distance between donor (D) and acceptor (A) ; becoming more efficient as D-A distance decreases
- ▶ $\Delta_r G$; e-transfer is more efficient as reaction becomes more exergonic, e.g., for efficient photo oxidation of S, reduction potential of S^* must be lower than that of Q
- ▶ **reorganization energy** (energy cost incurred by molecular rearrangements of D, A, and medium) :
electron transfer rate predicted to increase as reorganization energy is matched closely by reaction Gibbs energy

criteria governing relative efficiencies of collisional quenching, resonance energy transfer, and electron transfer:

Collisional quenching : particularly efficient when Q, e.g., I^- , receives energy from S^* and decays to ground state primarily by releasing heat

Electron transfer reactions : Marcus theory (1965) :

rates of electron transfer (from ground or excited states) depend on:

- ▶ distance between donor (D) and acceptor (A) ; becoming more efficient as D-A distance decreases
- ▶ $\Delta_r G$; e-transfer is more efficient as reaction becomes more exergonic, e.g., for efficient photo oxidation of S, reduction potential of S^* must be lower than that of Q
- ▶ **reorganization energy** (energy cost incurred by molecular rearrangements of D, A, and medium) :
electron transfer rate predicted to increase as reorganization energy is matched closely by reaction Gibbs energy

Electron transfer can be studied by time-resolved spectroscopy.

Oxidized and reduced products often have electronic absorption spectra distinct from those of neutral parent compounds. Rapid appearance of such known features in absorption spectrum after excitation by laser pulse may be taken as indication of quenching by electron transfer

Overall quantum yield of a photochemical reaction

Overall quantum yield of a photochemical reaction
For complex reactions involving secondary processes,

Overall quantum yield of a photochemical reaction

For complex reactions involving secondary processes,

e.g., chain reactions initiated by photolysis, many reactant molecules may be consumed as a result of absorption of single photon.

Overall quantum yield of a photochemical reaction

For complex reactions involving secondary processes,

e.g., chain reactions initiated by photolysis, many reactant molecules may be consumed as a result of absorption of single photon.

Overall quantum yield, Φ , # reactant molecules consumed per photon absorbed may exceed 1.




Overall quantum yield of a photochemical reaction

For complex reactions involving secondary processes,

e.g., chain reactions initiated by photolysis, many reactant molecules may be consumed as a result of absorption of single photon.

Overall quantum yield, Φ , # reactant molecules consumed per photon absorbed may exceed 1.

photolysis of HI:






Overall quantum yield of a photochemical reaction

For complex reactions involving secondary processes,

e.g., chain reactions initiated by photolysis, many reactant molecules may be consumed as a result of absorption of single photon.

Overall quantum yield, Φ , # reactant molecules consumed per photon absorbed may exceed 1.

photolysis of HI:



\therefore absorption of 1 photon leads to destruction of 2 HI molecules

\therefore Overall quantum yield = 2




Overall quantum yield of a photochemical reaction

For complex reactions involving secondary processes,

e.g., chain reactions initiated by photolysis, many reactant molecules may be consumed as a result of absorption of single photon.

Overall quantum yield, Φ , # reactant molecules consumed per photon absorbed may exceed 1.

photolysis of HI:



\therefore absorption of 1 photon leads to destruction of 2 HI molecules

\therefore Overall quantum yield = 2




In chain reactions, Φ may be very large : values $\sim 10^4$ are common

Overall quantum yield of a photochemical reaction

For complex reactions involving secondary processes, e.g., chain reactions initiated by photolysis, many reactant molecules may be consumed as a result of absorption of single photon.

Overall quantum yield, Φ , # reactant molecules consumed per photon absorbed may exceed 1.

photolysis of HI:



\therefore absorption of 1 photon leads to destruction of 2 HI molecules

\therefore Overall quantum yield = 2

In chain reactions, Φ may be very large : values $\sim 10^4$ are common
the chain acts as chemical amplifier of initial absorption step

4-heptanone irradiated

4-heptanone irradiated
for 100 s

4-heptanone irradiated
for 100 s
with 313 nm radiation

4-heptanone irradiated
for 100 s
with 313 nm radiation
with a power output of 50 W

4-heptanone irradiated
for 100 s
with 313 nm radiation
with a power output of 50 W
under conditions of total absorption,

4-heptanone irradiated
for 100 s
with 313 nm radiation
with a power output of 50 W
under conditions of total absorption,
2.8 mmol C_2H_4 formed.

4-heptanone irradiated
for 100 s
with 313 nm radiation
with a power output of 50 W
under conditions of total absorption,
2.8 mmol C₂H₄ formed.

amount (in moles) of photons absorbed, $n = \frac{P\Delta t}{\left(\frac{hc}{\lambda}\right)N_A}$

4-heptanone irradiated
for 100 s
with 313 nm radiation
with a power output of 50 W
under conditions of total absorption,
2.8 mmol C₂H₄ formed.

amount (in moles) of photons absorbed, $n = \frac{P\Delta t}{\left(\frac{hc}{\lambda}\right)N_A}$


$$\Phi = \frac{n_{C_2H_4}}{n} = \frac{n_{C_2H_4} N_A hc}{\lambda P \Delta t} = \frac{2.8 \times 10^{-3} \times 6.023 \times 10^{23} \times 6.636 \times 10^{-34} \times 2.998 \times 10^8}{3.13 \times 10^{-7} \times 50 \times 100} = 0.21$$

Rate laws of complex photochemical reactions:

Rate laws of complex photochemical reactions:
consider photochemical activation of chain reaction


Rate laws of complex photochemical reactions:
consider photochemical activation of chain reaction
 $\text{H}_2(\text{g}) + \text{Br}_2(\text{g}) \rightarrow 2\text{HBr}(\text{g})$

Rate laws of complex photochemical reactions:
consider photochemical activation of chain reaction



Earlier for thermal reaction,

Rate laws of complex photochemical reactions:
consider photochemical activation of chain reaction




Earlier for thermal reaction,



Initiation: $\text{Br}_2 + \text{M} \longrightarrow \text{Br}\cdot + \text{Br}\cdot + \text{M}$; $v = k_i[\text{Br}_2][\text{M}]$

Rate laws of complex photochemical reactions:

consider photochemical activation of chain reaction




Earlier for thermal reaction,





Rate laws of complex photochemical reactions:

consider photochemical activation of chain reaction



Earlier for thermal reaction,




Earlier, we had:



$$\frac{d[\text{HBr}]}{dt} = 2k_{p'}[H][\text{Br}_2] = \frac{2k_p \sqrt{\frac{k_i}{k_t}} [H_2][\text{Br}_2]^{\frac{3}{2}}}{[\text{Br}_2] + \frac{k_r}{k_{p'}} [\text{HBr}]} = \frac{k[H_2][\text{Br}_2]^{\frac{3}{2}}}{[\text{Br}_2] + k'[\text{HBr}]}$$

Rate laws of complex photochemical reactions:

consider photochemical activation of chain reaction



Earlier for thermal reaction,



Earlier, we had:

$$\frac{d[\text{HBr}]}{dt} = 2k_{p'}[H][\text{Br}_2] = \frac{2k_p \sqrt{\frac{k_i}{k_t}} [H_2][\text{Br}_2]^{\frac{3}{2}}}{[\text{Br}_2] + \frac{k_r}{k_{p'}} [\text{HBr}]} = \frac{k[H_2][\text{Br}_2]^{\frac{3}{2}}}{[\text{Br}_2] + k'[\text{HBr}]}$$

now: I_{abs} should take the place of $k_i[\text{Br}_2][\text{M}]$

$$\frac{d[\text{HBr}]}{dt} = 2k_{p'}[H][\text{Br}_2] = \frac{2k_p \sqrt{\frac{1}{k_t[M]}} [H_2][\text{Br}_2] \sqrt{I_{\text{abs}}}}{[\text{Br}_2] + \frac{k_r}{k_{p'}} [\text{HBr}]} = \frac{k[H_2][\text{Br}_2]}{[\text{Br}_2] + k'[\text{HBr}]} \sqrt{I_{\text{abs}}}$$

photosensitization:

photosensitization:

Reactions of a molecule that does not absorb directly can be made to occur if another absorbing molecule is present

photosensitization:

Reactions of a molecule that does not absorb directly can be made to occur if another absorbing molecule is present

because latter may be able to transfer its energy to former during a collision

photosensitization:

Reactions of a molecule that does not absorb directly can be made to occur if another absorbing molecule is present

because latter may be able to transfer its energy to former during a collision

e.g., generation of excited state O_2 in photodynamic therapy:

photosensitization:

Reactions of a molecule that does not absorb directly can be made to occur if another absorbing molecule is present

because latter may be able to transfer its energy to former during a collision

e.g., generation of excited state O_2 in photodynamic therapy:

laser radiation delivered to diseased tissue through a fibre optic cable

photosensitization:

Reactions of a molecule that does not absorb directly can be made to occur if another absorbing molecule is present

because latter may be able to transfer its energy to former during a collision

e.g., generation of excited state O_2 in photodynamic therapy:

laser radiation delivered to diseased tissue through a fibre optic cable

absorbed by a drug which, in its first excited triplet state 3P , photosensitizes the formation of excited 1O_2

photosensitization:

Reactions of a molecule that does not absorb directly can be made to occur if another absorbing molecule is present

because latter may be able to transfer its energy to former during a collision

e.g., generation of excited state O_2 in photodynamic therapy:

laser radiation delivered to diseased tissue through a fibre optic cable

absorbed by a drug which, in its first excited triplet state 3P , photosensitizes the formation of excited 1O_2

Absorption: $P + h\nu \rightarrow P^*$

photosensitization:

Reactions of a molecule that does not absorb directly can be made to occur if another absorbing molecule is present

because latter may be able to transfer its energy to former during a collision

e.g., generation of excited state O_2 in photodynamic therapy:

laser radiation delivered to diseased tissue through a fibre optic cable

absorbed by a drug which, in its first excited triplet state 3P , photosensitizes the formation of excited 1O_2

Absorption: $P + h\nu \rightarrow P^*$

ISC: $P^* \rightarrow ^3P$

photosensitization:

Reactions of a molecule that does not absorb directly can be made to occur if another absorbing molecule is present

because latter may be able to transfer its energy to former during a collision

e.g., generation of excited state O_2 in photodynamic therapy:

laser radiation delivered to diseased tissue through a fibre optic cable

absorbed by a drug which, in its first excited triplet state 3P , photosensitizes the formation of excited 1O_2

Absorption: $P + h\nu \rightarrow P^*$

ISC: $P^* \rightarrow ^3P$

Photosensitization: $^3P + ^3O_2 \rightarrow P + ^1O_2$

photosensitization:

Reactions of a molecule that does not absorb directly can be made to occur if another absorbing molecule is present

because latter may be able to transfer its energy to former during a collision

e.g., generation of excited state O_2 in photodynamic therapy:

laser radiation delivered to diseased tissue through a fibre optic cable

absorbed by a drug which, in its first excited triplet state 3P , photosensitizes the formation of excited 1O_2

Absorption: $P + h\nu \rightarrow P^*$

ISC: $P^* \rightarrow ^3P$

Photosensitization: $^3P + ^3O_2 \rightarrow P + ^1O_2$

1O_2 molecules are very reactive and destroy cellular components and it is thought that cell membranes are primary cellular targets

photosensitization:

Reactions of a molecule that does not absorb directly can be made to occur if another absorbing molecule is present

because latter may be able to transfer its energy to former during a collision

e.g., generation of excited state O_2 in photodynamic therapy:

laser radiation delivered to diseased tissue through a fibre optic cable

absorbed by a drug which, in its first excited triplet state 3P , photosensitizes the formation of excited 1O_2

Absorption: $P + h\nu \rightarrow P^*$

ISC: $P^* \rightarrow ^3P$

Photosensitization: $^3P + ^3O_2 \rightarrow P + ^1O_2$

1O_2 molecules are very reactive and destroy cellular components and it is thought that cell membranes are primary cellular targets

Photosensitization: $^3P + ^3O_2 \rightarrow P + ^1O_2$

photosensitization:

Reactions of a molecule that does not absorb directly can be made to occur if another absorbing molecule is present

because latter may be able to transfer its energy to former during a collision

e.g., generation of excited state O_2 in photodynamic therapy:

laser radiation delivered to diseased tissue through a fibre optic cable

absorbed by a drug which, in its first excited triplet state 3P , photosensitizes the formation of excited 1O_2

Absorption: $P + h\nu \rightarrow P^*$

ISC: $P^* \rightarrow ^3P$

Photosensitization: $^3P + ^3O_2 \rightarrow P + ^1O_2$

1O_2 molecules are very reactive and destroy cellular components and it is thought that cell membranes are primary cellular targets

Photosensitization: $^3P + ^3O_2 \rightarrow P + ^1O_2$

Oxidation reactions: $^1O_2 + \text{reactants} \rightarrow \text{products}$

photosensitization:

Reactions of a molecule that does not absorb directly can be made to occur if another absorbing molecule is present

because latter may be able to transfer its energy to former during a collision

e.g., generation of excited state O_2 in photodynamic therapy:

laser radiation delivered to diseased tissue through a fibre optic cable

absorbed by a drug which, in its first excited triplet state 3P , photosensitizes the formation of excited 1O_2

Absorption: $P + h\nu \rightarrow P^*$

ISC: $P^* \rightarrow ^3P$

Photosensitization: $^3P + ^3O_2 \rightarrow P + ^1O_2$

1O_2 molecules are very reactive and destroy cellular components and it is thought that cell membranes are primary cellular targets

Photosensitization: $^3P + ^3O_2 \rightarrow P + ^1O_2$

Oxidation reactions: $^1O_2 + \text{reactants} \rightarrow \text{products}$

example: $P \equiv \text{Haematoporphyrin}$; characteristics ?

Another example :

Another example :
generation of atomic H,

Another example :
generation of atomic H,
irradiation of H₂ gas containing trace of Hg vapour using radiation of 254 nm
from a Hg discharge lamp

Another example :
generation of atomic H,
irradiation of H₂ gas containing trace of Hg vapour using radiation of 254 nm
from a Hg discharge lamp
The Hg atoms are excited (to Hg*) by resonant absorption of
radiation



Another example :

generation of atomic H,

irradiation of H₂ gas containing trace of Hg vapour using radiation of 254 nm from a Hg discharge lamp

The Hg atoms are excited (to Hg*) by resonant absorption of radiation

and then collide with H₂ molecules.





Another example :

generation of atomic H,




irradiation of H₂ gas containing trace of Hg vapour using radiation of 254 nm from a Hg discharge lamp

The Hg atoms are excited (to Hg*) by resonant absorption of radiation

and then collide with H₂ molecules.



The latter reaction is the initiation step for other mercury photosensitized reactions, such as synthesis of formaldehyde from carbon monoxide and hydrogen





Another example :

generation of atomic H,



irradiation of H₂ gas containing trace of Hg vapour using radiation of 254 nm from a Hg discharge lamp

The Hg atoms are excited (to Hg*) by resonant absorption of radiation

and then collide with H₂ molecules.



The latter reaction is the initiation step for other mercury photosensitized reactions, such as synthesis of formaldehyde from carbon monoxide and hydrogen
 $\text{H} \cdot + \text{CO} \rightarrow \text{HCO} \cdot$



last step is termination by disproportionation rather than by combination