

catalysis:

catalysis:

decomposition of H_2O_2 in solution,

$E_{\text{act}} = 76 \text{ kJ mol}^{-1}$ at 298K.

catalysis:

decomposition of H_2O_2 in solution,

$E_{\text{act}} = 76 \text{ kJ mol}^{-1}$ at 298K.

When a little I^- ion is added, $E_{\text{act}} = 57 \text{ kJ mol}^{-1}$

k increases by a factor of 2000

catalysis:

decomposition of H_2O_2 in solution,


$E_{\text{act}} = 76 \text{ kJ mol}^{-1}$ at 298K.

When a little I^- ion is added, $E_{\text{act}} = 57 \text{ kJ mol}^{-1}$

k increases by a factor of 2000

With Enzyme catalase $E_{\text{act}} = 8 \text{ kJ mol}^{-1}$

k increases by a factor of 10^{15}



catalysis:

decomposition of H_2O_2 in solution,

$E_{\text{act}} = 76 \text{ kJ mol}^{-1}$ at 298K.



When a little I^- ion is added, $E_{\text{act}} = 57 \text{ kJ mol}^{-1}$

k increases by a factor of 2000

With Enzyme catalase $E_{\text{act}} = 8 \text{ kJ mol}^{-1}$

k increases by a factor of 10^{15}

Homogeneous : catalyst in same phase as reaction mixture



catalysis:

decomposition of H_2O_2 in solution,

$E_{\text{act}} = 76 \text{ kJ mol}^{-1}$ at 298K.


When a little I^- ion is added, $E_{\text{act}} = 57 \text{ kJ mol}^{-1}$

k increases by a factor of 2000

With Enzyme catalase $E_{\text{act}} = 8 \text{ kJ mol}^{-1}$


k increases by a factor of 10^{15}

Homogeneous : catalyst in same phase as reaction mixture



Heterogeneous : catalyst in different phase as reaction mixture,

e.g.,



catalysis:

decomposition of H_2O_2 in solution,

$E_{\text{act}} = 76 \text{ kJ mol}^{-1}$ at 298K.


When a little I^- ion is added, $E_{\text{act}} = 57 \text{ kJ mol}^{-1}$

k increases by a factor of 2000



With Enzyme catalase $E_{\text{act}} = 8 \text{ kJ mol}^{-1}$

k increases by a factor of 10^{15}

Homogeneous : catalyst in same phase as reaction mixture



Heterogeneous : catalyst in different phase as reaction mixture,



catalysis:

decomposition of H_2O_2 in solution,

$E_{\text{act}} = 76 \text{ kJ mol}^{-1}$ at 298K.


When a little I^- ion is added, $E_{\text{act}} = 57 \text{ kJ mol}^{-1}$

k increases by a factor of 2000


With Enzyme catalase $E_{\text{act}} = 8 \text{ kJ mol}^{-1}$

k increases by a factor of 10^{15}


Homogeneous : catalyst in same phase as reaction mixture



Heterogeneous : catalyst in different phase as reaction mixture,



metal provides surface upon which reactants bind



catalysis:

decomposition of H_2O_2 in solution,


$E_{\text{act}} = 76 \text{ kJ mol}^{-1}$ at 298K.

When a little I^- ion is added, $E_{\text{act}} = 57 \text{ kJ mol}^{-1}$


k increases by a factor of 2000

With Enzyme catalase $E_{\text{act}} = 8 \text{ kJ mol}^{-1}$


k increases by a factor of 10^{15}



Homogeneous : catalyst in same phase as reaction mixture



Heterogeneous : catalyst in different phase as reaction mixture,



metal provides surface upon which reactants bind

binding facilitates encounters between reactants and increases rxn rate

Example of **homogeneous catalysis**:

Example of **homogeneous catalysis**:

Br^- catalysed decomposition of H_2O_2 : $2\text{H}_2\text{O}_2(\text{aq}) \longrightarrow 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$

Example of **homogeneous catalysis**:

Br^- catalysed decomposition of H_2O_2 : $2\text{H}_2\text{O}_2(\text{aq}) \longrightarrow 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$

pre-equilibrium: $\text{H}_3\text{O}^+ + \text{H}_2\text{O}_2 \rightleftharpoons \text{H}_3\text{O}_2^+ + \text{H}_2\text{O}$; $K = \frac{[\text{H}_3\text{O}_2^+]}{[\text{H}_2\text{O}_2][\text{H}_3\text{O}^+]}$

Example of **homogeneous catalysis**:

Br^- catalysed decomposition of H_2O_2 : $2\text{H}_2\text{O}_2(\text{aq}) \longrightarrow 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$

pre-equilibrium: $\text{H}_3\text{O}^+ + \text{H}_2\text{O}_2 \rightleftharpoons \text{H}_3\text{O}_2^+ + \text{H}_2\text{O}$; $K = \frac{[\text{H}_3\text{O}_2^+]}{[\text{H}_2\text{O}_2][\text{H}_3\text{O}^+]}$

$\text{H}_3\text{O}_2^+ + \text{Br}^- \longrightarrow \text{HOBr} + \text{H}_2\text{O}$, $v = k[\text{H}_3\text{O}_2^+][\text{Br}^-]$

Example of **homogeneous catalysis**:

Br^- catalysed decomposition of H_2O_2 : $2\text{H}_2\text{O}_2(\text{aq}) \longrightarrow 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$

pre-equilibrium: $\text{H}_3\text{O}^+ + \text{H}_2\text{O}_2 \rightleftharpoons \text{H}_3\text{O}_2^+ + \text{H}_2\text{O}$; $K = \frac{[\text{H}_3\text{O}_2^+]}{[\text{H}_2\text{O}_2][\text{H}_3\text{O}^+]}$

$\text{H}_3\text{O}_2^+ + \text{Br}^- \longrightarrow \text{HOBr} + \text{H}_2\text{O}$, $v = k[\text{H}_3\text{O}_2^+][\text{Br}^-]$

$\text{HOBr} + \text{H}_2\text{O}_2 \longrightarrow \text{H}_3\text{O}^+ + \text{O}_2 + \text{Br}^-$; fast

Example of **homogeneous catalysis**:

Br^- -catalysed decomposition of H_2O_2 : $2\text{H}_2\text{O}_2(\text{aq}) \longrightarrow 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$

pre-equilibrium: $\text{H}_3\text{O}^+ + \text{H}_2\text{O}_2 \rightleftharpoons \text{H}_3\text{O}_2^+ + \text{H}_2\text{O}$; $K = \frac{[\text{H}_3\text{O}_2^+]}{[\text{H}_2\text{O}_2][\text{H}_3\text{O}^+]}$

$\text{H}_3\text{O}_2^+ + \text{Br}^- \longrightarrow \text{HOBr} + \text{H}_2\text{O}$, $v = k[\text{H}_3\text{O}_2^+][\text{Br}^-]$

$\text{HOBr} + \text{H}_2\text{O}_2 \longrightarrow \text{H}_3\text{O}^+ + \text{O}_2 + \text{Br}^-$; fast

rate: $\frac{d[\text{O}_2]}{dt} = k_{\text{eff}} [\text{H}_2\text{O}_2] [\text{H}_3\text{O}^+][\text{Br}^-]$; $k_{\text{eff}} = k.K$

Example of **homogeneous catalysis**:

Br^- -catalysed decomposition of H_2O_2 : $2\text{H}_2\text{O}_2(\text{aq}) \longrightarrow 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$

pre-equilibrium: $\text{H}_3\text{O}^+ + \text{H}_2\text{O}_2 \rightleftharpoons \text{H}_3\text{O}_2^+ + \text{H}_2\text{O}$; $K = \frac{[\text{H}_3\text{O}_2^+]}{[\text{H}_2\text{O}_2][\text{H}_3\text{O}^+]}$

$\text{H}_3\text{O}_2^+ + \text{Br}^- \longrightarrow \text{HOBr} + \text{H}_2\text{O}$, $v = k[\text{H}_3\text{O}_2^+][\text{Br}^-]$

$\text{HOBr} + \text{H}_2\text{O}_2 \longrightarrow \text{H}_3\text{O}^+ + \text{O}_2 + \text{Br}^-$; fast

rate: $\frac{d[\text{O}_2]}{dt} = k_{\text{eff}} [\text{H}_2\text{O}_2] [\text{H}_3\text{O}^+][\text{Br}^-]$; $k_{\text{eff}} = k.K$

Experimentally variation of rate with $[\text{Br}^-]$ and pH agrees with above

Example of **homogeneous catalysis**:

Br^- -catalysed decomposition of H_2O_2 : $2\text{H}_2\text{O}_2(\text{aq}) \longrightarrow 2\text{H}_2\text{O}(\text{l}) + \text{O}_2(\text{g})$

pre-equilibrium: $\text{H}_3\text{O}^+ + \text{H}_2\text{O}_2 \rightleftharpoons \text{H}_3\text{O}_2^+ + \text{H}_2\text{O}$; $K = \frac{[\text{H}_3\text{O}_2^+]}{[\text{H}_2\text{O}_2][\text{H}_3\text{O}^+]}$

$\text{H}_3\text{O}_2^+ + \text{Br}^- \longrightarrow \text{HOBr} + \text{H}_2\text{O}$, $v = k[\text{H}_3\text{O}_2^+][\text{Br}^-]$


$\text{HOBr} + \text{H}_2\text{O}_2 \longrightarrow \text{H}_3\text{O}^+ + \text{O}_2 + \text{Br}^-$; fast

rate: $\frac{d[\text{O}_2]}{dt} = k_{\text{eff}} [\text{H}_2\text{O}_2] [\text{H}_3\text{O}^+][\text{Br}^-]$; $k_{\text{eff}} = k.K$

Experimentally variation of rate with $[\text{Br}^-]$ and pH agrees with above


Other examples: Acid/base catalysis

Enzyme catalysis (Homogeneous)




Enzyme catalysis (Homogeneous)

lock-n-key: active site and substrate have complementary 3D structures



Enzyme catalysis (Homogeneous)


lock-n-key: active site and substrate have complementary 3D structures



induced fit model: binding induces a conformational change in active site

Enzyme catalysis (Homogeneous)

lock-and-key: active site and substrate have complementary 3D structures




induced fit model: binding induces a conformational change in active site

active site returns to its original state after products are released

Enzyme catalysis (Homogeneous)

lock-n-key: active site and substrate have complementary 3D structures



induced fit model: binding induces a conformational change in active site


active site returns to its original state after products are released

Many enzymes consist primarily of proteins

some featuring organic or inorganic co-factors in active sites

Enzyme catalysis (Homogeneous)

lock-n-key: active site and substrate have complementary 3D structures



induced fit model: binding induces a conformational change in active site

active site returns to its original state after products are released


Many enzymes consist primarily of proteins

some featuring organic or inorganic co-factors in active sites

Certain RNA molecules can also be biological catalysts, forming ribozymes

Enzyme catalysis (Homogeneous)

lock-n-key: active site and substrate have complementary 3D structures



induced fit model: binding induces a conformational change in active site

active site returns to its original state after products are released

Many enzymes consist primarily of proteins

some featuring organic or inorganic co-factors in active sites

Certain RNA molecules can also be biological catalysts, forming ribozymes

example of ribozyme \Rightarrow ribosome, large assembly of proteins and catalytically active RNA molecules responsible for synthesis of proteins in cell

1. For a given initial concentration of substrate, $[S]_0$,
initial rate of product formation \propto total concentration of enzyme, $[E]_0$

1. For a given initial concentration of substrate, $[S]_0$,
initial rate of product formation \propto total concentration of enzyme, $[E]_0$
2. For a given $[E]_0$ and low values of $[S]_0$,

rate of product formation $\propto [S]_0$.

1. For a given initial concentration of substrate, $[S]_0$,
initial rate of product formation \propto total concentration of enzyme, $[E]_0$
2. For a given $[E]_0$ and low values of $[S]_0$,

$$\text{rate of product formation} \propto [S]_0.$$

3. For a given $[E]_0$ and high values of $[S]_0$,
rate of product formation : independent of $[S]_0$
reaches maximum velocity, v_{\max}

Michaelis-Menten mechanism: $E + S \xrightleftharpoons[k'_a]{k_a} ES \xrightarrow{k_b} P + E;$

Michaelis-Menten mechanism: $E + S \xrightleftharpoons[k'_a]{k_a} ES \xrightarrow{k_b} P + E$;

$$\text{SSA} \implies \frac{d[ES]}{dt} = k_a[E][S] - k'_a[ES] - k_b[ES] = 0$$

Michaelis-Menten mechanism: $E + S \xrightleftharpoons[k'_a]{k_a} ES \xrightarrow{k_b} P + E$;

$$\text{SSA} \implies \frac{d[ES]}{dt} = k_a[E][S] - k'_a[ES] - k_b[ES] = 0$$

$$\text{or, } [ES] = \frac{k_a}{k'_a + k_b} [E][S] = K_M^{-1} [E][S]$$

Michaelis-Menten mechanism: $E + S \xrightleftharpoons[k'_a]{k_a} ES \xrightarrow{k_b} P + E$;

$$SSA \implies \frac{d[ES]}{dt} = k_a[E][S] - k'_a[ES] - k_b[ES] = 0$$

$$\text{or, } [ES] = \frac{k_a}{k'_a + k_b} [E][S] = K_M^{-1} [E][S]$$

$$\text{Using } [E] = [E]_0 - [ES]; \quad K_M [ES] = ([E]_0 - [ES]) [S]$$


Michaelis-Menten mechanism: $E + S \xrightleftharpoons[k'_a]{k_a} ES \xrightarrow{k_b} P + E$;

$$SSA \Rightarrow \frac{d[ES]}{dt} = k_a[E][S] - k'_a[ES] - k_b[ES] = 0$$

$$\text{or, } [ES] = \frac{k_a}{k'_a + k_b} [E][S] = K_M^{-1} [E][S]$$

$$\text{Using } [E] = [E]_0 - [ES]; \quad K_M [ES] = ([E]_0 - [ES]) [S]$$

$$\text{or, } (K_M + [S]) [ES] = [E]_0 [S] \Rightarrow [ES] = \frac{[E]_0}{1 + \frac{K_M}{[S]}} \approx \frac{[E]_0}{1 + \frac{K_M}{[S]_0}}$$




Michaelis-Menten mechanism: $E + S \xrightleftharpoons[k'_a]{k_a} ES \xrightarrow{k_b} P + E$;

$$SSA \Rightarrow \frac{d[ES]}{dt} = k_a[E][S] - k'_a[ES] - k_b[ES] = 0$$

$$\text{or, } [ES] = \frac{k_a}{k'_a + k_b} [E][S] = K_M^{-1} [E][S]$$

$$\text{Using } [E] = [E]_0 - [ES]; \quad K_M [ES] = ([E]_0 - [ES])[S]$$

$$\text{or, } (K_M + [S])[ES] = [E]_0[S] \Rightarrow [ES] = \frac{[E]_0}{1 + \frac{K_M}{[S]}} \approx \frac{[E]_0}{1 + \frac{K_M}{[S]_0}}$$



$$v = \frac{v_{\max}}{1 + \frac{K_M}{[S]_0}}; \quad v_{\max} = k_b[E]_0$$


Michaelis-Menten mechanism: $E + S \xrightleftharpoons[k'_a]{k_a} ES \xrightarrow{k_b} P + E$;

$$SSA \Rightarrow \frac{d[ES]}{dt} = k_a[E][S] - k'_a[ES] - k_b[ES] = 0$$

$$\text{or, } [ES] = \frac{k_a}{k'_a + k_b} [E][S] = K_M^{-1} [E][S]$$

$$\text{Using } [E] = [E]_0 - [ES]; \quad K_M [ES] = ([E]_0 - [ES])[S]$$

$$\text{or, } (K_M + [S])[ES] = [E]_0[S] \Rightarrow [ES] = \frac{[E]_0}{1 + \frac{K_M}{[S]}} \approx \frac{[E]_0}{1 + \frac{K_M}{[S]_0}}$$



$$v = \frac{v_{\max}}{1 + \frac{K_M}{[S]_0}}; \quad v_{\max} = k_b[E]_0$$

$$[S]_0 \ll K_M \Rightarrow v = \frac{k_b}{k_M} [S]_0 [E]_0$$

$$[S]_0 \gg K_M \Rightarrow v = v_{\max} = k_b[E]_0$$

$$\frac{1}{v} = \frac{1}{v_{\max}} + \left(\frac{K_M}{v_{\max}} \right) \frac{1}{[S]_0}$$


Michaelis-Menten mechanism: $E + S \xrightleftharpoons[k'_a]{k_a} ES \xrightarrow{k_b} P + E$;

$$SSA \Rightarrow \frac{d[ES]}{dt} = k_a[E][S] - k'_a[ES] - k_b[ES] = 0$$

$$\text{or, } [ES] = \frac{k_a}{k'_a + k_b} [E][S] = K_M^{-1} [E][S]$$

$$\text{Using } [E] = [E]_0 - [ES]; \quad K_M [ES] = ([E]_0 - [ES]) [S]$$

$$\text{or, } (K_M + [S]) [ES] = [E]_0 [S] \Rightarrow [ES] = \frac{[E]_0}{1 + \frac{K_M}{[S]}} \approx \frac{[E]_0}{1 + \frac{K_M}{[S]_0}}$$



$$v = \frac{v_{max}}{1 + \frac{K_M}{[S]_0}}; \quad v_{max} = k_b[E]_0$$

$$[S]_0 \ll K_M \Rightarrow v = \frac{k_b}{K_M} [S]_0 [E]_0$$

$$[S]_0 \gg K_M \Rightarrow v = v_{max} = k_b[E]_0$$

$$\frac{1}{v} = \frac{1}{v_{max}} + \left(\frac{K_M}{v_{max}} \right) \frac{1}{[S]_0}$$


Lineweaver-Burk plot

Lineweaver-Burk plot

Lineweaver-Burk plot

catalysis of hydration of CO_2 in red blood cells by carbonic anhydrase


$[\text{CO}_2]/(\text{mmol dm}^{-3})$	1.25	2.5	5	20
rate/ $(\text{mmol dm}^{-3} \text{ s}^{-1})$	2.78×10^{-2}	5.00×10^{-2}	8.33×10^{-2}	1.67×10^{-1}
$1/([\text{CO}_2]/(\text{mmol dm}^{-3}))$	0.800	0.400	0.200	0.0500
$1/(v/(\text{mmol dm}^{-3} \text{ s}^{-1}))$	36.0	20.0	12.0	6.0



Lineweaver-Burk plot

catalysis of hydration of CO_2 in red blood cells by carbonic anhydrase

$[\text{CO}_2]/(\text{mmol dm}^{-3})$	1.25	2.5	5	20
rate/ $(\text{mmol dm}^{-3} \text{ s}^{-1})$	2.78×10^{-2}	5.00×10^{-2}	8.33×10^{-2}	1.67×10^{-1}
$1/([\text{CO}_2]/(\text{mmol dm}^{-3}))$	0.800	0.400	0.200	0.0500
$1/(v/(\text{mmol dm}^{-3} \text{ s}^{-1}))$	36.0	20.0	12.0	6.0




$$v_{\max} = \frac{1}{4} = 0.25 \text{ mmol} \cdot \text{dm}^{-3} \text{ s}^{-1}$$

Lineweaver-Burk plot

catalysis of hydration of CO_2 in red blood cells by carbonic anhydrase

$[\text{CO}_2]/(\text{mmol dm}^{-3})$	1.25	2.5	5	20
rate/ $(\text{mmol dm}^{-3} \text{ s}^{-1})$	2.78×10^{-2}	5.00×10^{-2}	8.33×10^{-2}	1.67×10^{-1}
$1/([\text{CO}_2]/(\text{mmol dm}^{-3}))$	0.800	0.400	0.200	0.0500
$1/(v/(\text{mmol dm}^{-3} \text{ s}^{-1}))$	36.0	20.0	12.0	6.0




$$v_{\max} = \frac{1}{4} = 0.25 \text{ mmol} \cdot \text{dm}^{-3} \text{ s}^{-1}$$

$$K_M = \frac{40}{4} = 10 \text{ mmol} \cdot \text{dm}^{-3}$$

Lineweaver-Burk plot

catalysis of hydration of CO_2 in red blood cells by carbonic anhydrase

$[\text{CO}_2]/(\text{mmol dm}^{-3})$	1.25	2.5	5	20
rate/ $(\text{mmol dm}^{-3} \text{ s}^{-1})$	2.78×10^{-2}	5.00×10^{-2}	8.33×10^{-2}	1.67×10^{-1}
$1/([\text{CO}_2]/(\text{mmol dm}^{-3}))$	0.800	0.400	0.200	0.0500
$1/(v/(\text{mmol dm}^{-3} \text{ s}^{-1}))$	36.0	20.0	12.0	6.0



$$v_{\max} = \frac{1}{4} = 0.25 \text{ mmol} \cdot \text{dm}^{-3} \text{ s}^{-1}$$


$$K_M = \frac{40}{4} = 10 \text{ mmol} \cdot \text{dm}^{-3}$$

$$k_{\text{cat}} = \frac{v_{\max}}{[E]_0} = \frac{2.5 \times 10^{-4}}{2.3 \times 10^{-9}} = 1.1 \times 10^5 \text{ s}^{-1}$$

Lineweaver-Burk plot

catalysis of hydration of CO_2 in red blood cells by carbonic anhydrase

$[\text{CO}_2]/(\text{mmol dm}^{-3})$	1.25	2.5	5	20
rate/ $(\text{mmol dm}^{-3} \text{ s}^{-1})$	2.78×10^{-2}	5.00×10^{-2}	8.33×10^{-2}	1.67×10^{-1}
$1/([\text{CO}_2]/(\text{mmol dm}^{-3}))$	0.800	0.400	0.200	0.0500
$1/(v/(\text{mmol dm}^{-3} \text{ s}^{-1}))$	36.0	20.0	12.0	6.0



$$v_{\max} = \frac{1}{4} = 0.25 \text{ mmol} \cdot \text{dm}^{-3} \text{ s}^{-1}$$

$$K_M = \frac{40}{4} = 10 \text{ mmol} \cdot \text{dm}^{-3}$$

$$k_{\text{cat}} = \frac{v_{\max}}{[E]_0} = \frac{2.5 \times 10^{-4}}{2.3 \times 10^{-9}} = 1.1 \times 10^5 \text{ s}^{-1}$$

$$\epsilon = \frac{k_{\text{cat}}}{K_M} = \frac{1.1 \times 10^5}{10^{-2}} = 1.1 \times 10^7 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$$

turnover frequency or catalytic constant, $k_{cat} = \frac{\text{\#catalytic cycles performed by active site}}{\text{duration of interval}}$
has units of 1st order rate constant

turnover frequency or catalytic constant, $k_{cat} = \frac{\text{\#catalytic cycles performed by active site}}{\text{duration of interval}}$

has units of 1st order rate constant

numerically equivalent to k_b in Michaelis-Menten mechanism

$$k_{cat} = k_b = \frac{v_{max}}{[E]_0}$$

turnover frequency or catalytic constant, $k_{cat} = \frac{\text{\#catalytic cycles performed by active site}}{\text{duration of interval}}$

has units of 1st order rate constant

numerically equivalent to k_b in Michaelis-Menten mechanism

$$k_{cat} = k_b = \frac{v_{max}}{[E]_0}$$

$$\text{catalytic efficiency, } \varepsilon = \frac{k_{cat}}{K_M} = \frac{k_a k_b}{k'_a + k_b}$$

Higher $\varepsilon \implies$ more efficient

turnover frequency or catalytic constant, $k_{cat} = \frac{\text{\#catalytic cycles performed by active site}}{\text{duration of interval}}$

has units of 1st order rate constant

numerically equivalent to k_b in Michaelis-Menten mechanism

$$k_{cat} = k_b = \frac{v_{max}}{[E]_0}$$

$$\text{catalytic efficiency, } \varepsilon = \frac{k_{cat}}{K_M} = \frac{k_a k_b}{k'_a + k_b}$$

Higher $\varepsilon \implies$ more efficient

efficiency is max ($= k_a$) when $k_b \gg k'_a$.

turnover frequency or catalytic constant, $k_{cat} = \frac{\text{\#catalytic cycles performed by active site}}{\text{duration of interval}}$

has units of 1st order rate constant

numerically equivalent to k_b in Michaelis-Menten mechanism

$$k_{cat} = k_b = \frac{v_{max}}{[E]_0}$$

$$\text{catalytic efficiency, } \varepsilon = \frac{k_{cat}}{K_M} = \frac{k_a k_b}{k'_a + k_b}$$

Higher $\varepsilon \implies$ more efficient


efficiency is max ($= k_a$) when $k_b \gg k'_a$.

max efficiency related to max rate of diffusion of E and S in solution

\implies rate constants $\approx 10^8 - 10^9 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$ (catalytic perfection)

Inhibitor : decreases rate of product formation from substrate by binding to enzyme, to ES complex, or to enzyme and ES complex simultaneously

Inhibitor : decreases rate of product formation from substrate
by binding to enzyme, to ES complex, or to enzyme and ES complex simultaneously



Inhibitor : decreases rate of product formation from substrate
by binding to enzyme, to ES complex, or to enzyme and ES complex simultaneously





$$[E]_0 = [E] + [EI] + [ES] + [ESI]$$

Inhibitor : decreases rate of product formation from substrate

by binding to enzyme, to ES complex, or to enzyme and ES complex simultaneously

$$\text{define: } \alpha = 1 + \frac{[I]}{K_I}$$


$$\alpha' = 1 + \frac{[I]}{K'_I}$$



$$[E]_0 = [E] + [EI] + [ES] + [ESI]$$

Inhibitor : decreases rate of product formation from substrate

by binding to enzyme, to ES complex, or to enzyme and ES complex simultaneously



$$[E]_0 = [E] + [EI] + [ES] + [ESI]$$

$$\text{define: } \alpha = 1 + \frac{[I]}{K_I}$$


$$\alpha' = 1 + \frac{[I]}{K'_I}$$

$$[E]_0 = [E]\alpha + [ES]\alpha'$$

$$\text{using } K_M = \frac{[E][S]}{[ES]}$$

Inhibitor : decreases rate of product formation from substrate

by binding to enzyme, to ES complex, or to enzyme and ES complex simultaneously



$$[E]_0 = [E] + [EI] + [ES] + [ESI]$$

$$\text{define: } \alpha = 1 + \frac{[I]}{K_I}$$

$$\alpha' = 1 + \frac{[I]}{K'_I}$$

$$[E]_0 = [E]\alpha + [ES]\alpha'$$

$$\text{using } K_M = \frac{[E][S]}{[ES]}$$




$$[E]_0 = \frac{K_M [ES]}{[S]_0} \alpha + [ES] \alpha' = [ES] \left(\frac{\alpha K_M}{[S]_0} + \alpha' \right)$$

$$v = k_b [ES] = \frac{k_b [E]_0}{\frac{\alpha K_M}{[S]_0} + \alpha'} = \frac{v_{\max}}{\alpha' + \frac{\alpha K_M}{[S]_0}}$$

$$\text{or, } \frac{1}{v} = \frac{\alpha'}{v_{\max}} + \alpha \left(\frac{K_M}{v_{\max}} \right) \frac{1}{[S]_0}$$

Lineweaver-Burk plots characteristic of three modes of enzyme inhibition:




$$\frac{1}{v} = \frac{\alpha'}{v_{max}} + \alpha \left(\frac{K_M}{v_{max}} \right) \frac{1}{[S]}$$



Lineweaver-Burk plots characteristic of three modes of enzyme inhibition:

competitive inhibition: inhibitor binds only to active site of enzyme and inhibits attachment of substrate; $\alpha > 1$ and $\alpha' = 1$

$$\frac{1}{v} = \frac{\alpha'}{v_{max}} + \alpha \left(\frac{K_M}{v_{max}} \right) \frac{1}{[S]_0}$$






Lineweaver-Burk plots characteristic of three modes of enzyme inhibition:

competitive inhibition: inhibitor binds only to active site of enzyme and inhibits attachment of substrate; $\alpha > 1$ and $\alpha' = 1$

uncompetitive inhibition : inhibitor binds to enzyme site removed from active site, but only if substrate is already present. ESI reduces the concentration of ES (active complex). In this case $\alpha = 1$ (\because EI does not form) and $\alpha' > 1$.

$$\frac{1}{v} = \frac{\alpha'}{v_{max}} + \alpha \left(\frac{K_M}{v_{max}} \right) \frac{1}{[S]_0}$$



Lineweaver-Burk plots characteristic of three modes of enzyme inhibition:




competitive inhibition: inhibitor binds only to active site of enzyme and inhibits attachment of substrate; $\alpha > 1$ and $\alpha' = 1$

uncompetitive inhibition : inhibitor binds to enzyme site removed from active site, but only if substrate is already present. ESI reduces the concentration of ES (active complex). In this case $\alpha = 1$ (\because EI does not form) and $\alpha' > 1$.

non-competitive (mixed) inhibition : inhibitor binds to site other than active site, and its presence reduces ability of substrate to bind to active site.

Inhibition occurs at both E and ES sites : $\alpha > 1$ and $\alpha' > 1$.

$$\frac{1}{v} = \frac{\alpha'}{v_{max}} + \alpha \left(\frac{K_M}{v_{max}} \right) \frac{1}{[S]_0}$$



Lineweaver-Burk plots characteristic of three modes of enzyme inhibition:

competitive inhibition: inhibitor binds only to active site of enzyme and inhibits attachment of substrate; $\alpha > 1$ and $\alpha' = 1$


uncompetitive inhibition : inhibitor binds to enzyme site removed from active site, but only if substrate is already present. ESI reduces the concentration of ES (active complex). In this case $\alpha = 1$ (\because EI does not form) and $\alpha' > 1$.

non-competitive (mixed) inhibition : inhibitor binds to site other than active site, and its presence reduces ability of substrate to bind to active site.



Inhibition occurs at both E and ES sites : $\alpha > 1$ and $\alpha' > 1$.

Slope and y-intercept of Lineweaver-Burk plot increase upon addition of inhibitor. Fig. c : special case : $K_I = K'_I$; and $\alpha = \alpha'$, which results in intersection of lines on x-axis

$$\frac{1}{v} = \frac{\alpha'}{v_{max}} + \alpha \left(\frac{K_M}{v_{max}} \right) \frac{1}{[S]_0}$$





Autocatalysis:



$$v = k[A][P]$$



labelled by b

Autocatalysis:



labelled by b

Autocatalysis:




$$\frac{dx}{dt} = k([A]_0 - x)([P]_0 + x)$$

integrating:


$$\frac{1}{[A]_0 + [P]_0} \ln \frac{([P]_0 + x)[A]_0}{[P]_0([A]_0 - x)} = kt$$

labelled by b

Autocatalysis:



labelled by b



$$\frac{dx}{dt} = k([A]_0 - x)([P]_0 + x)$$

integrating:


$$\frac{1}{[A]_0 + [P]_0} \ln \frac{([P]_0 + x)[A]_0}{[P]_0([A]_0 - x)} = kt$$

$$\text{or, } \frac{x}{[P]_0} = \frac{e^{at} - 1}{1 + be^{at}}, \text{ where } a = ([A]_0 + [P]_0)k$$

$$\text{and } b = \frac{[P]_0}{[A]_0}$$


oscillating reactions: Feedback mechanism in which a product either increases or decreases the reaction rate

- can be sustained indefinitely only if the reaction is carried at far from equilibrium conditions (continuous supply and dissipation of species)



oscillating reactions: Feedback mechanism in which a product either increases or decreases the reaction rate


- can be sustained indefinitely only if the reaction is carried at far from equilibrium conditions (continuous supply and dissipation of species)





under anerobic conditions, oscillations in [NADH] can occur in glycolytic cycle in yeast cells because of inhibition of the enzyme phosphofructokinase by high levels of ATP and its activation by high levels of ADP

Lotka-Volterra (Prey-predator) mechanism:




Lotka-Volterra (Prey-predator) mechanism:



Lotka-Volterra (Prey-predator) mechanism:




Lotka-Volterra (Prey-predator) mechanism:





Belousov-Zhabotinski reaction (KBrO_3 , malonic acid, cerium (IV) salt in acidic solution):



Belousov-Zhabotinski reaction (KBrO₃, malonic acid, cerium (IV) salt in acidic solution):



Belousov-Zhabotinski reaction (KBrO_3 , malonic acid, cerium (IV) salt in acidic solution):




Belousov-Zhabotinski reaction (KBrO_3 , malonic acid, cerium (IV) salt in acidic solution):




product HBrO_2 is a reactant in first step and provides a feedback mechanism that enhances rate of formation of HBrO_2

<https://en.wikipedia.org/wiki/Brusselator>



<https://en.wikipedia.org/wiki/Brusselator>



<https://en.wikipedia.org/wiki/Oregonator>

