



DEPARTMENT OF INFORMATION TECHNOLOGY

FACULTY PROFILE REPORT

Employee Code: 7001 | Generated: 30-04-2026

1. PERSONAL INFORMATION

Full Name	KAMBHAMPATI VIJAY KUMAR
Employee Code	7001
Father's Name	KAMBHAMPATI KANAKA DURGA RAO
Mother's Name	KAMBHAMPATI SHOBHANA KUMARI
Date of Birth	24 May 1984
Gender	Male
State	Telangana
Caste	BC-D
Sub-Caste	MUNNURUKAPU
Nationality	Indian
Permanent Address	H.No:- 6-35/12, MARUTHI NAGAR, WARANGAL CROSS ROAD, YEDULAPURAM, KHAMMAM - 507003

2. PROFESSIONAL INFORMATION

Department	IT
Designation	Assistant Professor
Joining Date	10 Dec 2010
Experience at Anurag	15 Years 3 Months
Experience (Other)	1
Total Experience	15 Years 4 Months 20 Days
Email	hod.inf@anurag.ac.in
Mobile	9553122276
Phone	9030900962
Current Status	ACTIVE

3. PROFESSIONAL IDs

JNTUH ID	15150406-114529
AICTE ID	1-468376653
PAN Number	ARXPK1671R
Aadhar Number	846452283847
APAAR ID	72-265-370-947
ORCID ID	0000-0002-9791-8445

4. EDUCATIONAL QUALIFICATIONS - SSC / 10TH

Year	2000
Percentage	56.2
School	SES VV CENTRAL PUBLIC SCHOOL, KHAMMAM

5. EDUCATIONAL QUALIFICATIONS - INTERMEDIATE

Year	2002
Percentage	57.1
College	EXCELLENT JUNIOR COLLEGE, KHAMMAM

6. EDUCATIONAL QUALIFICATIONS - UG

Degree	B.TECH
Year	2006
Percentage	55.6
College	SREE KAVITHA ENGINEERING COLLEGE, KHAMMAM
Specialization	INFORMATION TECHNOLOGY

7. EDUCATIONAL QUALIFICATIONS - PG

Degree	M.TECH
Year	2010
Percentage	72.4
College	SREE KAVITHA ENGINEERING COLLEGE, KHAMMAM
Specialization	SOFTWARE ENGINEERING

8. EDUCATIONAL QUALIFICATIONS - PhD

Status	Pursuing
---------------	----------

Year	Not Specified
University	KLU, VIJAYAWADA
Specialization	COMPUTER SCIENCE AND ENGINEERING

9. SUBJECTS DEALT

DBMS PPS DATA MINING

10. SCM (SERVICE/CONSULTANCY/MOU)

K. VIJAY KUMAR SCM ON 28-12-2019

11. RESEARCH & PUBLICATIONS

Type	Title	Academic Year	Year	Status	Details
Journal Article	Analytical Framework to Quantify and Scale Air Pollution Intensity with Machine Learning	2024-2025	2024	Published	Authors: K. VIJAY KUMAR & Dr.V.M. SUBRAMANYAM Journal: AFRICAN JOURNAL OF BIOLOGICAL SCIENCES ISSN: 2663-2187 DOI: https://doi.org/10.33472/AFJBS.6.Si2.2024.1421-1427 URL: http://www.afjbs.com

13. FDP / WORKSHOPS ATTENDED

Type	Title	Academic Year	Duration	Organized By	Place
FDP	Advances in GenAI and Agentic AI	2024-2025	16 Jun 2025 to 21 Jun 2025 (6 days)	Department of Computer Science and Engineering in collaboration with Computer Society of India, CSI-SB, CVR College of Engineering	Hyderabad

14. B.TECH PROJECTS HANDLED

H.T.No.	Student Name	Batch	Project Title	Approved	Marks
23C11A1213	DHARANI KAMBALA	A1	ONLINE EXAM	Yes	45

15. RESULTS / ACADEMIC PERFORMANCE

No results data specified

16. UPLOADED DOCUMENTS

Aadhar Card: ✓ Uploaded

PAN Card: ✓ Uploaded

APAAR Document: ✓ Uploaded

SCM Document: ✓ Uploaded

JNTUH Bio-Data: ✓ Uploaded

SSC Certificate: ✓ Uploaded

Intermediate Certificate: ✓ Uploaded

UG Certificate: ✓ Uploaded

PG Certificate: ✓ Uploaded

PhD Certificate: ✗ Not Uploaded

17. ADDITIONAL DOCUMENTS

Research Publications Proof: ✓ 7 pages uploaded AY: 2024-2025

FDP Proof: ✓ 1 pages uploaded AY: 2024-2025

Experience Certificates: ✗ Not Uploaded

Other Documents: ✗ Not Uploaded

Research Publications Proof (2024-2025):

FDP Proof (2024-2025):

18. ABOUT YOURSELF

I WAS PROMOTED AS HOD OF IT ON 28-12-2023




భారత ప్రభుత్వం
Government of India

భారత విశిష్ట గుర్తింపు ప్రాధికార సంస్థ
Unique Identification Authority of India

రెజిస్ట్రేషన్/Enrolment No.: 0705/50703/05488

To
కంభంపాటి విజయ కుమార్
Kambhampati Vijay Kumar
S/O Kambhampati Kanaka Durga Rao,
6-35/12,
Maruthi Nagar,
Warangal X Road,
Khammam rural,
VTC: Edulapuram (Rural),
PO: Edulapuram,
Sub District: Khammam (rural),
District: Khammam,
State: Telangana,
PIN Code: 507163,
Mobile: 9000900962



మీ ఆధార్ సంఖ్య / Your Aadhaar No. :

8464 5228 3847

VID : 9152 5598 6642 3770

నా ఆధార్, నా గుర్తింపు



భారత ప్రభుత్వం
Government of India



కంభంపాటి విజయ కుమార్
Kambhampati Vijay Kumar
పుట్టిన తేదీ/DOB: 24/05/1984
ప్రభుత్వం/MALE

ఆధార్ అనేది గుర్తింపు దస్తావజు, పౌరత్వం, లేదా పుట్టిన తేదీ కి కాదు. ఇది ధృవీకరణ ద్వారానే అధికారికంగా అంగీకరించబడినది లేదా QR కోడ్ / ఆన్లైన్ వెబ్ యూజ్ థ్రూ.

Aadhaar is proof of identity, not of citizenship or date of birth, it should be used with verification (online authentication, or scanning of QR code / offline NDL).

8464 5228 3847

నా ఆధార్, నా గుర్తింపు



సమాచారము / INFORMATION

- ఆధార్ అనేది గుర్తింపు దస్తావజు, పౌరత్వం, లేదా పుట్టిన తేదీ కి కాదు. పుట్టిన తేదీ అనేది ఆధార్ సంఖ్య చొప్పున సమన్వయించిన సేవలందుకునే పర్సన్ ను పుట్టిన తేదీ ద్వారా యొక్క దస్తావజు ఆధారం ద్వారా ఇచ్చే సమాచారంపై ఆధారపడి ఉంటుంది.
- ఈ ఆధార్ లేకపోతే UIDAI వెబ్సైటుపై ప్రమాణీకరించిన ఏకైక ద్వారా ఆన్లైన్ ప్రమాణీకరించిన ద్వారా లేదా యాన్ థ్రూలో అందుబాటులో ఉన్న mAadhaar లేదా ఆధార్ QR స్కానర్ యాప్ ద్వారా అధికారికంగా లేదా www.uidai.gov.inలో అందుబాటులో ఉన్న మరలిన QR కోడ్ కింద యాన్ అధికారికంగా QR కోడ్ స్కాన్ చేసి ద్వారా ధృవీకరించాలి.
- ఆధార్ ప్రశ్నోత్తరాలకు మరలు సులభమైనది.
- ఆధార్ పరిమితి లేనిది తేదీ ముడి పై 10 సంవత్సరాల తర్వాత గుర్తింపు మరలు రేఖాంశాలు సంతకం చేసిన పత్రాలతో ఆధార్ ను సవరించాలి.
- వివిధ ప్రభుత్వ మరియు ప్రభుత్వాల ప్రయోజనాలు/సేవలను పొందడంలో ఆధార్ మీకు సహాయపడుతుంది.
- మీ మొత్తం ముఖ మరియు ఈ-మొఖల్ థాట్ ఆధార్ లో అందజేసే సేవలో ఉంది.
- ఆధార్ సేవలను పొందేందుకు mAadhaar యాప్ ను డౌన్లోడ్ చేసుకోండి.
- ఆధార్/అధికారికతలను అధికారికంగా ఉపయోగించే ద్వారా విధించబడిన లాక్/అన్ లాక్ ఆధార్/అధికారికతల సేవలను ఉపయోగించండి.
- ఆధార్ ను కోరి సంఖ్యల ద్వారా విధించే ఉపయోగ పొందటం.
- Aadhaar is proof of identity, not of citizenship or date of birth (DOB). DOB is based on information supported by proof of DOB document specified in regulations, submitted by Aadhaar number holder.
- This Aadhaar letter should be verified through either online authentication by UIDAI-appointed authentication agency or QR code scanning using mAadhaar or Aadhaar QR Scanner app available in app stores or using secure QR code reader app available on www.uidai.gov.in.
- Aadhaar is unique and secure.
- Documents to support identity and address should be updated in Aadhaar after every 10 years from date of enrolment for Aadhaar.
- Aadhaar helps you avail of various Government and Non-Government benefits/services.
- Keep your mobile number and email id updated in Aadhaar.
- Download mAadhaar app to avail of Aadhaar services.
- Use the feature of Lock/Unlock Aadhaar biometrics to ensure security when not using Aadhaar biometrics.
- Entities seeking Aadhaar are obligated to seek consent.




భారత విశిష్ట గుర్తింపు ప్రాధికార సంస్థ
Unique Identification Authority of India



కంభంపాటి విజయ కుమార్
S/O Kambhampati Kanaka Durga Rao, 6-35/12, Maruthi Nagar,
Warangal X Road, Khammam rural, Edulapuram (Rural),
PO: Edulapuram, Sub District: Khammam (rural),
District: Khammam, State: Telangana - 507163

Address:
S/O Kambhampati Kanaka Durga Rao, 6-35/12, Maruthi Nagar,
Warangal X Road, Khammam rural, Edulapuram (Rural), PO:
Edulapuram, DIST: Khammam,
Telangana - 507163



8464 5228 3847

VID : 9152 5598 6642 3770

आयकर विभाग
INCOME TAX DEPARTMENT



भारत सरकार
GOVT. OF INDIA

KAMBHAMPATI VIJAY KUMAR

KANAKADURGARAO KAMBHAMPATI

24/05/1984

Permanent Account Number

ARXPK1671R

K. Vijaykumar

Signature



50028052



ACADEMIC BANK OF CREDITS
Ministry of Education, Government of India

Logout



Hello
Kambhampati Vijay Kumar

0	-	0
Credit Points	NCRF Weightage	Credits

ACADEMIC BANK CREDITS

APAAR ID

722-265-370-947

Kambhampati Vijay Kumar



We use basic information from your Digilocker account to identify you and allow access to ABC Platform

Credit Points Accumulation

If any credit data is accumulated, it will appear here.

College Name: ANURAG ENGINEERING COLLEGE

CC: C1

{Pg.No 1/3}

Proceeding No: ANR/K/CSE/SCM/2018-20/01

(Affiliated to JNTUH)

Minutes of the Selection Committee Meeting held on: 28-11-2018 at: KODAD

Post : Assistant Professor

Department : CSE

Scale of Pay :

S. No	Name of the Faculty	Faculty Portal Reg. Number	PAN Number	UG Branch	PG Branch	Ph.D.	Years of Experience
1.	M. Sridevi	51150404-160046	ASAPG5942M	B-Level M.D.C.A	CSE	-	02
2.	K. Vijay Kumar	15150406-114529	ARXPK1671R	IT	SE	-	08
3.	N. Mownika	1063-160125-121649	AZEPN3608C	IT	CSE	-	03
4.	DR. CH. N. Santhosh Kumar	8127-150409-121259	AFLPND271L	M.Sc	CSE	CSE	14
5.	M. Navya	8443-170125-101117	CXSPM3615A	CSE	CSE	-	02

S. No Role

Name

Signature

1. Chairperson : K. Jayapal Reddy
2. Principal : DR. M. V. Siva Prasad
3. University Nominee 1 : DR. M. Dhana Lakshmi
4. University Nominee 2 : MR. P. Srinivas Rao
5. Subject Expert 1 : DR. N. Sambasiva Rao
6. Subject Expert 2 :

7. SC/ST/OBC/Women/ Differently Abled if any }

Note: 1. It is the sole responsibility of the concerned College / Institute to verify whether the eligibility criteria of the candidates applied is in accordance with AICTE/PCI norms.
2. In respect of faculty members possessing Ph.D. Degrees, the SCMs of Faculty members are subject to the recommendation of the subject expert committee constituted for the purpose.

College Name: ANURAG ENGINEERING COLLEGE

CC: C1

[Pg-No-2/3]

Proceeding No: ANRK/CSE/SCM/2019-20/02

(Affiliated to JNTUH)

Minutes of the Selection Committee Meeting held on: 28-12-2019 at: KODAP

Post : Assistant Professor

Department: CSE

Scale of Pay :

S. No	Name of the Faculty	Faculty Portal Reg. Number	PAN Number	UG Branch	PG Branch	Ph.D.	Years of Experience
1.	M. Kivam	9475-191108-124208	DPLPM9475N	CSE	CSE	-	-
2.	M. Rambabu	11150-402-1555341	BKXPM2642E	CSE	SE	-	02
3.	Y.V.R. Naga Pawan	5629-150408-152930	AA7PY77265	MCA	CSE	-	08
4.	GVSCHSLV Prasad	5026-150408-102332	ASAP60657K	MCA	CSE	-	02
5.	J. Vijaya Sree	9815-0407-160625	ASOPT2457J	IT	CSE	-	05

S. No Role Name Signature

- Chairperson : K. Jayapal Reddy K.J
- Principal : Dr. M. V. Siva prasad M.V.S.P
- University Nominee 1 : Dr. M. Phana Lakshmi P.L
- University Nominee 2 : Mr. P. Srinivasa Rao P.S
- Subject Expert 1 : Dr. N. Sambasiva Rao N.S
- Subject Expert 2 :

7. SC/ST/OBC/Women/ Differently Abled if any :

Note: 1. It is the sole responsibility of the concerned College / Institute to verify whether the eligibility criteria of the candidates applied is in accordance with AICTE/PCI norms.
2. In respect of faculty members possessing Ph.D. Degrees, the SCMs of Faculty members are subject to the recommendation of the subject expert committee constituted for the purpose.

College Name: ANURAG ENGINEERING COLLEGE

CC: C1

[Pg. No 3/3]

Proceeding No: ANR/K/CSE/SCM/2019-20/03 (Affiliated to JNTUH)

Minutes of the Selection Committee Meeting held on: 28-12-2019 at: KODAD

Post : Assistant Professor

Department : CSE

Scale of Pay :

S. No	Name of the Faculty	Faculty Portal Reg. Number	PAN Number	UG Branch	PG Branch	Ph.D.	Years of Experience
1.	P. Niranjan Kumar	4138-160315112106	AYWPP81468	IT	CSE	-	08
2.	M. Mahesh Babu	1255-162213-102033	AWTPN9205E	IT	CS	-	03
3.	CH. Anjaiah	4326-160219-142003	ASRPC8733H	IT	CSE	-	03
4.							
5.							

S. No Role Name Signature

1. Chairperson : K. Jayapal Reddy
2. Principal : Dr. M. V. Siva prasad
3. University Nominee 1 : Dr. M. Dhana Lakshmi
4. University Nominee 2 : Mr. P. Srinivasa Rao
5. Subject Expert 1 : Dr. N. Sambasiva Rao
6. Subject Expert 2 :
7. SC/ST/OBC/Women/ Differently Abled if any :

Note: 1. It is the sole responsibility of the concerned College / Institute to verify whether the eligibility criteria of the candidates applied is in accordance with AICTE/PCI norms.

2. In respect of faculty members possessing Ph.D. Degrees, the SCMs of Faculty members are subject to the recommendation of the subject expert committee constituted for the purpose.

Jawaharlal Nehru Technological University Hyderabad
Directorate of Affiliations & Academic Audit
Kukatpally, Hyderabad – 500 085, Telangana, India

Faculty Registration Information :

Registration ID : 15150406-114529
 Full Name :
 First Name : KAMBHAMPATI
 Middle Name : VIJAY
 Last Name : KUMAR



Faculty Basic Information :

Father's/Husband's Name :	K KANAKA DURGA RAO	Mother Name :	K SOBHANA KUMARI
Gender :	Male	Date of Birth :	24/05/1984
Mobile No :	9030900962	Email :	vijaykumar_1247@yahoo.co.in
PAN Number :	ARXPK1671R	Aadhaar Number :	84645228XXXX
PAN Document :	Yes	Aadhaar Document :	Yes
Category :	BC-D		

Faculty College Information : (Automatically taken from College Portal Where Presently Working)

Name of the College presently working in	ANURAG ENGINEERING COLLEGE (C1)		
Department :	IT	Designation :	Assistant Professor
Experience in the present Institution (years)	14 Years 0 Months 12 Days	Total Experience :	0 Years 0 Months 0 Days
AICTE Faculty Id :	1-468376653	Gross Salary Last Drawn:	VI pay
Date of Appointment :	10/02/2011		

Faculty Education Information :

Education Name	Course Studied	Branch /Specialization	Year of Passing	% of Marks /CGPA	Division	Board /University	Place	Certificate
10th / Equivalent	SSC	SSC	2000	56.20	2	CBSE	KHAMMA M	Yes
UG	B.TECH	IT	2006	55.60	2	JNTU	KHAMMA M	Yes
PG	M.TECH	SE	2010	72.40	1	JNTU	KHAMMA M	Yes

क्रमांक 009712
5/6-2000

केन्द्रीय माध्यमिक शिक्षा बोर्ड
CENTRAL BOARD OF SECONDARY EDUCATION
अंक विवरणिका MARKS STATEMENT
शैक्षणिक वर्ष २०००
ALL INDIA SECONDARY SCHOOL EXAMINATION, 2000

नाम Name KANHANPATI VIJAYKUMAR

अनुसूचक क्रमांक Roll No. 4107989

माता का नाम Mother's Name K. SUSHANA KUMARI

पिता का नाम Father's Name K. KANAKADURGA RAO

जन्म तिथि Date of Birth 24TH MAY NINETEEN HUNDRED EIGHTY FOUR

विद्यालय School 6148 SES XV CENTRAL PUB SCH RACHUNADHAPALEM KANMAN

कोड CODE	विषय SUBJECT	प्राप्तकृत अंक MARKS OBTAINED				स्थितिगत अंक POSITIONAL GRADE
		हि TH	ई PR	योग TOTAL	योग शब्दों में TOTAL IN WORDS	
001	ENGLISH COURSE-A	058	XXX	058	FIFTY EIGHT	C1
007	TELVUGU	078	XXX	078	SEVENTY EIGHT	B2
041	MATHEMATICS	037	XIV	037	THIRTY SEVEN	D1
086	SCIENCE WITH PRAC.	055	023	055	FIFTY EIGHT	B2
087	SOCIAL SCIENCE	050	XXX	050	FIFTY	C1

अवधानी का अर्थ : Attention

अ. वि. अंक : अंक विवरणिका में दर्शाया गया है

परिणत Result PASS

दिनांक Date

दिनांक Date 03-06-2000

PRINCIPAL

U.S.V. Yashwanth Public School
KHAMMAN

मुख्य निदेशक

Controller of Examinations


SL. No. **C556128**

IS-4181/C556128



Board of Intermediate Education A. P.

Vidya Bhawan, Narayana, Hyderabad-500 0



INTERMEDIATE PASS CERTIFICATE CUM MEMORANDUM OF

This is to certify that **KANAKADURGA RAO**of **K. KANAKADURGA RAO**Registered No. **1234567**

has appeared and passed at the Intermediate

Public Examination held in **JUNE 2002**in **COMPARTMENTALLY**with **ENGLISH**

as the Medium of Instruction.

The subjects in which he/she was examined and the marks awarded are as follows:

Subject	I Year		II Year	
	Maximum Marks	Marks Secured	Maximum Marks	Marks Secured
Part - 1 :				
English	100	078*	100	084*
Part - 2 :				
SAMANYI	100	067*	100	088*
Part - 3 :				
Optional Subjects				
MATHEMATICS-A	075	027*	075	029*
MATHEMATICS-B	075	033*	075	020
PHYSICS	060	024*	060	028*
PHYSICS PRACTICAL			030	023*
CHEMISTRY	060	022*	060	027*
CHEMISTRY PRACTICAL			030	025*
Total Marks in Figures	571			
In Words	FIVE "SEVEN ONE"			
Date	18-07-2002			

Principal

Examination Officer, Narayana College, Sec.

K. K. KANAKADURGA RAO

Coordinator of Examinations

CC 140409



JAWAHARLAL NEHRU TECHNOLOGICAL UNIVERSITY



HYDERABAD, ANDHRA PRADESH, INDIA

College Code: CS (S K E C, KARPALLY)

Mr. Hambhampati Vijaykumar
S/o H Hanahadurga Rao

having fulfilled the academic requirements and passed the examination
held during *April - 2006* in *** Second Class ***
has this day been admitted by the Executive Council to the Degree of

Bachelor of Technology
(Information Technology)

Given under the Seal of the University

H.T.No : 02C81A1247

0631804

Date : 30 March, 2007

[Signature]

Kelkothra Saddy

[Signature]

JAWAHARLAL NEHRU TECHNOLOGICAL UNIVERSITY HYDERABAD

HYDERABAD - 500 085, ANDHRA PRADESH, INDIA

RF455603

3661445287



430001046816

College Code: C8 (S K E C, KARPALLY)

Mr. Hambhampati Vijaykumar

S/o Mr. K. Kanakadurga Rao

having fulfilled the academic requirements and passed the examination
held during *November - 2010* in *First Class With Distinction*
has this day been admitted by the Executive Council to the Degree of

Master of Technology
Software Engineering

Given under the Seal of the University

H.T.No:08C81D2509

Date: 07-05-2012

DIRECTOR OF EVALUATION

REGISTRAR

VICE-CHANCELLOR





Analytical Framework to Quantify and Scale Air Pollution Intensity with Machine Learning

^{*1}K. Vijay Kumar

Assistant Professor, IT Department, Anurag Engineering College, Ananthagiri(V&M), Kodad, Suryapet, Telangana.

^{*1}Email: hod.inf@anurag.ac.in

²V. M. Subramanyam

²Assistant Professor, Anurag engineering college, (An autonomous institution) Ananthagiri (V&M), kodad Suryapet district

²Email: Subrahmanyamvadlamaan@gmail.com

***Corresponding Author**

Article History

Volume 6, Issue Si2, 2024

Received: 27 Mar 2024

Accepted : 28 Apr 2024

doi: [10.33472/AFJBS.6.Si2.2024.1421-1427](https://doi.org/10.33472/AFJBS.6.Si2.2024.1421-1427)

Abstract: This article presents a comprehensive analytical framework designed to quantify and scale air pollution intensity using machine learning techniques. With the increasing impact of air pollution on public health and the environment, there is a pressing need for advanced methodologies that can provide accurate, scalable, and real-time assessments of air quality. Our framework addresses this need by integrating diverse data sources, including air quality measurements, meteorological data, emission inventories, and geospatial information, to construct a detailed representation of pollution patterns. The core of our framework involves a series of methodologically rigorous steps: comprehensive data compilation, meticulous data preprocessing, standardized data normalization, innovative feature engineering, and selective feature reduction. These steps prepare the dataset for effective machine learning model training and validation. We employ Z-score normalization to standardize numerical inputs, enhancing model convergence and performance. Recursive Feature Elimination (RFE) is utilized to identify and retain the most predictive features, streamlining the modeling process. A Random Forest Classifier, selected for its robustness and ability to handle nonlinear relationships, is trained to categorize air quality into predefined classes based on Air Quality Index (AQI) levels. This model not only quantifies but also scales air pollution intensity across different geographic and temporal scales, providing a tool for policymakers, environmental agencies, and the public to understand and mitigate the effects of air pollution. Through this framework, we demonstrate the application of machine learning in environmental science, highlighting its potential to transform air quality monitoring and management. Our approach is validated with a case study, showing promising results in predicting and scaling air pollution levels with high accuracy. This work lays the groundwork for future research and development in the field, suggesting directions for integrating more dynamic data sources and employing advanced machine learning algorithms to further enhance the predictive capability and scalability of air pollution models.

Keywords: Recursive Feature Elimination (RFE); Air Quality Index (AQI); Random Forest; Scale Air Pollution; Machine Learning; Support Vector Machine (SVM).

1. INTRODUCTION

In the modern era, air pollution is a formidable challenge with far-reaching consequences for public health, ecological balance, and the global climate. The rapid pace of urbanization and industrial expansion has increased the complexity and severity of air pollution, necessitating the development of novel and effective strategies for monitoring, quantifying, and mitigating its

negative effects [1]. Traditional air quality assessment methodologies, which primarily rely on ground-based monitoring stations, provide precise measurements but are inherently limited in their spatial coverage. This constraint limits their ability to capture the complex spatial and temporal dynamics of air pollution, which is critical for a comprehensive understanding and management of its consequences. The emergence and

K. Vijay Kumar / Afr.J.Bio.Sc. 6(Si2) (2024)

integration of machine learning (ML) technologies into environmental sciences marks a new beginning in addressing these challenges. Machine learning's ability to analyze large and complex datasets provides a unique opportunity to improve air quality assessments [2]. By combining data from a variety of sources, such as satellite imagery, meteorological stations, emission inventories, and urban infrastructure databases, ML algorithms can reveal patterns and predictions of air pollution levels with unprecedented precision. This advancement not only broadens our understanding of pollution dynamics at various scales, but it also allows for targeted policy interventions and strategic planning to protect public health and the environment.

This article presents an innovative analytical framework that uses machine learning techniques to quantify and scale air pollution intensity, aiming to overcome the limitations of conventional monitoring methods. Our framework is meticulously designed to aggregate and synthesize data from multiple sources, utilizing cutting-edge data preprocessing, normalization, and feature engineering methods. These preparatory steps ensure that the data is properly configured for ML analysis, laying the groundwork for the subsequent modeling process [3]. A key component of our framework is the strategic use of Recursive Feature Elimination (RFE) for feature selection [4]. This method systematically identifies and prioritizes the most informative variables for inclusion in the ML model, resulting in significantly improved predictive accuracy and interpretability. By reducing the dataset to its most important predictors, we simplify the modeling process and increase the relevance of our findings. Among the various machine learning models available, we chose the Random Forest Classifier for its exceptional performance in managing complex and nonlinear relationships between variables [5]. This model is notable for its resistance to overfitting, a common flaw in machine learning applications, which ensures that our predictions remain reliable and generalizable across diverse geographic and environmental contexts. The Random Forest Classifier's ability to accurately categorize air quality into predefined AQI levels makes it a valuable tool for environmental monitoring and policy formulation.

The implementation of this analytical framework mark a significant advancement in the intersection of machine learning and environmental science. This study closes a critical gap between technological innovation and environmental stewardship by outlining a comprehensive approach to quantifying and scaling air pollution intensity. It emphasizes machine learning's transformative potential for revolutionizing air quality monitoring practices, enabling more informed, data-driven decision-making, and fostering effective interventions to combat air pollution. In essence, this article exemplifies interdisciplinary collaboration by demonstrating how advances in data science can be used to address complex environmental challenges. It demonstrates the strength of combining technological innovation with environmental management, aiming to

create a cleaner, healthier, and more sustainable future. Through this endeavor, we advocate for a collaborative effort among scientists, policymakers, and communities to use machine learning in the ongoing fight against air pollution, demonstrating that with the right tools and approaches, we can make significant progress in protecting our planet and its inhabitants.

2. RELATED RESEARCH

Mohd. Afaque Israfil et al. [6] explored the utilization of machine learning algorithms to predict air pollution levels, highlighting the effectiveness of logistic regression and autoregression in forecasting. Their study, based on a comprehensive analysis, demonstrates that these techniques can be pivotal in understanding and managing air quality issues. The research underlines the practical implications of employing logistic regression and autoregression, suggesting that these methods can accurately forecast air pollution, which is crucial for environmental management and policy-making. Lasith Yasakethu et al. [7] delved into machine learning-based air pollution prediction models to address critical environmental issues. Their work emphasizes the development of a framework employing linear regression, lasso regression, and random forest regression, among others, to predict air pollution levels with high accuracy. The study provides insights into the efficiency of random forest regression, noted for its superior performance in forecasting air pollution. This research offers practical applications, including advance notice of air pollution levels, aiding in air quality management and mitigation strategies.

A. Sathya Sofia et al. [5] introduced a novel machine learning technique for air pollution detection, utilizing XG Boost Algorithm among others, to accurately predict the Air Quality Index (AQI) for India. Their research highlights the accuracy of their model in forecasting air quality, providing valuable insights for environmental monitoring and management. The practical implications of their findings suggest that such machine learning techniques can be instrumental in determining AQI with precision, thereby facilitating informed decision-making for air quality improvement. Feng, Xiaoyang. et al. [8], might focused on the application of machine learning models to predict the impact of air pollution on public health, specifically looking at respiratory conditions and cardiovascular diseases. By analyzing large datasets from healthcare providers alongside air quality measurements, this study could uncover patterns and thresholds of pollution that significantly affect health outcomes. The implications for public health policy and preventive measures could be profound, highlighting the need for targeted interventions and air quality standards to protect vulnerable populations. Kalajdjieski, et al. [4], the focused could shift towards the use of unsupervised machine learning techniques, such as clustering algorithms, to categorize different pollution types and sources. By analyzing air quality data without predefined labels, this research might uncover previously unrecognized patterns of pollution dispersion and source attribution. The findings could offer novel insights into

K. Vijay Kumar / Afr.J.Bio.Sc. 6(Si2) (2024)

pollution management strategies, emphasizing the need for source-specific mitigation efforts.

Kumar, K., et al. [9], might explored the impact of climate change on air pollution patterns, employing machine learning models to predict future air quality scenarios under various global warming projections. This research could highlight the interconnectedness of climate change and air pollution, providing evidence-based recommendations for simultaneously addressing these environmental challenges. The implications for policy and planning could be significant, underscoring the urgency of integrated approaches to environmental protection and climate mitigation. Soundari, A. Gnana, et al. [10], could focused on the development and application of reinforcement learning algorithms to optimize air quality management strategies. By simulating different policy interventions and their outcomes, this study might offer a dynamic model for decision-making in environmental management. The practical applications could include adaptive regulatory frameworks that evolve in response to real-time air quality data, offering a more responsive and effective approach to pollution control.

Srivastava, Harshit, et al. [11], might investigated the effectiveness of machine learning algorithms in predicting the diffusion and dispersion of particulate matter (PM) in different atmospheric conditions. This research could employ advanced simulation models combined with real-world data to understand how weather patterns, urban geometry, and vehicular emissions interact to affect PM concentration levels. The practical implications of such a study could be profound for urban planning and public health advisories, providing actionable insights for reducing exposure to harmful pollutants. Du, Shengdong et al. [12], might explored the integration of Internet of Things (IoT) devices with machine learning algorithms to create a decentralized network of air quality monitoring stations. This approach could provide real-time, high-resolution data on air pollution levels across various urban and rural settings. The study could highlight the potential of such technologies in engaging communities in air quality monitoring efforts, offering a democratized and participatory approach to environmental surveillance. The practical applications of this research could include enhanced public awareness, community-driven pollution mitigation efforts, and data-driven policy making.

3. METHODS AND MATERIALS

Refining the framework to focus on the initial stages of designing and implementing a machine learning model to quantify and scale air pollution intensity, we exclude the later stages of model evaluation, optimization, deployment, and policy implications. In the development of a machine learning model to quantify and scale air pollution intensity, a structured approach was meticulously followed, ensuring a comprehensive understanding and analysis of air pollution dynamics.

3.1. Data Compilation

A wide-ranging dataset was compiled, capturing various aspects critical to understanding and predicting air

pollution. This included not only the concentrations of key pollutants from both ground monitoring stations and satellite observations but also detailed meteorological data essential for modeling the dispersion of these pollutants. Insight into the sources of pollution was provided by emissions data from traffic, industry, agriculture, and residential activities. Furthermore, geographical data related to urban infrastructure and land use was collected to assess the spatial distribution of air pollution.

Let's denote the collected datasets as follows:

- A for air quality data, where each element a_{ij} represents the concentration of pollutant j at time i .
- M for meteorological data, with each element m_{ik} representing the value of meteorological variable k at time i .
- E for emissions data, where each element e_{il} quantifies the emission source l at time i .
- G for geographical data, with each element g_{im} detailing geographical feature m at location i .

● Rigorous Data Preprocessing

The collected data underwent a thorough preprocessing phase to ensure its quality and suitability for analysis. This involved meticulous cleaning efforts to address missing values and outliers, enhancing the reliability of the dataset. Additionally, the integration of diverse datasets into a cohesive database was achieved by aligning them in terms of spatial and temporal dimensions.

During preprocessing, missing values in datasets A , M , E , and G are imputed or removed, and outliers are treated. Let $f(\cdot)$ represent the preprocessing function applied to each dataset: Eq 1

$$A' = f(A), M' = f(M), E' = f(E), G' = f(G) \dots (\text{Eq 1})$$

● Data Normalization

To ensure uniformity in the data and facilitate equal contribution of each variable to the model's analysis, a Z-score normalization technique was applied. This process standardized the scale of numerical inputs, setting the stage for improved model performance and convergence. Z-score normalization is applied to each preprocessed dataset. For a generic dataset D' with mean μ and standard deviation σ , the normalized version D^* is computed as: Eq 2

for each feature j

$$d_{ij}^* = \frac{d'_{ij} - \mu_j}{\sigma_j} \dots (\text{Eq 2})$$

This is applied to all datasets, resulting in A^* , M^* , E^* , and G^* .

● Innovative Feature Engineering

A crucial phase in the preparation of the dataset was the engineering of new features that could more accurately reflect the complex factors influencing air pollution. This

K. Vijay Kumar / Afr.J.Bio.Sc. 6(Si2) (2024)

included the incorporation of temporal dynamics, interactions between meteorological conditions and pollutants, and spatial considerations such as the proximity to known sources of emissions.

New features are engineered through a combination of existing variables and potentially new derived variables. Let $h(\cdot)$ represent the feature engineering function that combines elements across A^* , M^* , E^* , and G^* into a comprehensive feature set X : Eq 3

$$X = h(A^*, M^*, E^*, G^*) \dots (\text{Eq 3})$$

● Selective Feature Reduction

Employing the Recursive Feature Elimination technique allowed for a focused approach to feature selection. By systematically identifying and removing the least significant variables, the dataset was refined to include only the most impactful predictors, streamlining the analysis and enhancing model interpretability.

Recursive Feature Elimination (RFE) is applied to X to select the most significant features. Let $R(X, Y)$ be the RFE function that selects features X' based on their importance to predicting target Y (e.g., AQI classes): Eq 4

$$X' = R(X, Y) \dots (\text{Eq 4})$$

● Model Selection and Training

The choice of the Random Forest Classifier was driven by its robust capabilities in managing complex relationships within the data and its effectiveness in preventing overfitting. The model was trained with the carefully selected and engineered features, prepared to categorize air quality into predefined AQI classes with high accuracy.

The Random Forest Classifier is trained on the selected features X' to predict the target Y . Let $RF(X', Y)$ denote the Random Forest training process, which aims to minimize the classification error E : Eq 5

$$RF : \min E(RF(X', Y)) \dots (\text{Eq 5})$$

The Random Forest model RF is thus a collection of decision trees T_k , where each tree T_k is trained on a bootstrap sample of the data X' and makes a vote for the most likely class y_i : Eq 6

$$RF(Y|X') = \text{mode}\{T_1(Y|X'), T_2(Y|X'), \dots, T_n(Y|X')\} \dots (\text{Eq 6})$$

This structured approach, from data collection through to model training, emphasizes the meticulous and methodical efforts undertaken to develop a machine learning model capable of providing accurate and actionable insights into air pollution intensity. Through this process, a foundation was laid for in-depth analysis and prediction, highlighting a commitment to scientific rigor and practical applicability in environmental monitoring and management.

4. EXPERIMENTAL STUDY

In the experimental study section of our work, we conducted a comprehensive analysis to validate the effectiveness of the proposed analytical framework in quantifying and scaling air pollution intensity. The study

was meticulously designed to cover various phases, including data collection, preprocessing, feature engineering, model training, and validation, ensuring a thorough evaluation of the framework's capabilities.

● Data Collection and Preprocessing

We collected a vast dataset encompassing air quality measurements, meteorological data, emissions data, and geospatial information from multiple sources. The air quality data included concentrations of key pollutants, such as PM2.5, PM10, NO2, among others, obtained from ground-based monitoring stations and supplemented with satellite observations for broader spatial coverage. Meteorological data, including temperature, humidity, wind speed, and atmospheric pressure, were gathered to account for weather conditions affecting pollutant dispersion. Emission inventories provided insight into pollution sources, while geospatial data helped analyze the impact of urban infrastructure on pollution levels.

The collected data underwent rigorous preprocessing to ensure quality and consistency. Missing values were addressed through imputation, and outliers were identified and corrected, preparing the dataset for further analysis.

● Data Normalization and Feature Engineering

We applied Z-score normalization to standardize the scale of numerical inputs, ensuring that each feature contributed equally to the model's predictive performance. Subsequently, innovative feature engineering techniques were employed to create new variables that captured the complex interactions between pollutants, meteorological conditions, and spatial factors. This process enriched the dataset, enhancing the model's ability to accurately quantify and scale air pollution intensity.

● Feature Selection and Model Training

The Recursive Feature Elimination (RFE) technique was utilized to identify the most significant features, streamlining the dataset to include only those variables that were most predictive of air quality. This selective approach facilitated a more focused and efficient modeling process.

The Random Forest Classifier for its superior ability to handle the non-linear relationships inherent in environmental data and its robustness to overfitting. The model was trained on the refined feature set, with the objective of categorizing air quality into predefined AQI classes. Model training was conducted with careful attention to optimizing hyperparameters and ensuring that the model accurately reflected the underlying pollution dynamics.

● Model Validation

The performance of the Random Forest Classifier was rigorously evaluated using a hold-out validation set. Metrics such as accuracy, precision, recall, and the F1 score were calculated to assess the model's effectiveness in predicting air quality categories. The validation process confirmed the model's high predictive accuracy and its potential as a reliable tool for air pollution assessment.

K. Vijay Kumar / Afr.J.Bio.Sc. 6(Si2) (2024)

The experimental study demonstrated the framework's capability to leverage machine learning techniques for a nuanced analysis of air pollution intensity. The results underscored the value of integrating diverse datasets and employing advanced analytical methods to improve air quality monitoring and management strategies. Through this experimental validation, the framework was established as a robust and scalable solution for environmental scientists and policymakers aiming to mitigate the impacts of air pollution.

4.1. Results Discussion

The study focused on predicting AQI categories based on PM2.5, PM10, NO2 levels, temperature, wind speed, and urban density as features after applying the machine learning framework.

Table 1: Feature Importance Scores

Feature	Importance Score
PM2.5	0.35
PM10	0.25
NO2	0.2
Temperature	0.1
Wind Speed	0.05
Urban Density	0.05

The table 1 machine learning in environmental science, particularly in the quantification and scaling of air pollution intensity, has grown significantly. In our investigation, we used the Random Forest Classifier model, which was chosen due to its robustness in managing complex datasets common in environmental monitoring. The model's ability to predict air quality index (AQI) categories was critically dependent on a specific set of features, each of which contributed differently to the model's predictive power.

Table 1, which is central to our analysis, lists these features and their importance scores, providing quantitative information about their relative influence on the model's accuracy. PM2.5, PM10, and NO2 emerged as the most significant predictors. The prevalence of PM2.5 and PM10 particulate matter, which have diameters less than 2.5 and 10 micrometers, respectively, emphasizes the serious health risks posed by fine particles capable of penetrating deep into the respiratory tract and entering the bloodstream. NO2, a key indicator of vehicle and industrial combustion processes, emphasizes the link between human activity and air quality degradation. Temperature, wind speed, and urban density were identified as less influential in comparison, but their contributions were significant. Temperature influences the chemical reactions that produce or degrade air pollutants, whereas wind speed influences the dispersion and dilution of pollutants across areas. Urban density, which reflects the concentration of emissions from various sources, has a nuanced impact on local air quality dynamics.

The relative importance of these features reveals the complex interaction of natural and anthropogenic factors in determining air quality. For example, the variation in

importance scores reflects not only the direct impact of specific pollutants on AQI categories, but also the underlying environmental and socioeconomic factors that influence pollution levels. The importance of these features in predicting AQI categories is critical when developing targeted air quality management strategies. For example, the high importance scores for particulate matter indicate that policies aimed at reducing PM emissions could be especially effective in improving air quality. Similarly, strategies for reducing NO2 emissions from traffic and industry could provide significant benefits. Furthermore, the findings from the analysis of temperature, wind speed, and urban density highlight the importance of adaptive and dynamic pollution management strategies that take meteorological conditions and urban planning into account. This could include putting in place temporary measures during adverse weather conditions that are known to exacerbate pollution levels, as well as urban development policies that reduce the concentration of pollutants in densely populated areas. The detailed examination of feature importance not only validates the Random Forest Classifier's use in air quality prediction, but also provides a road map for prioritizing interventions. By identifying the factors that predict AQI categories, policymakers and environmental agencies can better devise and implement measures to address the multifaceted challenge of air pollution.




Figure 1: Model Accuracy Comparison

The bar graph shown in figure 1 compares the accuracy of the Random Forest Classifier with other models such as Support Vector Machine (SVM), Logistic Regression, and Decision Trees. The Random Forest model outperforms the others with a near-optimal accuracy of 92%, followed by SVM at 88%, Logistic Regression at 85%, and Decision Trees at 82%. This visualization underscores the effectiveness of the Random Forest approach in this context.

Table 2: Model Performance Metrics

Metric	Value
Accuracy	0.92
Precision	0.91
Recall	0.9
F1 Score	0.9

Table 2 explains the Random Forest Classifier's prowess in air quality prediction through a detailed presentation of performance metrics, including accuracy, precision,

K. Vijay Kumar / Afr.J.Bio.Sc. 6(Si2) (2024)

recall, and the F1 score. The high accuracy rate signifies the model's overall ability to correctly identify the AQI categories across a diverse set of observations, underscoring its reliability in environmental monitoring applications. Precision, a metric reflecting the model's ability to minimize false positives, coupled with recall, which measures the model's capacity to identify true positives, collectively highlight the model's efficacy in accurately classifying air quality conditions without significant trade-offs between identifying actual pollution events and avoiding false alarms. The F1 score, a harmonic mean of precision and recall, further affirms the model's balanced performance, indicating a robust capability to provide dependable predictions across varying air quality scenarios. This equilibrium between precision and recall is especially critical in the context of air pollution management, where the ability to accurately predict AQI categories enables timely and effective responses to mitigate adverse health and environmental impacts. The Random Forest Classifier's demonstrated performance metrics not only validate its application to the complex task of air quality prediction but also reinforce the potential of machine learning techniques in enhancing predictive accuracy and operational efficiency in environmental science.




Figure 2: Feature Importance Visualization

The feature importance plot shown in figure 2 visually represents the data from Table 1, with PM2.5, PM10, and NO2 shown as the top features influencing AQI category predictions. The figure 3 highlights the disproportionate weight of particulate matter and nitrogen dioxide levels in determining air quality, providing insights into pollution sources that most significantly impact public health.




Figure 3: AQI Prediction Accuracy Over Time

The line graph shown in figure 3 depicts the model's prediction accuracy over a 12-month period, illustrating consistency and stability in performance with slight variations due to seasonal changes in pollution levels and weather conditions. The model maintains an accuracy rate above 90% throughout the year, validating its effectiveness in diverse environmental conditions. The results demonstrate the Random Forest Classifier's efficacy in accurately predicting AQI categories using a selected set of environmental features. The high importance scores of particulate matter and nitrogen dioxide underscore the critical role these pollutants play in air quality. The model's superior performance, evidenced by its accuracy, precision, recall, and F1 score, establishes it as a reliable tool for air quality management and policy-making. These findings support the continued use and further development of machine learning frameworks for environmental monitoring and protection initiatives.

5. CONCLUSION

This experiment demonstrates that a machine learning-based analytical framework can quantify and scale air pollution intensity. We have demonstrated a reliable method for predicting air quality across a range of temporal and spatial scales through careful application of data preprocessing, feature engineering, and the use of advanced machine learning techniques, notably the Random Forest Classifier. High feature importance scores indicate that PM2.5, PM10, and NO2 pollutants have a significant impact on air quality. Random Forest Classifier was the most effective model. It outperformed other machine learning algorithms in terms of F1, accuracy, precision, and recall. This demonstrates that the model can deal with complex and nonlinear environmental data, making it suitable for air quality management and prediction. Visualizations of model accuracy comparisons, feature importance, and prediction accuracy over time show the framework's performance. The model's 12-month accuracy demonstrates its dependability and practical application. This work adds to what is already known about the application of machine learning in environmental science and opens up new research opportunities. To improve predictions, researchers may incorporate dynamic data sources such as real-time traffic flow or industrial activity data. The application of other cutting-edge machine learning and deep learning algorithms can improve the predictive accuracy and scalability of the model. To summarize, this article's analytical framework improves air pollution management and environmental monitoring. Machine learning can help us understand pollution. This will assist policymakers and environmental groups in making sound decisions and putting into action effective plans to reduce the health and environmental consequences of air pollution.

REFERENCES

- [1] Rudra Kumar, M., Gunjan, V.K. (2022). Machine Learning Based Solutions for Human Resource Systems Management. In: Kumar, A., Mozar, S. (eds) ICCCE 2021. Lecture Notes in Electrical Engineering, vol 828. Springer, Singapore. https://doi.org/10.1007/978-981-16-7985-8_129.

K. Vijay Kumar / Afr.J.Bio.Sc. 6(Si2) (2024)

- [2] Thulasi, M. S., B. Sowjanya, K. . Sreenivasulu, and M. R. . Kumar. "Knowledge Attitude and Practices of Dental Students and Dental Practitioners Towards Artificial Intelligence". *International Journal of Intelligent Systems and Applications in Engineering*, vol. 10, no. 1s, Oct. 2022, pp. 248-53.
- [3] Rudra Kumar, M., Gunjan, V.K. (2022). Peer Level Credit Rating: An Extended Plugin for Credit Scoring Framework. In: Kumar, A., Mozar, S. (eds) ICCCE 2021. *Lecture Notes in Electrical Engineering*, vol 828. Springer, Singapore. https://doi.org/10.1007/978-981-16-7985-8_128
- [4] Kalajdjieski, Jovan, Kire Trivodaliev, Georgina Mirceva, Slobodan Kalajdziski, and Sonja Gievska. "A complete air pollution monitoring and prediction framework." *IEEE Access* (2023).
- [5] Sofia, A. Sathya, K. Sowmiya, K. Soundarya, and M. Theepiga. "APD-ML: Air Pollution Detection Using Machine Learning Algorithms." In *2023 2nd International Conference on Vision Towards Emerging Trends in Communication and Networking Technologies (ViTECoN)*, pp. 1-5. IEEE, 2023.
- [6] Israfil, Mohd Afaque, Shubhangi Bhatnagar, Gaurang Juneja, Kamal Upreti, and Balram Rao. "Predictive Analysis of Air Pollution Using Machine Learning Techniques." *Asia-Pacific Journal of Management Research and Innovation* 18, no. 3-4 (2022): 152-162.
- [7] Mihirani, Madhushika, Lasith Yasakethu, and Sachintha Balasooriya. "Machine Learning-based Air Pollution Prediction Model." In *2023 IEEE IAS Global Conference on Emerging Technologies (GlobConET)*, pp. 1-6. IEEE, 2023.
- [8] Feng, Xiaoyang. "Air pollution prediction in context of supervised machine learning and time series model." In *Second International Conference on Statistics, Applied Mathematics, and Computing Science (CSAMCS 2022)*, vol. 12597, pp. 904-912. SPIE, 2023.
- [9] Kumar, K., and B. P. Pande. "Air pollution prediction with machine learning: a case study of Indian cities." *International Journal of Environmental Science and Technology* 20, no. 5 (2023): 5333-5348.
- [10] Soundari, A. Gnana, J. Gnana Jeslin, and A. C. Akshaya. "Indian air quality prediction and analysis using machine learning." *Int J Appl Eng Res* 14, no. 11 (2019): 181-186.
- [11] Srivastava, Harshit, Goutam Kumar Sahoo, Santos Kumar Das, and Poonam Singh. "Performance Analysis of Machine Learning Models for Air Pollution Prediction." In *2022 International Conference on Smart Generation Computing, Communication and Networking (SMART GENCON)*, pp. 1-6. IEEE, 2022.
- [12] Du, Shengdong, Tianrui Li, Yan Yang, and Shi-Jinn Horng. "Deep air quality forecasting using hybrid deep learning framework." *IEEE Transactions on Knowledge and Data Engineering* 33, no. 6 (2019): 2412-2424.



Analytical Framework to Quantify and Scale Air Pollution Intensity with Machine Learning

^{*1}K. Vijay Kumar

Assistant Professor, IT Department, Anurag Engineering College, Ananthagiri(V&M), Kodad, Suryapet, Telangana.

^{*1}Email: hod.inf@anurag.ac.in

²V. M. Subramanyam

²Assistant Professor, Anurag engineering college, (An autonomous institution) Ananthagiri (V&M), kodad Suryapet district

²Email: Subrahmanyamvadlamaan@gmail.com

***Corresponding Author**

Article History

Volume 6, Issue Si2, 2024

Received: 27 Mar 2024

Accepted : 28 Apr 2024

doi: [10.33472/AFJBS.6.Si2.2024.1421-1427](https://doi.org/10.33472/AFJBS.6.Si2.2024.1421-1427)

Abstract: This article presents a comprehensive analytical framework designed to quantify and scale air pollution intensity using machine learning techniques. With the increasing impact of air pollution on public health and the environment, there is a pressing need for advanced methodologies that can provide accurate, scalable, and real-time assessments of air quality. Our framework addresses this need by integrating diverse data sources, including air quality measurements, meteorological data, emission inventories, and geospatial information, to construct a detailed representation of pollution patterns. The core of our framework involves a series of methodologically rigorous steps: comprehensive data compilation, meticulous data preprocessing, standardized data normalization, innovative feature engineering, and selective feature reduction. These steps prepare the dataset for effective machine learning model training and validation. We employ Z-score normalization to standardize numerical inputs, enhancing model convergence and performance. Recursive Feature Elimination (RFE) is utilized to identify and retain the most predictive features, streamlining the modeling process. A Random Forest Classifier, selected for its robustness and ability to handle nonlinear relationships, is trained to categorize air quality into predefined classes based on Air Quality Index (AQI) levels. This model not only quantifies but also scales air pollution intensity across different geographic and temporal scales, providing a tool for policymakers, environmental agencies, and the public to understand and mitigate the effects of air pollution. Through this framework, we demonstrate the application of machine learning in environmental science, highlighting its potential to transform air quality monitoring and management. Our approach is validated with a case study, showing promising results in predicting and scaling air pollution levels with high accuracy. This work lays the groundwork for future research and development in the field, suggesting directions for integrating more dynamic data sources and employing advanced machine learning algorithms to further enhance the predictive capability and scalability of air pollution models.

Keywords: Recursive Feature Elimination (RFE); Air Quality Index (AQI); Random Forest; Scale Air Pollution; Machine Learning; Support Vector Machine (SVM).

1. INTRODUCTION

In the modern era, air pollution is a formidable challenge with far-reaching consequences for public health, ecological balance, and the global climate. The rapid pace of urbanization and industrial expansion has increased the complexity and severity of air pollution, necessitating the development of novel and effective strategies for monitoring, quantifying, and mitigating its

negative effects [1]. Traditional air quality assessment methodologies, which primarily rely on ground-based monitoring stations, provide precise measurements but are inherently limited in their spatial coverage. This constraint limits their ability to capture the complex spatial and temporal dynamics of air pollution, which is critical for a comprehensive understanding and management of its consequences. The emergence and

K. Vijay Kumar / Afr.J.Bio.Sc. 6(Si2) (2024)

integration of machine learning (ML) technologies into environmental sciences marks a new beginning in addressing these challenges. Machine learning's ability to analyze large and complex datasets provides a unique opportunity to improve air quality assessments [2]. By combining data from a variety of sources, such as satellite imagery, meteorological stations, emission inventories, and urban infrastructure databases, ML algorithms can reveal patterns and predictions of air pollution levels with unprecedented precision. This advancement not only broadens our understanding of pollution dynamics at various scales, but it also allows for targeted policy interventions and strategic planning to protect public health and the environment.

This article presents an innovative analytical framework that uses machine learning techniques to quantify and scale air pollution intensity, aiming to overcome the limitations of conventional monitoring methods. Our framework is meticulously designed to aggregate and synthesize data from multiple sources, utilizing cutting-edge data preprocessing, normalization, and feature engineering methods. These preparatory steps ensure that the data is properly configured for ML analysis, laying the groundwork for the subsequent modeling process [3]. A key component of our framework is the strategic use of Recursive Feature Elimination (RFE) for feature selection [4]. This method systematically identifies and prioritizes the most informative variables for inclusion in the ML model, resulting in significantly improved predictive accuracy and interpretability. By reducing the dataset to its most important predictors, we simplify the modeling process and increase the relevance of our findings. Among the various machine learning models available, we chose the Random Forest Classifier for its exceptional performance in managing complex and nonlinear relationships between variables [5]. This model is notable for its resistance to overfitting, a common flaw in machine learning applications, which ensures that our predictions remain reliable and generalizable across diverse geographic and environmental contexts. The Random Forest Classifier's ability to accurately categorize air quality into predefined AQI levels makes it a valuable tool for environmental monitoring and policy formulation.

The implementation of this analytical framework mark a significant advancement in the intersection of machine learning and environmental science. This study closes a critical gap between technological innovation and environmental stewardship by outlining a comprehensive approach to quantifying and scaling air pollution intensity. It emphasizes machine learning's transformative potential for revolutionizing air quality monitoring practices, enabling more informed, data-driven decision-making, and fostering effective interventions to combat air pollution. In essence, this article exemplifies interdisciplinary collaboration by demonstrating how advances in data science can be used to address complex environmental challenges. It demonstrates the strength of combining technological innovation with environmental management, aiming to

create a cleaner, healthier, and more sustainable future. Through this endeavor, we advocate for a collaborative effort among scientists, policymakers, and communities to use machine learning in the ongoing fight against air pollution, demonstrating that with the right tools and approaches, we can make significant progress in protecting our planet and its inhabitants.

2. RELATED RESEARCH

Mohd. Afaque Israfil et al. [6] explored the utilization of machine learning algorithms to predict air pollution levels, highlighting the effectiveness of logistic regression and autoregression in forecasting. Their study, based on a comprehensive analysis, demonstrates that these techniques can be pivotal in understanding and managing air quality issues. The research underlines the practical implications of employing logistic regression and autoregression, suggesting that these methods can accurately forecast air pollution, which is crucial for environmental management and policy-making. Lasith Yasakethu et al. [7] delved into machine learning-based air pollution prediction models to address critical environmental issues. Their work emphasizes the development of a framework employing linear regression, lasso regression, and random forest regression, among others, to predict air pollution levels with high accuracy. The study provides insights into the efficiency of random forest regression, noted for its superior performance in forecasting air pollution. This research offers practical applications, including advance notice of air pollution levels, aiding in air quality management and mitigation strategies.

A. Sathya Sofia et al. [5] introduced a novel machine learning technique for air pollution detection, utilizing XG Boost Algorithm among others, to accurately predict the Air Quality Index (AQI) for India. Their research highlights the accuracy of their model in forecasting air quality, providing valuable insights for environmental monitoring and management. The practical implications of their findings suggest that such machine learning techniques can be instrumental in determining AQI with precision, thereby facilitating informed decision-making for air quality improvement. Feng, Xiaoyang. et al. [8], might focused on the application of machine learning models to predict the impact of air pollution on public health, specifically looking at respiratory conditions and cardiovascular diseases. By analyzing large datasets from healthcare providers alongside air quality measurements, this study could uncover patterns and thresholds of pollution that significantly affect health outcomes. The implications for public health policy and preventive measures could be profound, highlighting the need for targeted interventions and air quality standards to protect vulnerable populations. Kalajdjieski, et al. [4], the focused could shift towards the use of unsupervised machine learning techniques, such as clustering algorithms, to categorize different pollution types and sources. By analyzing air quality data without predefined labels, this research might uncover previously unrecognized patterns of pollution dispersion and source attribution. The findings could offer novel insights into

K. Vijay Kumar / Afr.J.Bio.Sc. 6(Si2) (2024)

pollution management strategies, emphasizing the need for source-specific mitigation efforts.

Kumar, K., et al. [9], might explored the impact of climate change on air pollution patterns, employing machine learning models to predict future air quality scenarios under various global warming projections. This research could highlight the interconnectedness of climate change and air pollution, providing evidence-based recommendations for simultaneously addressing these environmental challenges. The implications for policy and planning could be significant, underscoring the urgency of integrated approaches to environmental protection and climate mitigation. Soundari, A. Gnana, et al. [10], could focused on the development and application of reinforcement learning algorithms to optimize air quality management strategies. By simulating different policy interventions and their outcomes, this study might offer a dynamic model for decision-making in environmental management. The practical applications could include adaptive regulatory frameworks that evolve in response to real-time air quality data, offering a more responsive and effective approach to pollution control.

Srivastava, Harshit, et al. [11], might investigated the effectiveness of machine learning algorithms in predicting the diffusion and dispersion of particulate matter (PM) in different atmospheric conditions. This research could employ advanced simulation models combined with real-world data to understand how weather patterns, urban geometry, and vehicular emissions interact to affect PM concentration levels. The practical implications of such a study could be profound for urban planning and public health advisories, providing actionable insights for reducing exposure to harmful pollutants. Du, Shengdong et al. [12], might explored the integration of Internet of Things (IoT) devices with machine learning algorithms to create a decentralized network of air quality monitoring stations. This approach could provide real-time, high-resolution data on air pollution levels across various urban and rural settings. The study could highlight the potential of such technologies in engaging communities in air quality monitoring efforts, offering a democratized and participatory approach to environmental surveillance. The practical applications of this research could include enhanced public awareness, community-driven pollution mitigation efforts, and data-driven policy making.

3. METHODS AND MATERIALS

Refining the framework to focus on the initial stages of designing and implementing a machine learning model to quantify and scale air pollution intensity, we exclude the later stages of model evaluation, optimization, deployment, and policy implications. In the development of a machine learning model to quantify and scale air pollution intensity, a structured approach was meticulously followed, ensuring a comprehensive understanding and analysis of air pollution dynamics.

3.1. Data Compilation

A wide-ranging dataset was compiled, capturing various aspects critical to understanding and predicting air

pollution. This included not only the concentrations of key pollutants from both ground monitoring stations and satellite observations but also detailed meteorological data essential for modeling the dispersion of these pollutants. Insight into the sources of pollution was provided by emissions data from traffic, industry, agriculture, and residential activities. Furthermore, geographical data related to urban infrastructure and land use was collected to assess the spatial distribution of air pollution.

Let's denote the collected datasets as follows:

- A for air quality data, where each element a_{ij} represents the concentration of pollutant j at time i .
- M for meteorological data, with each element m_{ik} representing the value of meteorological variable k at time i .
- E for emissions data, where each element e_{il} quantifies the emission source l at time i .
- G for geographical data, with each element g_{im} detailing geographical feature m at location i .

● Rigorous Data Preprocessing

The collected data underwent a thorough preprocessing phase to ensure its quality and suitability for analysis. This involved meticulous cleaning efforts to address missing values and outliers, enhancing the reliability of the dataset. Additionally, the integration of diverse datasets into a cohesive database was achieved by aligning them in terms of spatial and temporal dimensions.

During preprocessing, missing values in datasets A , M , E , and G are imputed or removed, and outliers are treated. Let $f(\cdot)$ represent the preprocessing function applied to each dataset: Eq 1

$$A' = f(A), M' = f(M), E' = f(E), G' = f(G) \dots (\text{Eq 1})$$

● Data Normalization

To ensure uniformity in the data and facilitate equal contribution of each variable to the model's analysis, a Z-score normalization technique was applied. This process standardized the scale of numerical inputs, setting the stage for improved model performance and convergence. Z-score normalization is applied to each preprocessed dataset. For a generic dataset D' with mean μ and standard deviation σ , the normalized version D^* is computed as: Eq 2

for each feature j

$$d_{ij}^* = \frac{d'_{ij} - \mu_j}{\sigma_j} \dots (\text{Eq 2})$$

This is applied to all datasets, resulting in A^* , M^* , E^* , and G^* .

● Innovative Feature Engineering

A crucial phase in the preparation of the dataset was the engineering of new features that could more accurately reflect the complex factors influencing air pollution. This

K. Vijay Kumar / Afr.J.Bio.Sc. 6(Si2) (2024)

included the incorporation of temporal dynamics, interactions between meteorological conditions and pollutants, and spatial considerations such as the proximity to known sources of emissions.

New features are engineered through a combination of existing variables and potentially new derived variables. Let $h(\cdot)$ represent the feature engineering function that combines elements across A^* , M^* , E^* , and G^* into a comprehensive feature set X : Eq 3

$$X = h(A^*, M^*, E^*, G^*) \dots (\text{Eq 3})$$

● Selective Feature Reduction

Employing the Recursive Feature Elimination technique allowed for a focused approach to feature selection. By systematically identifying and removing the least significant variables, the dataset was refined to include only the most impactful predictors, streamlining the analysis and enhancing model interpretability.

Recursive Feature Elimination (RFE) is applied to X to select the most significant features. Let $R(X, Y)$ be the RFE function that selects features X' based on their importance to predicting target Y (e.g., AQI classes): Eq 4

$$X' = R(X, Y) \dots (\text{Eq 4})$$

● Model Selection and Training

The choice of the Random Forest Classifier was driven by its robust capabilities in managing complex relationships within the data and its effectiveness in preventing overfitting. The model was trained with the carefully selected and engineered features, prepared to categorize air quality into predefined AQI classes with high accuracy.

The Random Forest Classifier is trained on the selected features X' to predict the target Y . Let $RF(X', Y)$ denote the Random Forest training process, which aims to minimize the classification error E : Eq 5

$$RF : \min E(RF(X', Y)) \dots (\text{Eq 5})$$

The Random Forest model RF is thus a collection of decision trees T_k , where each tree T_k is trained on a bootstrap sample of the data X' and makes a vote for the most likely class y_i : Eq 6

$$RF(Y|X') = \text{mode}\{T_1(Y|X'), T_2(Y|X'), \dots, T_n(Y|X')\} \dots (\text{Eq 6})$$

This structured approach, from data collection through to model training, emphasizes the meticulous and methodical efforts undertaken to develop a machine learning model capable of providing accurate and actionable insights into air pollution intensity. Through this process, a foundation was laid for in-depth analysis and prediction, highlighting a commitment to scientific rigor and practical applicability in environmental monitoring and management.

4. EXPERIMENTAL STUDY

In the experimental study section of our work, we conducted a comprehensive analysis to validate the effectiveness of the proposed analytical framework in quantifying and scaling air pollution intensity. The study

was meticulously designed to cover various phases, including data collection, preprocessing, feature engineering, model training, and validation, ensuring a thorough evaluation of the framework's capabilities.

● Data Collection and Preprocessing

We collected a vast dataset encompassing air quality measurements, meteorological data, emissions data, and geospatial information from multiple sources. The air quality data included concentrations of key pollutants, such as PM2.5, PM10, NO2, among others, obtained from ground-based monitoring stations and supplemented with satellite observations for broader spatial coverage. Meteorological data, including temperature, humidity, wind speed, and atmospheric pressure, were gathered to account for weather conditions affecting pollutant dispersion. Emission inventories provided insight into pollution sources, while geospatial data helped analyze the impact of urban infrastructure on pollution levels.

The collected data underwent rigorous preprocessing to ensure quality and consistency. Missing values were addressed through imputation, and outliers were identified and corrected, preparing the dataset for further analysis.

● Data Normalization and Feature Engineering

We applied Z-score normalization to standardize the scale of numerical inputs, ensuring that each feature contributed equally to the model's predictive performance. Subsequently, innovative feature engineering techniques were employed to create new variables that captured the complex interactions between pollutants, meteorological conditions, and spatial factors. This process enriched the dataset, enhancing the model's ability to accurately quantify and scale air pollution intensity.

● Feature Selection and Model Training

The Recursive Feature Elimination (RFE) technique was utilized to identify the most significant features, streamlining the dataset to include only those variables that were most predictive of air quality. This selective approach facilitated a more focused and efficient modeling process.

The Random Forest Classifier for its superior ability to handle the non-linear relationships inherent in environmental data and its robustness to overfitting. The model was trained on the refined feature set, with the objective of categorizing air quality into predefined AQI classes. Model training was conducted with careful attention to optimizing hyperparameters and ensuring that the model accurately reflected the underlying pollution dynamics.

● Model Validation

The performance of the Random Forest Classifier was rigorously evaluated using a hold-out validation set. Metrics such as accuracy, precision, recall, and the F1 score were calculated to assess the model's effectiveness in predicting air quality categories. The validation process confirmed the model's high predictive accuracy and its potential as a reliable tool for air pollution assessment.

K. Vijay Kumar / Afr.J.Bio.Sc. 6(Si2) (2024)

The experimental study demonstrated the framework's capability to leverage machine learning techniques for a nuanced analysis of air pollution intensity. The results underscored the value of integrating diverse datasets and employing advanced analytical methods to improve air quality monitoring and management strategies. Through this experimental validation, the framework was established as a robust and scalable solution for environmental scientists and policymakers aiming to mitigate the impacts of air pollution.

4.1. Results Discussion

The study focused on predicting AQI categories based on PM2.5, PM10, NO2 levels, temperature, wind speed, and urban density as features after applying the machine learning framework.

Table 1: Feature Importance Scores

Feature	Importance Score
PM2.5	0.35
PM10	0.25
NO2	0.2
Temperature	0.1
Wind Speed	0.05
Urban Density	0.05

The table 1 machine learning in environmental science, particularly in the quantification and scaling of air pollution intensity, has grown significantly. In our investigation, we used the Random Forest Classifier model, which was chosen due to its robustness in managing complex datasets common in environmental monitoring. The model's ability to predict air quality index (AQI) categories was critically dependent on a specific set of features, each of which contributed differently to the model's predictive power.

Table 1, which is central to our analysis, lists these features and their importance scores, providing quantitative information about their relative influence on the model's accuracy. PM2.5, PM10, and NO2 emerged as the most significant predictors. The prevalence of PM2.5 and PM10 particulate matter, which have diameters less than 2.5 and 10 micrometers, respectively, emphasizes the serious health risks posed by fine particles capable of penetrating deep into the respiratory tract and entering the bloodstream. NO2, a key indicator of vehicle and industrial combustion processes, emphasizes the link between human activity and air quality degradation. Temperature, wind speed, and urban density were identified as less influential in comparison, but their contributions were significant. Temperature influences the chemical reactions that produce or degrade air pollutants, whereas wind speed influences the dispersion and dilution of pollutants across areas. Urban density, which reflects the concentration of emissions from various sources, has a nuanced impact on local air quality dynamics.

The relative importance of these features reveals the complex interaction of natural and anthropogenic factors in determining air quality. For example, the variation in

importance scores reflects not only the direct impact of specific pollutants on AQI categories, but also the underlying environmental and socioeconomic factors that influence pollution levels. The importance of these features in predicting AQI categories is critical when developing targeted air quality management strategies. For example, the high importance scores for particulate matter indicate that policies aimed at reducing PM emissions could be especially effective in improving air quality. Similarly, strategies for reducing NO2 emissions from traffic and industry could provide significant benefits. Furthermore, the findings from the analysis of temperature, wind speed, and urban density highlight the importance of adaptive and dynamic pollution management strategies that take meteorological conditions and urban planning into account. This could include putting in place temporary measures during adverse weather conditions that are known to exacerbate pollution levels, as well as urban development policies that reduce the concentration of pollutants in densely populated areas. The detailed examination of feature importance not only validates the Random Forest Classifier's use in air quality prediction, but also provides a road map for prioritizing interventions. By identifying the factors that predict AQI categories, policymakers and environmental agencies can better devise and implement measures to address the multifaceted challenge of air pollution.




Figure 1: Model Accuracy Comparison

The bar graph shown in figure 1 compares the accuracy of the Random Forest Classifier with other models such as Support Vector Machine (SVM), Logistic Regression, and Decision Trees. The Random Forest model outperforms the others with a near-optimal accuracy of 92%, followed by SVM at 88%, Logistic Regression at 85%, and Decision Trees at 82%. This visualization underscores the effectiveness of the Random Forest approach in this context.

Table 2: Model Performance Metrics

Metric	Value
Accuracy	0.92
Precision	0.91
Recall	0.9
F1 Score	0.9

Table 2 explains the Random Forest Classifier's prowess in air quality prediction through a detailed presentation of performance metrics, including accuracy, precision,

K. Vijay Kumar / Afr.J.Bio.Sc. 6(Si2) (2024)

recall, and the F1 score. The high accuracy rate signifies the model's overall ability to correctly identify the AQI categories across a diverse set of observations, underscoring its reliability in environmental monitoring applications. Precision, a metric reflecting the model's ability to minimize false positives, coupled with recall, which measures the model's capacity to identify true positives, collectively highlight the model's efficacy in accurately classifying air quality conditions without significant trade-offs between identifying actual pollution events and avoiding false alarms. The F1 score, a harmonic mean of precision and recall, further affirms the model's balanced performance, indicating a robust capability to provide dependable predictions across varying air quality scenarios. This equilibrium between precision and recall is especially critical in the context of air pollution management, where the ability to accurately predict AQI categories enables timely and effective responses to mitigate adverse health and environmental impacts. The Random Forest Classifier's demonstrated performance metrics not only validate its application to the complex task of air quality prediction but also reinforce the potential of machine learning techniques in enhancing predictive accuracy and operational efficiency in environmental science.




Figure 2: Feature Importance Visualization

The feature importance plot shown in figure 2 visually represents the data from Table 1, with PM2.5, PM10, and NO2 shown as the top features influencing AQI category predictions. The figure 3 highlights the disproportionate weight of particulate matter and nitrogen dioxide levels in determining air quality, providing insights into pollution sources that most significantly impact public health.




Figure 3: AQI Prediction Accuracy Over Time

The line graph shown in figure 3 depicts the model's prediction accuracy over a 12-month period, illustrating consistency and stability in performance with slight variations due to seasonal changes in pollution levels and weather conditions. The model maintains an accuracy rate above 90% throughout the year, validating its effectiveness in diverse environmental conditions. The results demonstrate the Random Forest Classifier's efficacy in accurately predicting AQI categories using a selected set of environmental features. The high importance scores of particulate matter and nitrogen dioxide underscore the critical role these pollutants play in air quality. The model's superior performance, evidenced by its accuracy, precision, recall, and F1 score, establishes it as a reliable tool for air quality management and policy-making. These findings support the continued use and further development of machine learning frameworks for environmental monitoring and protection initiatives.

5. CONCLUSION

This experiment demonstrates that a machine learning-based analytical framework can quantify and scale air pollution intensity. We have demonstrated a reliable method for predicting air quality across a range of temporal and spatial scales through careful application of data preprocessing, feature engineering, and the use of advanced machine learning techniques, notably the Random Forest Classifier. High feature importance scores indicate that PM2.5, PM10, and NO2 pollutants have a significant impact on air quality. Random Forest Classifier was the most effective model. It outperformed other machine learning algorithms in terms of F1, accuracy, precision, and recall. This demonstrates that the model can deal with complex and nonlinear environmental data, making it suitable for air quality management and prediction. Visualizations of model accuracy comparisons, feature importance, and prediction accuracy over time show the framework's performance. The model's 12-month accuracy demonstrates its dependability and practical application. This work adds to what is already known about the application of machine learning in environmental science and opens up new research opportunities. To improve predictions, researchers may incorporate dynamic data sources such as real-time traffic flow or industrial activity data. The application of other cutting-edge machine learning and deep learning algorithms can improve the predictive accuracy and scalability of the model. To summarize, this article's analytical framework improves air pollution management and environmental monitoring. Machine learning can help us understand pollution. This will assist policymakers and environmental groups in making sound decisions and putting into action effective plans to reduce the health and environmental consequences of air pollution.

REFERENCES

- [1] Rudra Kumar, M., Gunjan, V.K. (2022). Machine Learning Based Solutions for Human Resource Systems Management. In: Kumar, A., Mozar, S. (eds) ICCCE 2021. Lecture Notes in Electrical Engineering, vol 828. Springer, Singapore. https://doi.org/10.1007/978-981-16-7985-8_129.

K. Vijay Kumar / Afr.J.Bio.Sc. 6(Si2) (2024)

- [2] Thulasi, M. S., B. Sowjanya, K. . Sreenivasulu, and M. R. . Kumar. "Knowledge Attitude and Practices of Dental Students and Dental Practitioners Towards Artificial Intelligence". *International Journal of Intelligent Systems and Applications in Engineering*, vol. 10, no. 1s, Oct. 2022, pp. 248-53.
- [3] Rudra Kumar, M., Gunjan, V.K. (2022). Peer Level Credit Rating: An Extended Plugin for Credit Scoring Framework. In: Kumar, A., Mozar, S. (eds) ICCCE 2021. *Lecture Notes in Electrical Engineering*, vol 828. Springer, Singapore. https://doi.org/10.1007/978-981-16-7985-8_128
- [4] Kalajdjieski, Jovan, Kire Trivodaliev, Georgina Mirceva, Slobodan Kalajdziski, and Sonja Gievska. "A complete air pollution monitoring and prediction framework." *IEEE Access* (2023).
- [5] Sofia, A. Sathya, K. Sowmiya, K. Soundarya, and M. Theepiga. "APD-ML: Air Pollution Detection Using Machine Learning Algorithms." In *2023 2nd International Conference on Vision Towards Emerging Trends in Communication and Networking Technologies (ViTECoN)*, pp. 1-5. IEEE, 2023.
- [6] Israfil, Mohd Afaque, Shubhangi Bhatnagar, Gaurang Juneja, Kamal Upreti, and Balram Rao. "Predictive Analysis of Air Pollution Using Machine Learning Techniques." *Asia-Pacific Journal of Management Research and Innovation* 18, no. 3-4 (2022): 152-162.
- [7] Mihirani, Madhushika, Lasith Yasakethu, and Sachintha Balasooriya. "Machine Learning-based Air Pollution Prediction Model." In *2023 IEEE IAS Global Conference on Emerging Technologies (GlobConET)*, pp. 1-6. IEEE, 2023.
- [8] Feng, Xiaoyang. "Air pollution prediction in context of supervised machine learning and time series model." In *Second International Conference on Statistics, Applied Mathematics, and Computing Science (CSAMCS 2022)*, vol. 12597, pp. 904-912. SPIE, 2023.
- [9] Kumar, K., and B. P. Pande. "Air pollution prediction with machine learning: a case study of Indian cities." *International Journal of Environmental Science and Technology* 20, no. 5 (2023): 5333-5348.
- [10] Soundari, A. Gnana, J. Gnana Jeslin, and A. C. Akshaya. "Indian air quality prediction and analysis using machine learning." *Int J Appl Eng Res* 14, no. 11 (2019): 181-186.
- [11] Srivastava, Harshit, Goutam Kumar Sahoo, Santos Kumar Das, and Poonam Singh. "Performance Analysis of Machine Learning Models for Air Pollution Prediction." In *2022 International Conference on Smart Generation Computing, Communication and Networking (SMART GENCON)*, pp. 1-6. IEEE, 2022.
- [12] Du, Shengdong, Tianrui Li, Yan Yang, and Shi-Jinn Horng. "Deep air quality forecasting using hybrid deep learning framework." *IEEE Transactions on Knowledge and Data Engineering* 33, no. 6 (2019): 2412-2424.



CVR COLLEGE OF ENGINEERING

(An UGC Autonomous Institution, NAAC A Grade)

(Approved by AICTE & Govt. of Telangana and Affiliated to JNTU, Hyderabad)
Vastunagar, Mangalpalli (V), Ibrahimpatnam (M), R.R (D), Telangana - 501 510



Certificate of Participation

This is to certify that **Mr. Khambhampati Vijay Kumar**, Assistant Professor, **ANURAG ENGINEERING COLLEGE, ANANTHAGIRI** has Participated in one week Faculty Development Program(FDP) on **Advances in GenAI and Agentic AI** from **16th to 21st June, 2025** organized by the **Department of Computer Science and Engineering** in collaboration with **Computer Society of India, CSI-SB, CVR College of Engineering, Hyderabad.**

Dr. C. Ramesh
Professor, CSE
Coordinator

Dr. M. Sridevi
Associate Professor, CSE
CSI-SBC

Dr. A Vani Vathsala
Professor and Head of CSE
Convener

Dr. K. Ramamohan Reddy
Principal