

Experimental anti-tumor therapy in 3-D: Spheroids – old hat or new challenge?

JUERGEN FRIEDRICH¹, REINHARD EBNER², & LEONI A. KUNZ-SCHUGHART³

¹Institute of Pathology, University of Regensburg, Germany, ²Avalon Pharmaceuticals Inc., Germantown, Maryland, USA, and ³OncoRay, Center for Radiation Research in Oncology, Faculty of Medicine Carl Gustav Carus, Dresden, Germany

(Received 16 April 2007; revised 11 September 2007; accepted 4 October 2007)

Abstract

Purpose: To give a state-of-the-art overview on the promise of three-dimensional (3-D) culture systems for anticancer drug development, with particular emphasis on multicellular tumor spheroids (MCTS).

Results and conclusions: Cell-based assays have become an integral component in many stages of routine anti-tumor drug testing. However, they are almost always based on homogenous monolayer or suspension cultures and thus represent a rather artificial cellular environment. 3-D cultures – such as the well established spheroid culture system – better reflect the *in vivo* behavior of cells in tumor tissues and are increasingly recognized as valuable advanced tools for evaluating the efficacy of therapeutic intervention. The present article summarizes past and current applications and particularly discusses technological challenges, required improvements and recent progress with the use of the spheroid model in experimental therapeutics, as a basis for sophisticated drug/therapy screening. A brief overview is given focusing on the nomenclature of spherical 3-D cultures, their potential to mimic many aspects of the pathophysiological situation in tumors, and currently available protocols for culturing and analysis. A list of spheroid-forming epithelial cancer cell lines of different origin is provided and the recent trend to use spheroids for testing combination treatment strategies is highlighted. Finally, various spheroid co-culture approaches are presented that have been established to study heterologous cell interactions in solid tumors and thereby are able to reflect the cellular tumor environment with increasing accuracy. The intriguing observation that in order to retain certain tumor initiating cell properties, some primary tumor cell populations must be maintained exclusively in 3-D culture is mentioned, adding a new but fascinating challenge for future therapeutic campaigns.

Keywords: 3-D culture, multicellular tumor spheroids, anti-tumor therapy

Introduction

Establishment and adaptation of cell-based assays which rapidly detect the most active drug candidates from large pools of potential effectors has become an integral component in many stages of anti-tumor drug development and testing. However, most cell-based assays apply monolayer or suspension cultures which present a largely artificial cellular environment with limited relevance for *in vivo* studies, and often with little value in predicting clinical efficacy of both generally cytotoxic as well as molecularly targeted drugs. Three-dimensional (3-D) cultures are gaining increased recognition as important additions to the strategic toolkit for evaluating the efficacy of therapeutic intervention. In cancer research and treatment in particular, they are becoming more and more accepted as essential research tools

which ideally complement the techniques already available.

Examples of 3-D cell models employed in basic and applied tumor biology include multilayer cell systems cultured on top of porous membranes, used for example to study drug transport and binding, therapy resistance and cell invasion (Hicks et al. 1997, Padron et al. 2000). Matrix-embedded cultures of single cells or cell aggregates using various artificial and natural extracellular matrix compounds have proven helpful in investigating tumor cell-matrix interactions, tumor cell differentiation, migration and invasion processes (Casciari et al. 1994, Muralidhar et al. 2001, Debnath & Brugge 2005, Lee et al. 2007). Hollowfiber bioreactors with cells cultured within a network of perfused artificial capillaries to analyze cell metabolism and therapy resistance (Gillies et al. 1993, Legallais et al. 2001,

Dulong et al. 2002) and more complex systems such as *ex vivo* cultures of pieces or slices from different tissues or tumors excised and cultured *in vitro* (Berglund et al. 2002, Ochs et al. 2003) have also been employed. All of these 3-D *in vitro* culture systems are of intermediate complexity, mimicking different aspects of the human tumor tissue environment, and should thus be considered as advanced models for routine anti-tumor drug testing. One of the culture systems that attracted major attention throughout the past 10 years is the model of Bissell and co-workers, who established a laminin-rich ECM 'embedded' assay, in which breast epithelial cells form either polarized, growth-arrested acinus-like or disorganized, proliferative and non-polar colonies, depending on their non-malignant or malignant status. This model system, its procedures and alternative protocols have previously been reviewed in detail (Gudjonsson et al. 2003, Nelson & Bissell 2005, Lee et al. 2007). There is also increasing effort to engineer advanced culture devices to improve and miniaturize such systems for screening purposes. Here, microfluidic devices are one of the most promising approaches (e.g., Bhadriraju & Chen 2002, Walker & Beebe 2002, Chaw et al. 2007, Kim et al. 2007, Warrick et al. 2007). For example, a recently developed tubeless microfluidics systems that can be operated by standard liquid handling automation has been successfully combined with the IrECM breast tissue architecture model (e.g., Hayes, Skoien et al, 13th Annual Conference of the Society for Biomolecular Sciences, 2007, Poster/Abstract PSTZ1H002). Fully recapitulating acinar morphogenesis, this combined platform has obvious potential as an attractive model for high-throughput screening in the near future.

The present article, however, emphasizes the potential and recent progress with the 'classical' 3-D cell system in cancer research: the multicellular spheroid model (MCS). Multicellular tumor spheroids (MCTS) reflect some pathophysiological phenomena in tumor tissues that critically affect therapeutic efficacy. They have frequently been applied to anti-tumor therapy testing throughout the past four decades, e.g., to study and/or identify mechanisms of radio- or drug resistance, as reviewed in a number of comprehensive articles and article series with different focus. But most importantly, the spheroid model is suitable to be incorporated readily into mainstream and routine drug testing operations, as suggested earlier (Mueller-Klieser 2000, Kunz-Schughart et al. 2004). In this article, the power of this *in vitro* model system for sophisticated drug testing is explained, and technical basics and recent advances with the spheroid model are briefly discussed. Spheroid-based approaches to drug screening under conditions that reflect tumor tissue

conditions with increasing accuracy are highlighted in the following chapters.

Aggregates, spheres, spheroids, organoids, tumoroids: Spherical 3-D cultures reflecting tumor pathophysiology

Numerous review and book articles have summarized the history of the spheroid model, starting with its introduction in cancer research by radiobiologists in the early 1970s (Sutherland et al. 1970, 1971a, 1971b). Since then, the classical system of Sutherland and co-workers of spherical multicellular cultures of animal cell lines has been extended, e.g., by the use of a variety of human cell lines and has also been modified in other ways as detailed later. Unfortunately, much of the corresponding literature is not easily identified via standard reference searching because of varying nomenclature used by different authors. Tumoroids (Vamvakidou et al. 2007), for example, are no more or less than mixed spheroid cultures (Djordjevic & Lange 1990, Lange et al. 1992, Djordjevic et al. 1993, Kunz-Schughart 1999, Djordjevic & Lange 2004, 2006) or so-called mosaic spheroids (Boyd et al. 2002, 2004) consisting of two or more types of cells. Other synonyms describing 3-D cultures with spherical structure include aggregate, sphere or organoid. Aggregates may be more or less spherical but in most cases, the term is exclusively used to describe a loose package of cells and not the formation of compact spherical cultures (Ivascu & Kubbies 2006). On the other hand, some so-called spheroids in the literature are no more than spherical loose aggregates, may easily fall apart if advanced analytical processing is required, and may not express all pathophysiological characteristics known to be present in compact multicellular tumor spheroids. And last but not least, the term sphere is frequently used in biochemical papers, sometimes for real spheroids and sometimes for crude aggregates as well as in the field of stem/progenitor cell research which also includes the search for and study of so-called tumor stem cell populations (Ricci-Vitiani et al. 2007).

These inconsistencies have to be kept in mind when setting up spheroid-based therapeutic test systems. Here, the three-dimensional structure is mandatory to achieve the system's advantage of mimicking avascular tumor nodules or microregions of solid tumors with respect to growth kinetics, micromilieu and restoration of *in vivo* morphological and functional features of tumor cells as depicted in the cartoon of Figure 1. Cells located in the spheroid periphery reflect the situation of actively cycling tumor cells adjacent to capillaries *in vivo*. In contrast, innermost cells become quiescent and eventually die via apoptosis or necrosis to form a secondary necrotic core. The concentric arrangement of




Figure 1. Multicellular tumor spheroids (upper panel left; d = diameter) display growth characteristics and numerous pathophysiological features of avascular tumor nodules and/or developing micrometastases (lower panel) and also inside out resemble the situation from the capillary towards the periphery in tumor microregions (upper panel right) (modified from Kunz-Schughart 1999).

heterogeneous cell populations in spheroids and the establishment of pathophysiological gradients is comparable to the situation in micrometastases, avascular tumor microregions or inter-capillary tumor regions as detailed earlier (Mueller-Klieser 1987, Sutherland 1988). Beyond a critical size of $500 - 600 \mu\text{m}$, central secondary necrosis develops in most but not all spheroid types from permanent cell lines. Typically surrounded by a viable rim of cells of about $100 - 300 \mu\text{m}$, these necroses mainly relate to

the limited inward and outward diffusion of positive and negative regulators which cause a lack of oxygen and/or nutrients, accumulation of waste products and low pH (Acker et al. 1987, Carlsson & Acker 1988). Studies on steady state-levels of glucose, lactate, and energy-rich phosphates as well as mitochondrial activity in spheroids indicate an adaptation of metabolism of inner spheroid regions to environmental stress, maintaining intracellular homeostasis until shortly before cell death

(Freyer et al. 1990, 1991, Walenta et al. 1990, Bredel-Geissler et al. 1992, Teutsch et al. 1995). However, it has also become evident that tumor cells can critically differ in their sensitivity to this pathophysiological stress situation, as should be expected from their highly variable capacity for energy production under both aerobic and anaerobic conditions. Multicellular tumor spheroids (MCTS) may also contain an extensive extracellular matrix (ECM) and possess a complex three-dimensional network of cell-to-matrix and cell-to-cell interactions. This not only affects the distribution and function of biological response modifiers such as hormones and growth factors but is also relevant to the penetration and action of drugs.

Technology in brief

Two major aspects in spheroid technology are relevant for integration of this model system in routine drug testing: Easy and rapid handling of both culturing and analysis. Various techniques for spheroid culturing have been described earlier and are used in routine animal cell culturing (Mueller-Klieser 1987, Lund-Johansen et al. 1992, Kunz-Schughart et al. 2000). Figure 2 details the most frequently applied methods both for the generation of large amounts of spheroids or spheres and for single spheroid monitoring. In fact, culture technology has not much changed over the past twenty years, with spinner flask cultures remaining one of the most attractive and robust approaches to obtain a large pool of spheroids of particular size-ranges under optimum, stirred nutrient supply conditions. Here, spherical aggregates of tumor cells form spontaneously in the absence of attachment to another substrate directly in the flask. Alternatively, spheroids can be obtained during an initiation interval in non-adherent, e.g. bacteriological or agar-coated, so called 'mother dishes' from which they are selectively transferred into the flasks. Comparable approaches to generate large amounts of spheroids and to avoid effects caused by stagnant medium are roller tubes or roller bottles and the gyratory shaker. An interesting newly developed technology with similar potential is the rotating wall vessel ('NASA bioreactor') that provides even more constant culture conditions due to continuous medium perfusion and nutrient flow rate. This rotary cell culture system was designed to mimic micro-gravity and maintain cells in suspension under very low shear stress (Ingram et al. 1997) and has been employed to also grow tumor spheroids of different complexity (e.g., Clejan et al. 2001, Song et al. 2004, Vamvakidou et al. 2007). However, while being recommended for some normal and primary cell culturing under optimized culture conditions, this

technology does not provide major advantages for drug testing over the spinner flask technique due to the more cumbersome system set-up. Furthermore, the system is significantly more expensive than a spinner flask, while any advantages other than reduced shear force have yet to be demonstrated.

One has to keep in mind that all rotating culture systems require relatively high quantities of media, conflicting with the usual demand for minimum amounts of new drug candidates to be applied in a drug test system. Therefore, it may indeed be required to use stationary spheroid culture systems at least for drug exposure. Stationary culturing technologies include the growth of spheroid cultures in 96-well plates which allows monitoring and manipulation of single spheroids, since the inoculation of aggregating tumor cells into agar-coated 96-well plates results in the formation of one single spheroid per well. The well-to-well variation in spheroid diameter after a defined initiation period is less than 5% for most spheroid types and remains extraordinarily small throughout spheroid growth if 50% of the supernatant is renewed at least every other day. More frequent medium renewals can improve the supply conditions. High reproducibility in spheroid formation, the potential for analyzing individual spheroids before and after treatment, and the limited amount of drug required qualifies the 96-well plate approach as a practical therapy test system. However, one also has to realize that some of the limitations that are inevitable in high-throughput test settings may of course complicate interpretation of some drug study data. For example, early investigations have shown that the diffusion-limited supply under stationary conditions results in significant external nutrient gradients, and these change as a function of time as the spheroid grows. Also, supply of drugs to the spheroid may be more diffusion-limited than in rotary culture, depending on the concentration and cellular uptake rate. A further consideration specifically relevant for drug testing is that drug effects themselves (e.g., cytotoxicity) can alter the metabolic rates of treated spheroids versus controls, which may result in different external (and internal) nutrient environments.

In numerous previous studies, flat bottom wells coated with 1.5–3% agar/agarose in serum-free medium to form a non-adherent, concave surface have proven to be appropriate for liquid overlay spheroid culturing (Carlsson & Yuhás 1984), but modification such as the application of round bottom wells coated with 0.5% poly-HEMA (hydroxyethyl methacrylate) in 95% ethanol are also used (Tong et al. 1992, Hoevel et al. 2004, Ivascu & Kubbies 2006). Unfortunately, individual coating of plates is still required today, since current commercially available non-adherent 96-well plates do not




Figure 2. Techniques frequently applied to grow multicellular spheroids from cell lines or primary cells, spheroid co-cultures and fragment spheroids from tissue materials.

guarantee a reproducible formation of single spheroids per well. This drawback may also have resulted in the recent use of the hanging drop method for culturing tumor spheroids (Kelm et al. 2003, Del Duca et al. 2004), a methodology that was primarily established for culturing embryoid bodies. The technique, while avoiding the coating of plates for spheroid formation, is however only useful for short-term culturing, i.e., spheroid initiation, for as long as spheroids are not fed and/or treated.

A broad range of human tumor cell lines has been studied in spherical culture using the various culturing methods discussed. These include epithelial cell lines of different tumor entities but also brain tumors such as gliomas and glioblastomas, melanomas and sarcomas. We have been particularly interested in epithelial tumors and thus provide an overview of human carcinoma cell lines that have been shown to form spheroids in previous studies and that may be applicable for therapeutic intervention modeling (Table I). The list also contains information as to whether or not the respective cell line is included in the 60-cell line screen of the US National Cancer Institute Developmental Therapeutics Program (NCI-DTP; <http://dtp.nci.nih.gov/branches/btb/ivclsp.html>) which was implemented about 20 years ago. The NCI60 cell line-screen has been found to be of some value as predictor of xenograft and/or clinical drug activity (Newell 2001, Johnson et al. 2001, Monga & Sausville 2002) and has become a gold standard in drug screening (Newell 2001, Johnson et al. 2001). In the meantime, many drugs and drug candidates have been tested in these cell lines and a comprehensive database is freely available, e.g., providing parameters such as IC_{50} values in monolayer cultures. Nevertheless, tremendous controversy remains in this area with several reports stressing the limited predictive value of routine *in vitro* and *in vivo* drug screening models for clinical efficacy (Sausville 2000, Pelkonen et al. 2001, Balis 2002, Kerbel 2003, Voskoglou-Nomikos et al. 2003). Because 3-D culture systems such as spheroids better reflect the pathophysiology of human tumors, this debate lends further arguments in favor of the application of 3-D culture systems for evaluating therapeutic efficacy. Because a screening of the referenced literature shows that not all spherical cultures are compact spheroids that can easily be applied for drug testing as discussed earlier, we want to highlight a recent article from Ivascu and Kubbies (2006), who described the rapid generation of single tumor spheroids for high throughput cell function and toxicity analysis by the addition of extracellular matrix compounds during spheroid initiation. In their study, 2.5% reconstituted basement membrane was superior to fibronectin, laminin, collagen type

IV, collagen type I, heparan sulphate proteoglycane, chondroitin sulphate, and diverse mixtures. Despite the fact that the addition of ECM per se may alter the expression profile and function of tumor cells, potentially affecting the sensitivity to therapeutic intervention, this is one of the few options to grow non-adherent tumor cells in spheroid culture. As an alternative, the application of methylcellulose according to the endothelial cell spheroid model introduced by Korff and Augustin (1998) may be considered.

MC(T)S and experimental therapeutics

Despite tumor cell type dependent variations in spheroid growth and histomorphology, several physiological properties determined by the spherical geometry within MCTS as described above are general in nature. Therefore, MCTS have become a powerful tool to study therapeutic problems associated with metabolic and proliferative gradients, in particular the responsiveness and effects of chronically hypoxic tumor cells and 3-D cell-cell and cell-matrix interactions, e.g., in radio- and chemo-resistance. Needless to say, studies related to radiotherapy dominated the field of spheroid work in experimental therapy for its first 20 years, mainly as a consequence of their earlier implementation in radiobiological cancer research. This work has been reviewed in diverse dated but still informative articles which also highlight and discuss the phenomenon of hypoxia and cell-cell contact related radioresistance (Carlsson & Nederman 1989, Freyer 1992, Buchsbaum et al. 1993, Olive & Durand 1994, Mueller-Klieser 1997, Santini & Rainaldi 1999). In the 1990s, the number of scientific reports on this topic decreased, although throughout several groups remained active in this area. During the same time, the spheroid technology increasingly spread into other fields of therapeutic research ranging from chemotherapy, radioimmunotherapy, cell- and antibody-based immunotherapy, hyperthermia, gene therapy and photodynamic treatment (Olive & Durand 1994, Mueller-Klieser 1997, Desoize et al. 1998, Mueller-Klieser 2000, Desoize & Jardillier 2000). The aspect of drug resistance was examined in particular detail by Olive and coworkers who have frequently used the comet assay to identify DNA-damaged cells (Durand & Olive 1992, Olive & Durand 1994).

Today, a number of classical therapeutics have been tested in spheroid culture, but the various spheroid types applied for drug exposure as well as the different analytical tools and endpoints may limit the potential for comparing individual, independent studies. Recently, Mellor et al. (2005) compared the effect of 5-fluorouracil (5-FU), vinblastine,

Table I. Overview of some epithelial tumor cell lines of different origin that form spherical cultures with representative reference list.

Entity	Cell line	NIH-DTP screen	Literature
Squamous cell carcinoma	1483	—	(Sturgis et al. 1994)
	A431	—	(Ng et al. 1987; Casciari et al. 1988; Kwok & Sutherland 1991)
	CaSki	—	(Casciari et al. 1988; Kwok & Sutherland 1991; Lohr et al. 1995)
	FaDu	—	(Stuschke et al. 1992)
	HEp-2	—	(Kohno et al. 1988)
	HN-1	—	(Schwachofer et al. 1989)
	MDA 686LN	—	(Sturgis et al. 1994)
	MDA 886Ln	—	(Sacks et al. 1989; Nishikawa et al. 1990; Sturgis et al. 1994)
Lung/NSCLC	SQ5	—	(Nakayama 1991)
	A549*	+	(Ivascu & Kubbies 2006)
	NCI-H125	—	(Schwachofer et al. 1991)
	NCI-H322M	+	(Ivascu & Kubbies 2006)
	NCI-H40*	+	(Ivascu & Kubbies 2006)
SCLC	SK-MES-1	—	(Audran et al. 1994)
	DMS53	—	(Cate et al. 1984; Douple et al. 1985)
Colon	COLO 205*	+	(Ivascu & Kubbies 2006)
	DLD-1	+	(Nicholson et al. 1997; Mellor et al. 2005)
	HCT-15*	+	(Ivascu & Kubbies 2006)
	HCT-116	+	Information by word of mouth
	HT29	+	(Lees et al. 1981; Sutton et al. 1982; Sutherland et al. 1986; Simiantonaki et al. 2002; Lowthers et al. 2003)
	LoVo	—	(Soranzo et al. 1986; Soranzo & Ingrosso 1988; Pervez et al. 1989; Dolfini et al. 2005)
	LS174T	—	(Horan et al. 1985; Langmuir et al. 1989)
	SW620*	+	(Ivascu & Kubbies 2006)
Pancreas	WiDr [§]	—	(Chan et al. 1991; Banks et al. 1994; Jansen et al. 1994; Buffa et al. 2001; Mueller-Klieser et al. 2002)
	MIAPaCa2	—	(McLeod et al. 1997)
	Panc1	—	(McLeod et al. 1997)
Liver/hepatocellular carcinoma	Hep3B	—	(Lin et al. 2006)
	HepG2*	—	(Gorlach et al. 1994; Selden et al. 2000; Kelm et al. 2003; Ivascu & Kubbies 2006; Lambert et al. 2006; Lin et al. 2006)
	Huh-7	—	(Tonary & Pezacki 2006)
Breast	BT20	—	(Muller-Holzner et al. 1996)
	BT474	—	(Kunz-Schughart et al. 2001; Seidl et al. 2002)
	MCF-7	+	(Pourreau-Schneider et al. 1984; Olea et al. 1992; Kunz-Schughart et al. 2001; Kelm et al. 2003; Ivascu & Kubbies 2006)
	MDA-MB-231	+	(Ivascu & Kubbies 2006; Muir et al. 2006)
	MDA-MB-361	—	(Hoevel et al. 2004; Ivascu & Kubbies 2006)
	MDA-MB-468	+	(Ivascu & Kubbies 2006)
	SK-BR-3	—	(Kunz-Schughart et al. 2001; Ivascu & Kubbies 2006)
	T47D	+	(Ronen et al. 1990; Kunz-Schughart et al. 2001; Kan et al. 2001; Seidl et al. 2002; Ivascu & Kubbies 2006)
	MDA-MB-435 [†]	+	(Stuschke et al. 1995; Ivascu & Kubbies 2006)
	NCI/ADR ^{RES‡}	+	(Walker et al. 2004; Ivascu & Kubbies 2006)
Ovary	IGR-OV1*	+	(Ivascu & Kubbies 2006)*
	MCAS	—	(Sonoda et al. 2003)
	OVCAR-3	+	(Bardies et al. 1992; Filippovich et al. 1996)
	OVCAR-5	+	(Casey et al. 2001)
	SKOV3*	+	(Makhija et al. 1999; Frankel et al. 2000; Casey et al. 2001; Ivascu & Kubbies 2006)*
Cervix	HeLa-S3	—	(Sommers & Alfieri 1998)
	SiHa	—	(Olive et al. 1996; Olive & Banath 1997; Durand & Sham 1998; Oloumi et al. 2000; Oloumi et al. 2002)
Kidney	786-0	+	(Lieubeau-Teillet et al. 1998)
	A498	+	(Heuser et al. 2003)
	(HEK)-293	—	(Talts et al. 2000; Ivascu & Kubbies 2006)
	UO-31	+	(Ivascu & Kubbies 2006)

(continued)

Table I. (Continued).

Entity	Cell line	NIH-DTP screen	Literature
Bladder	J82	—	(Lottner et al. 2004)
	MGH-U1	—	(Tannock & Kopelyan 1986; Kerr & Smart 1991; Erlichman & Wu 1992)
	MGH-U3	—	(LaRue et al. 1997; Kilani et al. 2003)
	RT-4	—	(Konur et al. 1998)
	RT-112	—	(Huygens et al. 2003; Kamuhabwa et al. 2003)
	T24	—	(Kawai et al. 2001)
	UROtsa	—	(Lottner et al. 2004)
Prostate	DU-145	+	(Essand et al. 1993; Frankel et al. 2000; Enmon, Jr. et al. 2002; Wartenberg et al. 2002; Wartenberg et al. 2003)
	LNCAP*	—	(Ballangrud et al. 1999; Ballangrud et al. 2001; Konduri et al. 2001; Enmon et al. 2003; Ivascu & Kubbies 2006)*
	PC-3*	+	(Donaldson et al. 1990; Hedlund et al. 1999; Enmon, Jr. et al. 2002; Ivascu & Kubbies 2006)*
Skin/melanoma	A375	—	(Cheng et al. 1991; Mills et al. 2002; Martiny-Baron et al. 2004)
	BEX-c	—	(Hystad & Rofstad 1994; Rofstad et al. 1996)
	BRO	—	(Schwachofer et al. 1990; Schwachofer et al. 1991)
	C32	—	(Yuhass et al. 1984)
	C918	—	(Braun et al. 2007)
	C8161	—	(Strasberg et al. 2004)
	CaCL 73–36	—	(McFadden & Kwok 1988; Kwok et al. 1988; Kwok et al. 1989)
	DC-10	—	(Feder-Mengus et al. 2007)
	HBL	—	(Feder-Mengus et al. 2007)
	HMV-I	—	(Kubota et al. 1984; Kuwashima et al. 1988; Kuwashima et al. 1993)
	Hs294	—	(Offner et al. 1996)
	IGRI	—	(Gorlach et al. 1994; Porwol et al. 1996)
	Mel108	—	(Gottfried et al. 2006)
	MelIm	—	(Gottfried et al. 2006)
	MeWo	—	(Offner et al. 1996; Jorgensen et al. 2005)
	NA8	—	(Ghosh et al. 2005; Feder-Mengus et al. 2007)
	OM467	—	(Parry et al. 1992)
	OM482	—	(Parry et al. 1992)
	SB-2	—	(Mills et al. 2002)
	SK-MEL-28	+	(Freyer 1988; Freyer & Schor 1989; Gorlach et al. 1994; Offner et al. 1996; LaRue et al. 2004)
	ST-ML-11	—	(Offner et al. 1996)
	ST-ML-12	—	(Offner et al. 1992; Offner et al. 1993; Offner et al. 1996; Simiantonaki et al. 2002)
	ST-ML-14	—	(Offner et al. 1996)
	ST-ML-25	—	(Offner et al. 1996)
	WIX-c	—	(Hystad & Rofstad 1994; Rofstad et al. 1996)
	WM9	—	(Francia et al. 2005)
	WM35	—	(Shellman et al. 2003; Francia et al. 2005; Jorgensen et al. 2005)
	WM45.1	—	(Jorgensen et al. 2005)
	WM115	—	(Fang et al. 2005; Monzani et al. 2007)
	WM164	—	(Rieber & Rieber 2006)
	WM239	—	(Francia et al. 2005; Jorgensen et al. 2005)
	WM902B	—	(Jorgensen et al. 2005)
	WM1341B	—	(Jorgensen et al. 2005)
	WM3517	—	(Fang et al. 2005)
	WM3523	—	(Fang et al. 2005)
	WM3539	—	(Fang et al. 2005)

*with reconstituted basement membrane rBM in Ivascu & Kubbies 2006.

§WiDr cells are reported to be a derivative of HT29 (Chen et al. 1987).

†MDA-MB-435 breast origin is questioned as cells express melanocyte-specific genes (Ellison et al. 2002; Rae et al. 2004).

‡NCI/ADR^{Res} cells were recently identified to be derived from the ovarian cell line OVCAR-8 (Garraway et al. 2005).

doxorubicin, and cisplatin in a proliferating and quiescent spheroid culture model of DLD-1 colon adenocarcinoma cells. Their analyses of drug efficacy included spheroid volume growth and cell survival as

determined via a methylenblue staining procedure. A critical difference was documented not only between the various drugs but more importantly for the same drug in spheroids with distinct proliferation status

(Mellor et al. 2005). A similar study using HT-29 and HCT116 colon cancer cells has been performed by our group to demonstrate differences in drug efficacy and half-maximal inhibitory concentration (IC_{50}) values, respectively, of 5-FU, irinotecan and the new drug SymdaxTM in monolayer and spheroid culture with incubation intervals of 24 and 72 h (Friedrich et al., 2002). Orlandi et al. compared the effect of idarubicin and idarubicinol in MCF-7 breast cancer cell monolayers and MCTS, demonstrating significant activity in both culture systems with IC_{50} values (24 h exposure) that were only slightly higher in spheroids (Orlandi et al. 2005). In this study, evidence for drug resistance in spheroids seemed critically dependent on the time of exposure. It may be speculated that this finding is not due to cellular drug resistance but to an insufficient or delayed distribution of the drugs within spheroids which of course leads to a reduced efficacy. An attenuated resistance to doxorubicin in three-dimensional aggregates of the human breast carcinoma cells MDA-MB-231 was observed in a study by Muir et al. (2006) by administration of the nitric oxide mimetics diethylenetriamine/nitric oxide adduct (DETA/NO) and glyceryl trinitrate (GTN). Their data imply that the chemosensitization following NO mimetic exposure is not due to increased penetration of the drug nor associated with an increase in proliferative activity and suggest the involvement of a cyclic guanosine monophosphate (cGMP)-dependent signaling (Muir et al. 2006). It should, however, be noted that in other studies MDA-MB-231 have been described to only form loose aggregates and not compact spheroids (Ivascu & Kubbies 2006). Limitations in drug penetration are thus unlikely to occur, i.e., an effect on the penetration kinetics of a drug may not be visible. However, the study clearly reflects the present trend to use the spheroid model for testing combinatorial therapeutic strategies.

One field of interest is of course the combination of various drugs and/or radiosensitizers with single and fractionated irradiation regimens. Just briefly, supra-additive effects of cisplatin, gemcitabine or 5-Fluorouracil with irradiation were recently recorded in HepG2 hepatocellular carcinoma spheroids (Lambert et al. 2004). The same combinations were studied in glioma and glioblastoma cell line (GaMg, U87, Gli-6) and/or fragment spheroids from primary tissues, and the effect was compared with that observed in monolayer cultures (Genc et al. 2004, Fehlauer et al. 2005, Fehlauer et al. 2006). In the U87 glioma spheroid model the selective mTOR (mammalian target of rapamycin) inhibitor rapamycin and the cyclin-dependent kinase inhibitor roscovitine were shown to critically enhance the efficacy of fractionated irradiation. Interestingly, the additive effect of rapamycin was not present in two-dimen-

sional (2-D) cultures (Eshleman et al. 2002). Radio-sensitization and additive or supra-additive effects were also achieved with the xanthin derivate pentoxifylline, studied in T98G glioma spheroids. Here, the largest impact on radiation response occurred if intermittent doses were applied (Eley et al. 2002). The Heat-shock protein Hsp-90 inhibitor 17-N-Allyl-amino-17-Demethoxy-Geldanamycin (17-AAG) was examined in LNCaP and CWR22Rv1 prostate carcinoma spheroids and showed radiosensitizing potential (Enmon et al. 2003). As a last example, the study of Khaitan et al. (2006) shall be mentioned, which compared radiomodification by 2-Deoxyglucose (2-DG) in monolayer cultures and spheroids of a human glioma cell line (BMG-1). 2-DG – also the base substance for the most frequently applied clinical radiotracer for positron emission tomography PET ($[^{18}F]$ FDG) – enhanced the radiation-induced cell death of spheroids more drastically than that of monolayer cells. The study demonstrated that 2-DG uptake is enhanced in spheroid as opposed to monolayer cells of the BMG-1 glioma cell line. Accordingly, the radiosensitization effect leading, via different mechanisms, to transient G2 delay, apoptosis induction and late onset of DNA breaks is more pronounced in the 3-D setting. These few examples just illustrate the range of application of spheroids in the study of combination therapy and shall encourage scientists at academic institutions and in the pharmaceutical industry to use the model for future-oriented treatment strategies.

MC(T)S in drug screening: Progress and challenges

As detailed earlier, many complex methodologies have been applied to analyze intact and viable, frozen or paraffin-embedded and also dissociated spheroid cultures (Kunz-Schughart et al. 1998). Today, all modern molecular techniques can be exploited to study spheroid cells on the cellular, protein, RNA and DNA level. However, with respect to drug and therapy screening, there is still a lack of automated and easy-handling analytical tools and a need for defined, standard practice endpoints. Spheroid volume growth, which can be mathematically described in many spheroid types by a Gompertz equation (Marusic et al. 1994, Kunz-Schughart et al. 1996) or other mathematical models (Tabatabai 2005), and growth delay following treatment are just one set of parameters that should routinely be determined. This of course requires long-term measurement of spheroid diameters. Nevertheless, monitoring of these basic volume growth parameters is still the most popular technique to evaluate effects of growth-promoting or growth-suppressing factors and drugs in MCTS, is technically easy, and

provides a highly convenient endpoint for quantitative studies (e.g., Figure 3a). Simple readings of the growth or shrinkage of spheroids can be accomplished by standard phase-contrast microscopy and computer image analysis.

An interesting technique for analyzing and sorting intact spheroids is flow cytometry, first demonstrated by Freyer and colleagues (1987). To date, this technique has seen limited application (Freyer et al. 1987, 1989), primarily due to the requirement for instrument modification to handle large particles on most commercial flow cytometers. However, a recently launched commercial instrument (Union Biometrica COPASBiosorter, <http://www.unionbio.com/>) that is capable of both light scattering and fluorescence analysis and sorting of large particles up to 1500 μm may encourage broader use of this technology for spheroid sizing, analysis and selection. It is recognized that high throughput flow cytometric analysis may require relatively large numbers of structurally intact spheroids and may thus not necessarily be suitable for every treatment modality. However, the option of selecting intact spheroids of defined sizes remains a clear challenge. As an example, such an approach could foster increased use of MCTS pre-cultured under optimal rotary culture and nutrient supply conditions, e.g., prior to selective transfer into sufficient culture (plate) systems for testing small amounts of new drug candidates.

Since spheroid growth delay and re-growth per se do not allow an independent separation of effects on viability versus effects on growth rate or recruitment rate of quiescent cells to re-enter the cell cycle, other analytical endpoints should be considered. Parameters of interest with potential for automated analysis include cell viability and survival, respectively, outgrowth or migratory activity of spheroid cells, and the (patho)physiological status within spheroids. A multicellular spheroid-based sensor based on real-time impedimetric measurement was presented a few years ago for testing anti-cancer therapeutics. But the proposed biohybrid system, with spheroids to be positioned in a measurement capillary for monitoring impedance spectra as a measure for spheroid morphology, cell integrity and physiology has not yet entered routine analysis, probably due to the highly delicate handling and data interpretation required (Thielecke et al. 2001). The system has been used recently to study genetically altered and methotrexate treated MCF-7 spheroids (Bartholoma et al. 2005). Its applicability in drug screening, however, has yet to be validated. More importantly, there have been a number of attempts to simplify the measurement of cell number and viability in MCTS. Respiratory activity is generally regarded to reflect cellular status and to




Figure 3. The effect of 5-FU treatment (72 h treatment interval) as determined by spheroid volume growth kinetics (a) and directly after treatment using the recently validated APH cell viability assay (b). The APH assay is applicable in spheroids grown under serum and serum-free conditions. The effect of 5-FU in serum-free cultured HT29 spheroids was slightly higher than for serum-conditioned spheroid cultures in this particular experiment but the prominent difference between monolayer and spheroid cells was restored.

correlate with cell viability. Therefore, to verify anticancer drug sensitivity, Torisawa et al. (2005) monitored respiratory activity non-invasively in

MCF-7 MCTS by scanning electrochemical microscopy (SECM) and showed some correlation with the MTT equivalent WST-8 assay which was performed following spheroid dissociation (MTT: 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide; WST-8: 2-(2-methoxy-4-nitrophenyl)-3-(4-nitrophenyl)-5-(2,4-disulfophenyl)-2H tetrazolium monosodium). Also, a number of researchers have applied routine cell viability assays such as the MTT or similar test systems, but in most cases their utility for spheroids larger than 200 μm has either not been well tested, or spheroid dissociation was required. This step, however, may result in artificial cell loss in particular following therapeutic intervention and should be avoided. A slightly antiquated systematic study using various methods for assaying cytotoxic effects in spheroids involved only small spheroids without central necrosis (Kosaka et al. 1996). Thus, the potential of those methods for evaluating drug effects on hypoxic and cell-cycle arrested tumor cell populations in non-peripheral, viable regions present in most spheroid types with a diameter of $\geq 500 \mu\text{m}$ is limited. In another study, cell death in spheroids was determined by monitoring of lactate dehydrogenase (LDH) activity. However, here the methodology was adapted for the location of cell death processes in spheroid sections based on image cytometry and is thus not likely to be useful as a high throughput screening tool (De Witt Hamer et al. 2005). Aside from the common classical LDH measurement in supernatants of treated vs. untreated cells, various commercially available cytotoxicity/viability assays based on absorption, luminescence or fluorescence are routinely applied in monolayer cultures as well. Measurements are performed in supernatants or on intact or lysed cells, depending on the parameter measured, such as membrane integrity, intracellular activity or energetic status. We recently tested a number of such assay systems for their applicability to MCTS and found the Acid Phosphatase (APH) assay to be one of the most promising approaches amenable to high-throughput settings for non-dissociated spheroid cultures. We observed that the assay is reliable for monitoring cell viability in single spheroids, does not require spheroid dissociation and is linear and highly sensitive over a large cell concentration range and spheroid size, respectively, as shown for two colon carcinoma cell line spheroid types (Friedrich et al. 2007). The assay is thus simple, rapid and high-throughput compatible and has already been used to examine the effect of various drugs in spheroid culture. Moreover, our classical colon cancer spheroid model was recently adapted to serum-free conditions for highly standardized cell culturing as a prerequisite for reproducible and economical drug testing. This design also allows us to study the

impact of serum on drug efficacy. The APH assay can be applied to both culture conditions as documented in Figure 3b. The reduced efficacy of 5-FU in spheroids as compared to monolayers is not critically altered by serum supplementation. The dose response curves recorded in this experiment confirmed earlier data (Friedrich et al. 2007). Comparison with the growth kinetics clearly shows that cell viability is not necessarily reflected by spheroid integrity or regrowth over a period of 32 days. The potential for easy viability assessment with the APH assay as a function of time in culture after spheroid treatment will be useful in future studies intended to also verify if cell viability is indeed maintained throughout culturing and how this relates to growth delay and regrowth capacity for various treatment regimens.

It is clearly recognized that further technological improvements and increased standardization are required for a truly high-throughput drug screening platform with MCTS. However, significant progress over the past few years including the validation of the APH assay for rapid and reproducible determination of cell viability has essentially augmented the potential of spheroids for this purpose. The application of spheroids as tumor models for monitoring local drug penetration capacity, tissue distribution and binding properties is another challenge. Indeed, one advantage of the MCTS culture system is that drug candidates *must* penetrate in order to be highly effective. However, the analytical tools for such penetration studies, be they confocal laser scanning (Martin et al. 2003, Kostarelos et al. 2004, Mellor et al. 2005) or even multiphoton microscopy based, are still awaiting more intense validation as to their practical operating conditions. At this time, they have limited potential to be automated into rapid, high through-put approaches for spheroid analysis, especially if spheroid sectioning or dissociation has to be avoided. As mentioned above, incorporation of flow cytometric analysis should aid in the measurement of additional cell parameters, such as viability or metabolic activity.

MC(T)S – from homology to heterology

Recent progress has now cleared the way for using multicellular tumor spheroids as a model of tumor pathophysiology in anti-tumor drug screening. However, at the same time it also became evident again that the spheroid mono-culture system does not reflect the tumor cell environment in all aspects. Therefore, a number of complex spheroid-based systems to mimic cellular heterogeneity in tumor tissues have been developed as illustrated in Figure 4a. While these approaches are not yet established for drug screening, they are remarkable

tools to study heterologous cell-interactions in tumors *in vitro* under well-defined conditions and are thus considered major steps to gain insight into

particular drug and treatment related problems. A challenge specific to spheroid co-cultures is that the generation of spheroids with uniform fractions of the




Figure 4. Overview of potential spheroid co-culturing strategies of tumor and stromal cell types to study heterologous interactions in tumor tissue (a). The most frequent cell types used for co-culturing are tumor cells (T), immune cells (I), fibroblasts (F) and endothelial cells (EC) but other cell types are also applied, in particular in non-tumor approaches. Immune cells are primarily applied in suspensions while fibroblasts and EC have been used both in suspension and in pre-cultured adherent systems (spheroid, monolayer) for co-culturing with tumor cells and tumor cell spheroids, respectively. The combination of spheroids with multilayer co-culture systems (e.g., EC + F multilayers) is entertained but not yet realized. Representative literature for multicellular tumor spheroid co-culturing: (1) Tofilon et al. 1987, Thomsen et al. 1992, Frenzel et al. 1995, Nakamura et al. 1999, Boyd et al. 2001, 2002, Djordjevic et al. 2006, Spoettl et al. 2006; (2) Hauptmann et al. 1993, Konur et al. 1998, Siegert et al. 1999, Heimdahl et al. 2001, Olsnes et al. 2002, Furbert-Harris et al. 2003, Silzle et al. 2003, Gottfried et al. 2006; (3) Schuster et al. 1994, Kunz-Schughart et al. 1998, 2001; (4) Wartenberg et al. 2001, 2003, Wartenberg 2006, Gunther et al. 2007; (5) Broudy Boyé et al. 1994, Offner et al. 1996. Histomorphology of spheroid co-cultures is frequently documented in median sections following cryoconservation or paraffin-embedding and histological processing. Some examples of immunohistochemically or immunofluorescently stained median sections of spheroid co-cultures consisting of two or three cell types cultured with various strategies are shown in (b). Cell types and culture technique as detailed in the respective images.




Figure 4. (Continued).

different cell types has in some cases been difficult. Such uniformity would in principle be needed to use the system as a drug screening platform. The problem could be minimized by careful seeding of starting cell populations and/or subsequent spheroid analyses at the onset of treatment. Flow cytometric analysis of differentially labeled cell types in intact spheroids with or without spheroid sorting has some potential to help reduce heterogeneity as well, although this has not yet been tested out for screening purposes, as discussed earlier.

Co-culturing of different therapy-responsive tumor cells is one of the strategies of interest. Mixed, mosaic, or hybrid spheroids have been developed primarily from established cell lines or cell clones, e.g. defined by their difference in sensitivity to drugs such as cisplatin or BCNU (bis-chlorethyl-nitrosourea), or by their differentiation pattern, as described in bladder cancer or breast tumor cells and spheroids, respectively (Tofilon et al. 1987,

Bradley & Pitts 1994, Frenzel et al. 1995, Song et al. 2004, Persaud et al. 2005). Lately, a three cell line mixed spheroid system (termed tumoroid by the authors) based on three well known breast tumor cell lines, MCF-7, ZR-75-1 and MDA-MB-231, labeled with different fluorescent CellTracker Probes and cultured in the rotating wall vessel bioreactor has raised considerable attention (Vamvakidou et al. 2007). However, when using such systems, one has to bear in mind the potentially different behavior of the cell types in 3-D culture, e.g., with respect to cell attachment, survival and proliferative as well as migratory activity. These characteristics critically affect the histomorphology and cellular distribution in a mixed spheroid, even if identical cell numbers are applied for each line. The proportion of the cells not only at the time of seeding but rather at the time of treatment and their location within the spheroid are just two of the critical parameters that may modulate the

spheroid's response to drug exposure. Taking into account the historical shift in the drug development world from generally cytotoxic towards target-specific drug candidates, there may be a particular use for the 3-D co-culturing of genetically well defined tumor cell clones that differ exclusively in the gene of interest. Such a transfectant mosaic spheroid model composed of two cell populations derived from a single cell line and differing by just one gene was recently used in an interesting study series on target-specific treatment modalities, i.e. focusing on targeted radiotherapy and evaluation of the 'radiological bystander effect' conferred by a particular beta-emitting radiopharmaceutical (Boyd et al. 2002, 2004).

Other approaches of spheroid co-culturing include various stromal cell types. Under investigation are immune cells such as monocytes, macrophages, dendritic cells (DC), T-lymphocytes or even NK (natural killer) cells. These latter cell types are primarily added as cell suspensions to tumor or fibroblast spheroids to monitor migration and maturation or activation processes affected by the tumor environment. With one such system, Spoettl et al. recently studied intestinal differentiation of monocytes by using a spheroid model of transfected HT29 colon cancer cells specifically overexpressing one of two chemokines of interest (Spoettl et al. 2006, 2007). One of these chemokines (MCP-1 = monocyte chemoattractant protein-1) is also hypothesized to play a role in monocyte migration into breast tumor-derived stromal fibroblast spheroids (Silzle et al. 2003). The adverse impact of tumor cell lactic acid production on the differentiation of DC has been shown in spheroids of human J82 bladder and MelIm melanoma cells (Gottfried et al. 2006). The MelIm melanoma spheroid model has also been employed in a study to verify that tumor derived lactic acid critically suppresses the function of human cytotoxic T lymphocytes (CTL) (Fischer et al. 2007). Since the present article focuses on drug testing in spheroids, the numerous basic tumor-biological studies performed over the past 15 years that are based on tumor-immune cells co-cultured with spheroids cannot be comprehensively summarized. However, we want to draw attention to the recent review article 'Brave little world: spheroids as an *in vitro* model to study tumor-immune-cell interactions' by Gottfried et al. (2006). With respect to therapy, it is noteworthy that the spheroid model has been found useful in the development of mathematical models to use macrophages as vehicles for drug delivery to hypoxic tumor areas (Owen et al. 2004).

Different spheroid co-culturing techniques have also been applied to monitor interactions between tumor cells and stromal fibroblast. Experimental

settings range from mixed spheroids, MCTS cultured on fibroblast monolayers (Brouty-Boye et al. 1994), MCTS co-cultured with pre-established fibroblast spheroids – also termed confrontation cultures (Schuster et al. 1994, Kunz-Schughart et al. 1998), fibroblast suspension added to MCTS, to the incubation of fibroblast spheroids with tumor cell suspensions (Brabletz et al. 2004). The latter technique was found useful to reflect several important pathological phenomena, for example the migration of non-aggregating tumor cells into fibroblastic stroma at the invasion front. It is hypothesized that fibroblasts in tumor tissues and in spheroid co-cultures, which are known to produce high amounts of ECM molecules, release numerous cytokines and growth factors and also directly interact with the tumor cells (Kunz-Schughart & Knuechel 2002, Kunz-Schughart & Freyer 2002), which may critically affect therapeutic efficacy, e.g. by altering the distribution of macromolecular drug candidates. Undoubtedly, such co-culture models may be useful to verify the impact of the stromal environment on drug efficacy. However, the practical details of their application in drug screening are not yet readily envisioned. On the contrary, their use in radiotherapy testing is more obvious, largely for historical reasons. In fact, the idea to score the effects of radiation and of other treatments on tumor cell survival and clonogenicity via the application of fibroblast-tumor cell hybrid spheroids using supraethally irradiated fibroblasts as feeder cells is not new, though the technology has only been applied in a limited number of cases (Djordjevic & Lange 1990, Lange et al. 1992, Djordjevic et al. 1998). In a recent approach, Djordjevic and co-workers (Djordjevic & Lange 2006) once more presented experimental indications for a bystander-like effect in such hybrid spheroids. Here, the presence of even highly-irradiated neighboring cells reduced clonogenic survival of (tumor) test cells in the 3D spheroid co-culture system. Clonogenicity of test cells was evaluated via growth analysis of whole spheroids composed of defined proportions of non-clonogenic feeder cells and a minority of these test cells (1–10%) (Djordjevic & Lange 2004, 2006). The observations with this model evidently demonstrate that using the (most) appropriate *in vitro* test system is a critical aspect for radiation sensitivity testing, and may ultimately guide the selection of predictive assays for clinical efficacy.

In addition to immune cells and fibroblasts, the tumor stroma of course contains varying amounts of endothelial cells (EC). Thus, co-culturing spheroids with EC is another field of research (Oudar 2000) which is bound to become more relevant in the near future, especially in the testing

of combined therapeutic approaches with anti-angiogenic compounds. The technique of culturing MCTS on confluent EC with an underlying ECM has been used in the past to study aspects of tumor cell evasion (Vermeulen et al. 1996). Mixed cultures of tumor and EC to study tumor angiogenesis or the application of MCTS and endothelial cell suspensions to allow EC migration, carried out over many years, have had little success. However, in 2004, Timmins et al. described an EC network in HCT116 colon cancer and even more pronounced in HepG2 hepatocellular carcinoma spheroids by using the hanging drop culturing technique (Timmins et al. 2004). It is not clear whether such EC networks may be developed in other tumor spheroid types. Observations in fibroblast spheroids co-cultured with EC suspensions indicate that at least for some tumor cell types the fibroblastic stroma may be either required or subsidiary for the formation of networking tubular structures in spheroids (Kunz-Schughart et al. 2006). In contrast to tumor or fibroblast EC co-cultured spheroids showing EC network formation, the co-culture of EC and smooth muscle cells in spheroids results in an entirely different morphology, with EC located in the spheroid periphery surrounding a smooth muscle cell core (Korff et al. 2001). This co-culture approach is based on a pure EC spheroid system developed earlier (Korff & Augustin 1998), which depends on allowing cell-cell contacts and can be maintained in culture by the supplementation of methocylcellulose (methocel). Such EC spheroids can be collected and embedded into three-dimensional ECM gels for monitoring EC migration processes and to identify modulators of angiogenesis, or they can be co-cultured with other spheroid types, even with organotypic tumor or fragment spheroids. Such approaches have gained some experimental importance in the study and manipulation of angiogenesis in gliomas and other brain tumors (Goldbrunner et al. 2000, 2004). Finally, it should be mentioned, that tumor-induced angiogenesis has also been studied in an interesting system of MCTS in confrontation culture with embryoid bodies (Wartenberg et al. 2001, 2006, Gunther et al. 2007). This *in vitro* system, however, has some limitations due to the necessary addition of non-human pluripotent, i.e. mouse, embryonic stem cells.

Some examples of immunohistochemically or immunofluorescently stained median sections from different types of histologically processed spheroid co-cultures are shown in Figure 4b. In addition to such co-cultures from cell lines, cell clones and primary cells, there is growing interest centering around the use of so-called fragment spheroids grown from tumor biopsies/tissues as already

mentioned above (de Ridder et al. 2000, Heimdal et al. 2001, Kross et al. 2005, Olsnes et al. 2005). This seems of particular relevance for an individualized, patient-specific therapy testing and may be less feasible for massive parallel drug screening. When using such systems for drug testing, one has to for example consider the potential interindividual variance of multiple fragment spheroids derived from different sections of a single tumor. While such variation may in fact be representative of the tumor's real heterogeneity, it may also produce numerous analytical problems for establishing a robust drug testing platform. Having said that, tumor biopsy material may on the other hand contain so-called initiating tumor cells and tumor stem/progenitor cell populations, which are obviously of particular interest for new treatment modalities and the technique may thus have some not yet fully appreciated benefits. This hypothesis gained increased attention with the recent observations indicating that particular tumor initiating cell populations, not only of brain tumors but also of colorectal tumors, could exclusively be maintained in culture when grown as spheres (Zhang et al. 2005, Ricci-Vitiani et al. 2007). This intriguing aspect will no doubt receive thorough consideration in some future drug screening initiatives.

Acknowledgements

The authors gratefully acknowledge the technical assistance of Mrs Marit Wondrak. The experimental work documented herein was supported by the Society for Biomolecular Sciences and by the German Federal Ministry of Education and Research through grants 0313143 and 01ZZ0502 to LKS.

References

- Acker H, Carlsson J, Mueller-Klieser W, Sutherland RM. 1987. Comparative pO₂ measurements in cell spheroids cultured with different techniques. *British Journal of Cancer* 56:325–327.
- Audran R, Dazord L, Toujas L. 1994. Interactions between human macrophages and tumor cells in three-dimensional cultures. *Cancer Immunology, Immunotherapy* 39:299–304.
- Balis FM. 2002. Evolution of anticancer drug discovery and the role of cell-based screening. *Journal of the National Cancer Institute* 94:78–79.
- Ballangrud AM, Yang WH, Charlton DE, McDevitt MR, Hamacher KA, Panageas KS, Ma D, Bander NH, Scheinberg DA, Sgouros G. 2001. Response of LNCaP spheroids after treatment with an alpha-particle emitter (213Bi)-labeled anti-prostate-specific membrane antigen antibody (J591). *Cancer Research* 61:2008–2014.
- Ballangrud AM, Yang WH, Dnistrian A, Lampen NM, Sgouros G. 1999. Growth and characterization of LNCaP prostate cancer cell spheroids. *Clinical Cancer Research* 5:3171–3176.

- Banks SD, Waters KA, Barrett LL, Dickerson S, Pendergast W, Smith GK. 1994. Destruction of WiDr multicellular tumor spheroids with the novel thymidylate synthase inhibitor 1843U89 at physiological thymidine concentrations. *Cancer Chemotherapy and Pharmacology* 33: 455–459.
- Bardies M, Thedrez P, Gestin JF, Marcille BM, Guerreau D, Faivre-Chauvet A, Mahe M, Sai-Maurel C, Chatal JF. 1992. Use of multi-cell spheroids of ovarian carcinoma as an intraperitoneal radio-immunotherapy model: Uptake, retention kinetics and dosimetric evaluation. *International Journal of Cancer* 50:984–991.
- Bartholoma P, Impidjati, Reininger-Mack A, Zhang Z, Thielecke H, Robitzki A. 2005. A more aggressive breast cancer spheroid model coupled to an electronic capillary sensor system for a high-content screening of cytotoxic agents in cancer therapy: 3-dimensional *in vitro* tumor spheroids as a screening model. *Journal of Biomolecular Screening* 10:705–714.
- Berglund A, Glimelius B, Bergh J, Brodin O, Fjallskog ML, Hagberg H, von HA, Larsson R, Tholander B, de la TM, Astrom G, Oberg K, Paro G, Nygren P. 2002. Selection of chemotherapy by *ex vivo* assessment of tumor sensitivity to cytotoxic drugs: Results of a clinical trial. *Medical Oncology* 19:151–159.
- Bhadiraju K, Chen CS. 2002. Engineering cellular microenvironments to improve cell-based drug testing. *Drug Discovery Today* 11:612–620.
- Boyd M, Mairs RJ, Cunningham SH, Mairs SC, McCluskey A, Livingstone A, Stevenson K, Brown MM, Wilson L, Carlin S, Wheldon TE. 2001. A gene therapy/targeted radiotherapy strategy for radiation cell kill by [131I] meta-iodobenzylguanidine. *Journal of Gene Medicine* 3:165–172.
- Boyd M, Mairs SC, Stevenson K, Livingstone A, Clark AM, Ross SC, Mairs RJ. 2002. Transfectant mosaic spheroids: A new model for evaluation of tumour cell killing in targeted radiotherapy and experimental gene therapy. *Journal of Gene Medicine* 4:567–576.
- Boyd M, Mairs RJ, Keith WN, Ross SC, Welsh P, Akabani G, Owens J, Vaidyanathan G, Carruthers R, Dorrens J, Zalutsky MR. 2004. An efficient targeted radiotherapy/gene therapy strategy utilising human telomerase promoters and radioastatine and harnessing radiation-mediated bystander effects. *Journal of Gene Medicine* 6:937–947.
- Brabletz T, Spaderna S, Kolb J, Hlubek F, Faller G, Bruns CJ, Jung A, Nentwich J, Duluc I, Domon-Dell C, Kirchner T, Freund JN. 2004. Down-regulation of the homeodomain factor Cdx2 in colorectal cancer by collagen type I: An active role for the tumor environment in malignant tumor progression. *Cancer Research* 64:6973–6977.
- Bradley C, Pitts J. 1994. The use of genetic marking to assess the interaction of sensitive and multidrug-resistant cells in mixed culture. *British Journal of Cancer* 70:795–798.
- Braun RD, Gadianu M, Vistisen KS, Roberts RL, Berkowitz BA. 2007. Manganese-enhanced MRI of human choroidal melanoma xenografts. *Investigative Ophthalmology & Visual Science* 48:963–967.
- Bredel-Geissler A, Karbach U, Walenta S, Vollrath L, Mueller-Klieser W. 1992. Proliferation-associated oxygen consumption and morphology of tumor cells in monolayer and spheroid culture. *Journal of Cellular Physiology* 153:44–52.
- Brouty-Boye D, Mainguene C, Magnien V, Israel L, Beaupain R. 1994. Fibroblast-mediated differentiation in human breast carcinoma cells (MCF-7) grown as nodules *in vitro*. *International Journal of Cancer* 56:731–735.
- Buchsbaum DJ, Langmuir VK, Wessels BW. 1993. Experimental radioimmunotherapy. *Medical Physics* 20:551–567.
- Buffa FM, Davidson SE, Hunter RD, Nahum AE, West CM. 2001. Incorporating biologic measurements (SF(2), CFE) into a tumor control probability model increases their prognostic significance: A study in cervical carcinoma treated with radiation therapy. *International Journal of Radiation Biology and Related Studies in Physics, Chemistry, and Medicine* 50:1113–1122.
- Carlsson J, Acker H. 1988. Relations between pH, oxygen partial pressure and growth in cultured cell spheroids. *International Journal of Cancer* 42:715–720.
- Carlsson J, Nederman T. 1989. Tumour spheroid technology in cancer therapy research. *European Journal of Cancer and Clinical Oncology* 25:1127–1133.
- Carlsson J, Yuhas JM. 1984. Liquid-overlay culture of cellular spheroids. *Recent Results Cancer Research* 95:1–23.
- Casciari JJ, Hollingshead MG, Alley MC, Mayo JG, Malspeis L, Miyauchi S, Grever MR, Weinstein JN. 1994. Growth and chemotherapeutic response of cells in a hollow-fiber *in vitro* solid tumor model. *Journal of the National Cancer Institute* 86:1846–1852.
- Casciari JJ, Sotirchos SV, Sutherland RM. 1988. Glucose diffusivity in multicellular tumor spheroids. *Cancer Research* 48:3905–3909.
- Casey RC, Burleson KM, Skubitz KM, Pambuccian SE, Oegema TR, Jr., Ruff LE, Skubitz AP. 2001. Beta 1-integrins regulate the formation and adhesion of ovarian carcinoma multicellular spheroids. *American Journal of Physiology* 159:2071–2080.
- Cate CC, Douple EB, Andrews KM, Pettengill OS, Curphey TJ, Sorenson GD, Maurer LH. 1984. Calcitonin as an indicator of the response of human small cell carcinoma of the lung cells to drugs and radiation. *Cancer Research* 44:949–954.
- Chan WS, West CM, Moore JV, Hart IR. 1991. Photocytotoxic efficacy of sulphonated species of aluminium phthalocyanine against cell monolayers, multicellular spheroids and *in vivo* tumours. *British Journal of Cancer* 64:827–832.
- Chaw KC, Manimaran M, Tay EH, Swaminathan S. 2007. Multi-step microfluidic device for studying cancer metastasis. *Lab on a Chip* 7:1041–1047.
- Chen TR, Drabkowski D, Hay RJ, Macy M, Peterson W Jr. 1987. WiDr is a derivative of another colon adenocarcinoma cell line, HT-29. *Cancer Genetics and Cytogenetics* 27:125–134.
- Cheng FM, Hansen EB, Taylor CR, Epstein AL. 1991. Diffusion and binding of monoclonal antibody TNT-1 in multicellular tumor spheroids. *Journal of the National Cancer Institute* 83:200–204.
- Clejan S, O'Connor K, Rosensweig N. 2001. Tri-dimensional prostate cell cultures in simulated microgravity and induced changes in lipid second messengers and signal transduction. *Journal of Cellular and Molecular Medicine* 5:60–73.
- de Ridder L, Cornelissen M, de Ridder D. 2000. Autologous spheroid culture: A screening tool for human brain tumour invasion. *Critical Reviews in Oncology/Hematology* 36:107–122.
- De Witt Hamer PC, Jonker A, Leenstra S, Ruijter JM, Van Noorden CJ. 2005. Quantification of viability in organotypic multicellular spheroids of human malignant glioma using lactate dehydrogenase activity: A rapid and reliable automated assay. *Journal of Histochemistry and Cytochemistry* 53:23–34.
- Del Duca D, Werbowetski T, Del Maestro RF. 2004. Spheroid preparation from hanging drops: Characterization of a model of brain tumor invasion. *Journal of Neuro Oncology* 67:295–303.
- Debnath J, Brugge JS. 2005. Modelling glandular epithelial cancers in three-dimensional cultures. *Nature Reviews Cancer* 5:675–688.
- Desoize B, Gimonet D, Jardiller JC. 1998. Cell culture as spheroids: An approach to multicellular resistance. *Anticancer Research* 18:4147–4158.

- Desoize B, Jardillier J. 2000. Multicellular resistance: A paradigm for clinical resistance? *Critical Reviews in Oncology/Hematology* 36:193–207.
- Djordjevic B, Lange CS. 1990. Clonogenicity of mammalian cells in hybrid spheroids: A new assay method. *Radiation and Environmental Biophysics* 29:31–46.
- Djordjevic B, Lange CS, Allison RR, Rotman M. 1993. Response of primary colon cancer cells in hybrid spheroids to 5-fluorouracil. *Cancer Investigation* 11:291–298.
- Djordjevic B, Lange CS, Schwartz MS, Rotman M. 1998. Clonogenic inactivation of colon cancer-derived cells treated with 5-fluorouracil and indomethacin in hybrid spheroids. *Acta Oncologica* 37:735–739.
- Djordjevic B, Lange CS. 2004. Hybrid spheroids as a tool for prediction of radiosensitivity in tumor therapy. *Indian Journal of Experimental Biology* 42:443–447.
- Djordjevic B, Lange CS. 2006. Cell-cell interactions in spheroids maintained in suspension. *Acta Oncologica* 45:412–420.
- Dolfini E, Elli L, Roncoroni L, Costa B, Colleoni MP, Lorusso V, Ramponi S, Braidotti P, Ferrero S, Falini ML, Bardella MT. 2005. Damaging effects of gliadin on three-dimensional cell culture model. *World Journal of Gastroenterology* 11:5973–5977.
- Donaldson JT, Tucker JA, Keane TE, Walther PJ, Webb KS. 1990. Characterization of a new model of human prostatic cancer: The multicellular tumor spheroid. *International Journal of Cancer* 46:238–244.
- Douple EB, Cate CC, Curphey TJ, Pettengill OS, Sorenson GD, Maurer LH. 1985. Evaluation of drug efficacy *in vitro* using human small cell carcinoma of the lung spheroids. *Cancer* 56:1918–1925.
- Dulong JL, Legallais C, Darquy S, Reach G. 2002. A novel model of solute transport in a hollow-fiber bioartificial pancreas based on a finite element method. *Biotechnology and Bioengineering* 78:576–582.
- Durand RE, Olive PL. 1992. Evaluation of bioreductive drugs in multicell spheroids. *International Journal of Radiation Biology and Related Studies in Physics, Chemistry, and Medicine* 22:689–692.
- Durand RE, Sham E. 1998. The lifetime of hypoxic human tumor cells. *International Journal of Radiation Biology and Related Studies in Physics, Chemistry, and Medicine* 42:711–715.
- Eley KW, Benedict SH, Chung TD, Kavanagh BD, Broaddus WC, Schmidt-Ullrich RK, Lin PS. 2002. The effects of pentoxifylline on the survival of human glioma cells with continuous and intermittent stereotactic radiosurgery irradiation. *International Journal of Radiation Biology and Related Studies in Physics, Chemistry, and Medicine* 54:542–550.
- Ellison G, Klinowska T, Westwood RF, Docter E, French T, Fox JC. 2002. Further evidence to support the melanocytic origin of MDA-MB-435. *Molecular Pathology* 55:294–299.
- Enmon R, Yang WH, Ballangrud AM, Solit DB, Heller G, Rosen N, Scher HI, Sgouros G. 2003. Combination treatment with 17-N-allylamino-17-demethoxy geldanamycin and acute irradiation produces supra-additive growth suppression in human prostate carcinoma spheroids. *Cancer Research* 63:8393–8399.
- Enmon RM Jr, O'Connor KC, Song H, Lacks DJ, Schwartz DK. 2002. Aggregation kinetics of well and poorly differentiated human prostate cancer cells. *Biotechnology and Bioengineering* 80:580–588.
- Erlachman C, Wu A. 1992. Resistance of MGH-U1 bladder cancer spheroids to vincristine. *Anticancer Research* 12:1233–1236.
- Eshleman JS, Carlson BL, Mladek AC, Kastner BD, Shide KL, Sarkaria JN. 2002. Inhibition of the mammalian target of rapamycin sensitizes U87 xenografts to fractionated radiation therapy. *Cancer Research* 62:7291–7297.
- Essand M, Nilsson S, Carlsson J. 1993. Growth of prostatic cancer cells, DU 145, as multicellular spheroids and effects of estramustine. *Anticancer Research* 13:1261–1268.
- Fang D, Nguyen TK, Leishear K, Finko R, Kulp AN, Hotz S, Van Belle PA, Xu X, Elder DE, Herlyn M. 2005. A tumorigenic subpopulation with stem cell properties in melanomas. *Cancer Research* 65:9328–9337.
- Feder-Mengus C, Ghosh S, Weber WP, Wyler S, Zajac P, Terracciano L, Oertli D, Heberer M, Martin I, Spagnoli GC, Reschner A. 2007. Multiple mechanisms underlie defective recognition of melanoma cells cultured in three-dimensional architectures by antigen-specific cytotoxic T lymphocytes. *British Journal of Cancer* 96:1072–1082.
- Fehlauer F, Muench M, Rades D, Stalpers LJ, Leenstra S, van dV, Slotman B, Smid EJ, Sminia P. 2005. Effects of irradiation and cisplatin on human glioma spheroids: Inhibition of cell proliferation and cell migration. *Journal of Cancer Research and Clinical Oncology* 131:723–732.
- Fehlauer F, Muench M, Smid EJ, Slotman B, Richter E, van dV, Sminia P. 2006. Combined modality therapy of gemcitabine and irradiation on human glioma spheroids derived from cell lines and biopsy tissue. *Oncology Reports* 15:97–105.
- Filippovich IV, Sorokina N, Robillard N, Faivre-Chauvet A, Bardies M, Chatal JF. 1996. Cell death induced by a ¹³¹I-labeled monoclonal antibody in ovarian cancer multicell spheroids. *Nuclear Medicine and Biology* 23:623–626.
- Fischer K, Hoffmann P, Voelkl S, Meidenbauer N, Ammer J, Edinger M, Gottfried E, Schwarz S, Rothe G, Hoves S, Renner K, Timischl B, Mackensen A, Kunz-Schughart L, Andreesen R, Krause SW, Kreutz M. 2007. Inhibitory effect of tumor cell derived lactic acid on human T cells. *Blood* 109:3812–3819.
- Feder-Mengus C, Ghosh S, Weber WP, Wyler S, Zajac P, Terracciano L, Oertli D, Heberer M, Martin I, Spagnoli GC, Reschner A. 2007. Multiple mechanisms underlie defective recognition of melanoma cells cultured in three-dimensional architectures by antigen-specific cytotoxic T lymphocytes. *British Journal of Cancer* 96:1072–1082.
- Francia G, Green SK, Bocci G, Man S, Emmenegger U, Ebos JM, Weinerman A, Shaked Y, Kerbel RS. 2005. Down-regulation of DNA mismatch repair proteins in human and murine tumor spheroids: Implications for multicellular resistance to alkylating agents. *Molecular Cancer Therapeutics* 4:1484–1494.
- Frankel A, Man S, Elliott P, Adams J, Kerbel RS. 2000. Lack of multicellular drug resistance observed in human ovarian and prostate carcinoma treated with the proteasome inhibitor PS-341. *Clinical Cancer Research* 6:3719–3728.
- Friedrich J, Eder W, Castaneda J, Doss M, Huber E, Edner R, Kunz-Schughart LA. 2007. A reliable tool to determine cell viability in complex 3-d culture: the acid phosphatase assay. *Journal of Biomolecular Screening* 12:925–937.
- Frenzel KR, Saller RM, Kummermehr J, Schultz-Hector S. 1995. Quantitative distinction of cisplatin-sensitive and -resistant mouse fibrosarcoma cells grown in multicell tumor spheroids. *Cancer Research* 55:386–391.
- Freyer DR. 1992. Tumor spheroids in radiobiology research. Bjerkvig R, editor. Boca Raton, USA: CRC Press. pp 217–275.
- Freyer JP, Wilder ME, Jett JH. 1987. Viable sorting of intact multicellular spheroids by flow cytometry. *Cytometry* 8:427–436.
- Freyer JP. 1988. Role of necrosis in regulating the growth saturation of multicellular spheroids. *Cancer Research* 48:2432–2439.
- Freyer JP, Schor PL. 1989. Regrowth kinetics of cells from different regions of multicellular spheroids of four cell lines. *Journal of Cellular Physiology* 138:384–392.

- Freyer JP, Fillak D, Jett JH. 1989. Use of xanthan gum to suspend large particles during flow cytometric analysis and sorting. *Cytometry* 6:803–806.
- Freyer JP, Fink NH, Schor PL, Coulter JR, Neeman M, Sillerud LO. 1990. A system for viably maintaining a stirred suspension of multicellular spheroids during NMR spectroscopy. *NMR in Biomedicine* 3:195–205.
- Freyer JP, Schor PL, Jarrett KA, Neeman M, Sillerud LO. 1991. Cellular energetics measured by phosphorous nuclear magnetic resonance spectroscopy are not correlated with chronic nutrient deficiency in multicellular tumor spheroids. *Cancer Research* 51:3831–3837.
- Furber-Harris PM, Laniyan I, Harris D, Dunston GM, Vaughn T, Abdelnaby A, Parish-Gause D, Oredipe OA. 2003. Activated eosinophils infiltrate MCF-7 breast multicellular tumor spheroids. *Anticancer Research* 23:71–78.
- Garraway LA, Widlund HR, Rubin MA, Getz G, Berger AJ, Ramaswamy S, Beroukhi R, Milner DA, Granter SR, Du J, Lee C, Wagner SN, Li C, Golub TR, Rimm DL, Meyerson ML, Fisher DE, Sellers WR. 2005. Integrative genomic analyses identify MITF as a lineage survival oncogene amplified in malignant melanoma. *Nature* 436:117–122.
- Genc M, Castro KN, Barten-van RA, Stalpers LJ, Haveman J. 2004. Enhancement of effects of irradiation by gemcitabine in a glioblastoma cell line and cell line spheroids. *Journal of Cancer Research and Clinical Oncology* 130:45–51.
- Ghosh S, Spagnoli GC, Martin I, Ploegert S, Demougins P, Heberer M, Reschner A. 2005. Three-dimensional culture of melanoma cells profoundly affects gene expression profile: A high density oligonucleotide array study. *Journal of Cellular Physiology* 204:522–531.
- Gillies RJ, Galons JP, McGovern KA, Scherer PG, Lien YH, Job C, Ratcliff R, Chapa F, Cerdan S, Dale BE. 1993. Design and application of NMR-compatible bioreactor circuits for extended perfusion of high-density mammalian cell cultures. *NMR in Biomedicine* 6:95–104.
- Goldbrunner RH, Bendszus M, Tonn JC. 2004. Models for angiogenesis in gliomas. *Cancer Treatment and Research* 117:115–135.
- Goldbrunner RH, Wagner S, Roosen K, Tonn JC. 2000. Models for assessment of angiogenesis in gliomas. *Journal of Neuro Oncology* 50:53–62.
- Gorlach A, Fandrey J, Holtermann G, Acker H. 1994. Effects of cobalt on haem proteins of erythropoietin-producing HepG2 cells in multicellular spheroid culture. *FEBS Letters* 348:216–218.
- Gorlach A, Herter P, Hentschel H, Frosch PJ, Acker H. 1994. Effects of rIFN beta and rIFN gamma on growth and morphology of two human melanoma cell lines: Comparison between two- and three-dimensional culture. *International Journal of Cancer* 56:249–254.
- Gottfried E, Kunz-Schughart LA, Andreessen R, Kreutz M. 2006. Brave little world: Spheroids as an *in vitro* model to study tumor-immune-cell interactions. *Cell Cycle* 5:691–695.
- Gottfried E, Kunz-Schughart LA, Ebner S, Mueller-Klieser W, Hoves S, Andreessen R, Mackensen A, Kreutz M. 2006. Tumor-derived lactic acid modulates dendritic cell activation and antigen expression. *Blood* 107:2013–2021.
- Gudjonsson T, Rønnov-Jessen L, Villadsen R, Bissell MJ, Petersen OW. 2003. To create the correct microenvironment: Three-dimensional heterotypic collagen assays for human breast epithelial morphogenesis and neoplasia. *Methods* 30:247–255.
- Gunther S, Ruhe C, Derikito MG, Bose G, Sauer H, Wartenberg M. 2007. Polyphenols prevent cell shedding from mouse mammary cancer spheroids and inhibit cancer cell invasion in confrontation cultures derived from embryonic stem cells. *Cancer Letters* 250:25–35.
- Hauptmann S, Zwadlo-Klarwasser G, Jansen M, Klosterhalfen B, Kirkpatrick CJ. 1993. Macrophages and multicellular tumor spheroids in co-culture: A three-dimensional model to study tumor-host interactions. Evidence for macrophage-mediated tumor cell proliferation and migration. *American Journal of Pathology* 143:1406–1415.
- Hedlund TE, Duke RC, Miller GJ. 1999. Three-dimensional spheroid cultures of human prostate cancer cell lines. *Prostate* 41:154–165.
- Heimdal JH, Olsnes C, Olofsson J, Aarstad HJ. 2001. Monocyte and monocyte-derived macrophage secretion of MCP-1 in co-culture with autologous malignant and benign control fragment spheroids. *Cancer Immunology, Immunotherapy* 50:300–306.
- Heimdal JH, Aarstad HJ, Olsnes C, Olofsson J. 2001. Human autologous monocytes and monocyte-derived macrophages in co-culture with carcinoma F-spheroids secrete IL-6 by a non-CD14-dependent pathway. *Scandinavian Journal of Immunology* 53:162–170.
- Heuser M, Ringert RH, Zoeller G, Hemmerlein B. 2003. Dynamic assessment of angiogenesis in renal cell carcinoma spheroids by intravital microscopy. *Journal of Urology* 169:1267–1270.
- Hicks KO, Ohms SJ, van Zijl PL, Denny WA, Hunter PJ, Wilson WR. 1997. An experimental and mathematical model for the extravascular transport of a DNA intercalator in tumours. *British Journal of Cancer* 76:894–903.
- Hoevel T, Macek R, Swisshelm K, Kubbies M. 2004. Reexpression of the TJ protein CLDN1 induces apoptosis in breast tumor spheroids. *International Journal of Cancer* 108:374–383.
- Horan HP, Colcher D, Salomon D, Ridge J, Noguchi P, Schlom J. 1985. Influence of spatial configuration of carcinoma cell populations on the expression of a tumor-associated glycoprotein. *Cancer Research* 45:833–840.
- Huygens A, Huyghe D, Bormans G, Verbruggen A, Kamuhabwa AR, Roskams T, De Witte PA. 2003. Accumulation and photocytotoxicity of hypericin and analogs in two- and three-dimensional cultures of transitional cell carcinoma cells. *Photochemistry and Photobiology* 78:607–614.
- Hystad ME, Rofstad EK. 1994. Oxygen consumption rate and mitochondrial density in human melanoma monolayer cultures and multicellular spheroids. *International Journal of Cancer* 57:532–537.
- Ingram M, Techy GB, Saroufeem R, Yazan O, Narayan KS, Goodwin TJ, Spaulding GF. 1997. Three-dimensional growth patterns of various human tumor cell lines in simulated microgravity of a NASA bioreactor. *In Vitro Cellular & Developmental Biology – Animal* 33:459–466.
- Ivascu A, Kubbies M. 2006. Rapid generation of single-tumor spheroids for high-throughput cell function and toxicity analysis. *Journal of Biomolecular Screening* 11:922–932.
- Jansen M, Dykstra M, Lee JI, Stables J, Topley P, Knick VC, Mullin RJ, Duch DS, Smith GK. 1994. Effect of purine synthesis inhibition on WiDr spheroids *in vitro* or on WiDr or colon 38 tumors *in vivo*. Complete growth inhibition but not regression. *Biochemical Pharmacology* 47:1067–1078.
- Johnson JI, Decker S, Zaharevitz D, Rubinstein LV, Venditti JM, Schepartz S, Kalyandrug S, Christian M, Arbuck S, Hollingshead M, Sausville EA. 2001. Relationships between drug activity in NCI preclinical *in vitro* and *in vivo* models and early clinical trials. *British Journal of Cancer* 84:1424–1431.
- Jorgensen K, Skrede M, Cruciani V, Mikalsen SO, Slipicevic A, Florenes VA. 2005. Phorbol ester phorbol-12-myristate-13-acetate promotes anchorage-independent growth and survival of melanomas through MEK-independent activation of ERK1/2. *Biochemical and Biophysical Research Communications* 329:266–274.

- Kamuhabwa AA, Huygens A, De Witte PA. 2003. Photodynamic therapy of transitional cell carcinoma multicellular tumor spheroids with hypericin. *International Journal of Oncology* 23:1445–1450.
- Kan O, Griffiths L, Baban D, Iqbal S, Uden M, Spearman H, Slingsby J, Price T, Esapa M, Kingsman S, Kingsman A, Slade A, Naylor S. 2001. Direct retroviral delivery of human cytochrome P450 2B6 for gene-directed enzyme prodrug therapy of cancer. *Cancer Gene Therapy* 8:473–482.
- Kawai K, Hayashi H, Ozaki Y, Saijo K, Liu SQ, Akaza H, Ohno T. 2001. Assessment of the number of local cytotoxic T lymphocytes required for degradation of micrometer-size tumor spheroids. *Cytotechnology* 37:31–40.
- Kelm JM, Timmins NE, Brown CJ, Fussenegger M, Nielsen LK. 2003. Method for generation of homogeneous multicellular tumor spheroids applicable to a wide variety of cell types. *Biotechnology and Bioengineering* 83:173–180.
- Kerbel RS. 2003. Human tumor xenografts as predictive preclinical models for anticancer drug activity in humans: Better than commonly perceived – but they can be improved. *Cancer Biology & Therapy* 2:S134–S139.
- Kerr DJ, Smart HE. 1991. The relationship between cytotoxic drug exposure and tumour cell kill, *in vitro* and *in vivo*. *In Vivo* 5:385–388.
- Khaitan D, Chandna S, Arya MB, Dwarakanath BS. 2006. Differential mechanisms of radiosensitization by 2-deoxy-D-glucose in the monolayers and multicellular spheroids of a human glioma cell line. *Cancer Biology & Therapy* 5:1142–1151.
- Kim JH, Ahn SI, Kim JH, Zin WC. 2007. Evaporation of water droplets on polymer surfaces. *Langmuir* 23:6163–6169.
- Kilani RT, Tamimi Y, Hanel EG, Wong KK, Karmali S, Lee PW, Moore RB. 2003. Selective reovirus killing of bladder cancer in a co-culture spheroid model. *Virus Research* 93:1–12.
- Kohno N, Ohnuma T, Biller HF, Holland JF. 1988. Effects of cisplatin plus fluorouracil vs cisplatin plus cytarabine on head and neck squamous multicellular tumor spheroids. *Archives of Otolaryngology – Head & Neck Surgery* 114:157–161.
- Konduri SD, Tasiou A, Chandrasekar N, Rao JS. 2001. Overexpression of tissue factor pathway inhibitor-2 (TFPI-2), decreases the invasiveness of prostate cancer cells *in vitro*. *International Journal of Oncology* 18:127–131.
- Konur A, Kreutz M, Knuchel R, Krause SW, Andreesen R. 1998. Cytokine repertoire during maturation of monocytes to macrophages within spheroids of malignant and non-malignant urothelial cell lines. *International Journal of Cancer* 78:648–653.
- Korff T, Augustin HG. 1998. Integration of endothelial cells in multicellular spheroids prevents apoptosis and induces differentiation. *Journal of Cell Biology* 143:1341–1352.
- Korff T, Kimmina S, Martiny-Baron G, Augustin HG. 2001. Blood vessel maturation in a 3-dimensional spheroidal coculture model: Direct contact with smooth muscle cells regulates endothelial cell quiescence and abrogates VEGF responsiveness. *FASEB Journal* 15:447–457.
- Kosaka T, Fukaya K, Tsuboi S, Pu H, Ohno T, Tsuji T, Namba M. 1996. Comparison of various methods of assaying the cytotoxic effects of ethanol on human hepatoblastoma cells (HUH-6 line). *Acta Medica Okayama* 50:151–156.
- Kostarelos K, Emfietzoglou D, Papakostas A, Yang WH, Ballangrud A, Sgouros G. 2004. Binding and interstitial penetration of liposomes within avascular tumor spheroids. *International Journal of Cancer* 112:713–721.
- Kross KW, Heimdal JH, Olsnes C, Olofsson J, Aarstad HJ. 2005. Head and neck squamous cell carcinoma spheroid- and monocyte spheroid-stimulated IL-6 and monocyte chemotactic protein-1 secretion are related to TNM stage, inflammatory state and tumor macrophage density. *Acta oto-laryngologica* 125:1097–1104.
- Kubota N, Matsui K, Sato S, Inada T. 1984. Radiation responses of HMV-I human malignant melanoma cells grown *in vitro* as multicellular spheroids. *Journal of Radiation Research* 25:215–224.
- Kunz-Schughart LA. 1999. Multicellular tumor spheroids: Intermediates between monolayer culture and *in vivo* tumor. *Cell Biology International* 23:157–161.
- Kunz-Schughart LA, Freyer JP. 2002. Phosphorous metabolites and steady-state energetics of transformed fibroblasts during three-dimensional growth. *American Journal of Physiology: Cell Physiology* 283:C1287–C1297.
- Kunz-Schughart LA, Mueller-Klieser W. 2000. Three-dimensional culture. In: *Animal cell culture* (3rd ed.; Ed. Masters JRW). Oxford University Press. pp. 123–148.
- Kunz-Schughart LA, Freyer JP, Hofstaedter F, Ebner R. 2004. The use of 3-D cultures for high-throughput screening: The multicellular spheroid model. *Journal of Biomolecular Screening* 9:273–285.
- Kunz-Schughart LA, Groebe K, Mueller-Klieser W. 1996. Three-dimensional cell culture induces novel proliferative and metabolic alterations associated with oncogenic transformation. *International Journal of Cancer* 66:578–586.
- Kunz-Schughart LA, Heyder P, Schroeder J, Knuechel R. 2001. A heterologous 3-D coculture model of breast tumor cells and fibroblasts to study tumor-associated fibroblast differentiation. *Experimental Cell Research* 266:74–86.
- Kunz-Schughart LA, Knuechel R. 2002. Tumor-associated-fibroblasts (part II): Functional impact on tumor tissue. *Histology and Histopathology* 17:623–637.
- Kunz-Schughart LA, Kreutz M, Knuechel R. 1998. Multicellular spheroids: A three-dimensional *in vitro* culture system to study tumour biology. *International Journal of Experimental Pathology* 79:1–23.
- Kunz-Schughart LA, Schroeder JA, Wondrak M, van RF, Lehle K, Hofstaedter F, Wheatley DN. 2006. Potential of fibroblasts to regulate the formation of three-dimensional vessel-like structures from endothelial cells *in vitro*. *American Journal of Physiology: Cell Physiology* 290: C1385–C1398.
- Kuwashima Y, Majima H, Okada S. 1988. Cure, cell killing, growth delay and fragmentation of X-irradiated human melanoma HMV-I multicellular spheroids. *International Journal of Radiation Biology* 54:91–104.
- Kuwashima Y, Yamada T, Saio M, Takami T. 1993. Formation and growth of multicellular spheroids in media containing low concentrations of agarose. *Cancer Letters* 71:31–33.
- Kwok CS, Cole SE, Liao SK. 1988. Uptake kinetics of monoclonal antibodies by human malignant melanoma multicellular spheroids. *Cancer Research* 48:1856–1863.
- Kwok CS, Crivici A, MacGregor WD, Unger MW. 1989. Optimization of radioimmunotherapy using human malignant melanoma multicell spheroids as a model. *Cancer Research* 49:3276–3281.
- Kwok TT, Sutherland RM. 1991. The influence of cell-cell contact on radiosensitivity of human squamous carcinoma cells. *Radiation Research* 126:52–57.
- Lambert B, De RL, De VF, Slegers G, De G, V, Van de WC, Thierens H. 2006. Assessment of supra-additive effects of cytotoxic drugs and low dose rate irradiation in an *in vitro* model for hepatocellular carcinoma. *Canadian Journal of Physiology and Pharmacology* 84:1021–1028.
- Lambert B, De RL, Slegers G, De GV, Dierckx RA, Thierens H. 2004. Screening for supra-additive effects of cytotoxic drugs and gamma irradiation in an *in vitro* model for hepatocellular carcinoma. *Canadian Journal of Physiology and Pharmacology* 82:146–152.

- Lange CS, Djordjevic B, Brock WA. 1992. The hybrid spheroid clonogenic assay for the intrinsic radio- and chemo-sensitivities of human tumors. *International Journal of Radiation Oncology, Biology, Physics* 24:511–517.
- Langmuir VK, McGann JK, Buchegger F, Sutherland RM. 1989. 131I-anticarcinoembryonic antigen therapy of LS174T human colon adenocarcinoma spheroids. *Cancer Research* 49:3401–3406.
- LaRue H, Parent-Vaugeois C, Bergeron A, Champetier S, Fradet Y. 1997. Influence of spatial configuration on the expression of carcinoembryonic antigen and mucin antigens in human bladder cancer. *International Journal of Cancer* 71:986–992.
- LaRue KE, Khalil M, Freyer JP. 2004. Microenvironmental regulation of proliferation in multicellular spheroids is mediated through differential expression of cyclin-dependent kinase inhibitors. *Cancer Research* 64:1621–1631.
- Lee GY, Kenny PA, Lee EH, Bissell MJ. 2007. Three-dimensional culture models of normal and malignant breast epithelial cells. *Nature Methods* 4:359–365.
- Lees RK, Sordat B, MacDonald HR. 1981. Multicellular tumor spheroids of human colon carcinoma origin. Kinetic analysis of infiltration and in situ destruction in a xenogeneic (murine) host. *Experimental Cell Biology* 49:207–219.
- Legallais C, David B, Dore E. 2001. Bioartificial livers (BAL): Current technological aspects and future developments. *Journal of Membrane Science* 181:81–95.
- Lieubeau-Teillet B, Rak J, Jothy S, Iliopoulos O, Kaelin W, Kerbel RS. 1998. von Hippel-Lindau gene-mediated growth suppression and induction of differentiation in renal cell carcinoma cells grown as multicellular tumor spheroids. *Cancer Research* 58:4957–4962.
- Lin RZ, Chou LF, Chien CC, Chang HY. 2006. Dynamic analysis of hepatoma spheroid formation: Roles of E-cadherin and beta1-integrin. *Cell and Tissue Research* 324:411–422.
- Lohr F, Wenz F, Haas S, Flentje M. 1995. Comparison of proliferating cell nuclear antigen (PCNA) staining and BrdUrd-labelling index under different proliferative conditions *in vitro* by flow cytometry. *Cell Proliferation* 28:93–104.
- Lottner C, Knuechel R, Bernhardt G, Brunner H. 2004. Distribution and subcellular localization of a water-soluble hematoporphyrin-platinum(II) complex in human bladder cancer cells. *Cancer Letters* 215:167–177.
- Lowthers EL, Richard CL, Blay J. 2003. Differential sensitivity to short-chain ceramide analogues of human intestinal carcinoma cells grown in tumor spheroids versus monolayer culture. *In Vitro Cellular & Developmental Biology – Animal* 39:340–342.
- Lund-Johansen M, Bjerkvig R, Rucklidge GJ. 1992. Tumor spheroids from monolayer culture. R Bjerkvig, editor. Boca Raton, USA: CRC Press. pp 4–18.
- Makhija S, Taylor DD, Gibb RK, Gercel-Taylor C. 1999. Taxol-induced bcl-2 phosphorylation in ovarian cancer cell monolayer and spheroids. *International Journal of Oncology* 14:515–521.
- Martin C, Walker J, Rothnie A, Callaghan R. 2003. The expression of P-glycoprotein does influence the distribution of novel fluorescent compounds in solid tumour models. *British Journal of Cancer* 89:1581–1589.
- Martiny-Baron G, Korff T, Schaffner F, Esser N, Eggstein S, Marme D, Augustin HG. 2004. Inhibition of tumor growth and angiogenesis by soluble EphB4. *Neoplasia* 6:248–257.
- Marusic M, Bajzer Z, Vuk-Pavlovic S, Freyer JP. 1994. Tumor growth *in vivo* and as multicellular spheroids compared by mathematical models. *Bulletin of Mathematical Biology* 56:617–631.
- McFadden R, Kwok CS. 1988. Mathematical model of simultaneous diffusion and binding of antitumor antibodies in multicellular human tumor spheroids. *Cancer Research* 48:4032–4037.
- McLeod EJ, Beischer AD, Hill JS, Kaye AH. 1997. Multicellular tumor spheroids grown from pancreatic carcinoma cell lines: Use as an orthotopic xenograft in athymic nude mice. *Pancreas* 14:237–248.
- Mellor HR, Ferguson DJ, Callaghan R. 2005. A model of quiescent tumour microregions for evaluating multicellular resistance to chemotherapeutic drugs. *British Journal of Cancer* 93:302–309.
- Mills L, Tellez C, Huang S, Baker C, McCarty M, Green L, Gudas JM, Feng X, Bar-Eli M. 2002. Fully human antibodies to MCAM/MUC18 inhibit tumor growth and metastasis of human melanoma. *Cancer Research* 62:5106–5114.
- Monga M, Sausville EA. 2002. Developmental therapeutics program at the NCI: Molecular target and drug discovery process. *Leukemia* 16:520–526.
- Monzani E, Facchetti F, Galmozzi E, Corsini E, Benetti A, Cavazzin C, Gritti A, Piccinini A, Porro D, Santinami M, Invernici G, Parati E, Alessandri G, La Porta CA. 2007. Melanoma contains CD133 and ABCG2 positive cells with enhanced tumorigenic potential. *European Journal of Cancer* 43:935–946.
- Mueller-Klieser W. 1997. Three-dimensional cell cultures: From molecular mechanisms to clinical applications. *American Journal of Physiology* 273:C1109–C1123.
- Mueller-Klieser W. 2000. Tumor biology and experimental therapeutics. *Critical Reviews in Oncology/Hematology* 36:123–139.
- Mueller-Klieser W. 1987. Multicellular spheroids. A review on cellular aggregates in cancer research. *Journal of Cancer Research and Clinical Oncology* 113:101–122.
- Mueller-Klieser W, Schreiber-Klais S, Walenta S, Kreuter MH. 2002. Bioactivity of well-defined green tea extracts in multicellular tumor spheroids. *International Journal of Oncology* 21:1307–1315.
- Muir CP, Adams MA, Graham CH. 2006. Nitric oxide attenuates resistance to doxorubicin in three-dimensional aggregates of human breast carcinoma cells. *Breast Cancer Research and Treatment* 96:169–176.
- Muller-Holzer E, Marth C, Zeimet AG, Hofstadter F, Daxenbichler G. 1996. Effects of sex- and glucocorticoid steroids on breast cancer cells grown as either multicellular tumor spheroids or monolayers. *Journal of Steroid Biochemistry and Molecular Biology* 58:13–19.
- Muralidhar RV, Jayachandran G, Singh P. 2001. Development of high-density cultivation systems by bioencapsulation. *Current Science* 81:263–269.
- Nakamura K, Hanibuchi M, Yano S, Tanaka Y, Fujino I, Inoue M, Takezawa T, Shitara K, Sone S, Hanai N. 1999. Apoptosis induction of human lung cancer cell line in multicellular heterospheroids with humanized antiganglioside GM2 monoclonal antibody. *Cancer Research* 59:5323–5330.
- Nakayama T. 1991. Effects of split dose irradiation on monolayer cells and multicellular tumor spheroid. *Nippon Igaku Hoshasen Gakkai Zasshi* 51:79–87.
- Nelson CM, Bissell MJ. 2005. Modeling dynamic reciprocity: Engineering three-dimensional culture models of breast architecture, function, and neoplastic transformation. *Seminars in Cancer Biology* 15:342–352.
- Newell DR. 2001. Flasks, fibres and flanks—pre-clinical tumour models for predicting clinical antitumour activity. *British Journal of Cancer* 84:1289–1290.
- Ng CE, Keng PC, Sutherland RM. 1987. Characterization of radiation sensitivity of human squamous carcinoma A431 cells. *British Journal of Cancer* 56:301–307.
- Nicholson KM, Bibby MC, Phillips RM. 1997. Influence of drug exposure parameters on the activity of paclitaxel in multicellular spheroids. *European Journal of Cancer* 33:1291–1298.

- Nishikawa K, Newman RA, Murray L, Khokhar AR, Rosenblum MG. 1990. Detection of cellular platinum using the monoclonal antibody 1C1. *Molecular Biotherapy* 2:235–241.
- Ochs RL, Fensterer J, Ohori NP, Wells A, Gabrin M, George LD, Kornblith P. 2003. Evidence for the isolation, growth, and characterization of malignant cells in primary cultures of human tumors. *In Vitro Cellular & Developmental Biology – Animal* 39:63–70.
- Offner FA, Bigalke I, Schiefer J, Wirtz HC, Klosterhalfen B, Feichtinger H, Kirkpatrick CJ. 1993. Interaction of human malignant melanoma tumor spheroids with endothelium and reconstituted basement membrane: Modulation by RGDS. *International Journal of Cancer* 54:506–512.
- Offner FA, Schiefer J, Wirtz HC, Bigalke I, Pavelka M, Hollweg G, Ensinger C, Klosterhalfen B, Mittermayer C, Kirkpatrick CJ. 1996. Tumour-cell-endothelial interactions: Free radicals are mediators of melanoma-induced endothelial cell damage. *Virchows Archiv* 428:99–106.
- Offner FA, Wirtz HC, Schiefer J, Bigalke I, Klosterhalfen B, Bittinger F, Mittermayer C, Kirkpatrick CJ. 1992. Interaction of human malignant melanoma (ST-ML-12) tumor spheroids with endothelial cell monolayers. Damage to endothelium by oxygen-derived free radicals. *American Journal of Pathology* 141:601–610.
- Olea N, Villalobos M, Ruiz de Almodovar JM, Pedraza V. 1992. MCF-7 breast cancer cells grown as multicellular spheroids *in vitro*: Effect of 17 beta-estradiol. *International Journal of Cancer* 50:112–117.
- Olive PL, Banath JP. 1997. Multicell spheroid response to drugs predicted with the comet assay. *Cancer Research* 57:5528–5533.
- Olive PL, Durand RE. 1994. Drug and radiation resistance in spheroids: Cell contact and kinetics. *Cancer and Metastasis Reviews* 13:121–138.
- Olive PL, Vikse CM, Banath JP. 1996. Use of the comet assay to identify cells sensitive to tirapazamine in multicell spheroids and tumors in mice. *Cancer Research* 56:4460–4463.
- Oloumi A, Lam W, Banath JP, Olive PL. 2002. Identification of genes differentially expressed in V79 cells grown as multicell spheroids. *International Journal of Radiation Biology* 78:483–492.
- Oloumi A, MacPhail SH, Johnston PJ, Banath JP, Olive PL. 2000. Changes in subcellular distribution of topoisomerase IIalpha correlate with etoposide resistance in multicell spheroids and xenograft tumors. *Cancer Research* 60:5747–5753.
- Olsnes C, Heimdal JH, Kross K, Olofsson J, Aarstad HJ. 2002. Mechanisms for monocyte activation in co-culture with autologous tumor spheroids. *Cell Immunol* 219:11–21.
- Olsnes C, Heimdal JH, Kross KW, Olofsson J, Aarstad HJ. 2005. Viable head and neck tumor spheroids stimulate *in vitro* autologous monocyte MCP-1 secretion through soluble substances and CD14/lectin-like receptors. *European Archives of Oto-Rhino-Laryngology* 262:953–960.
- Orlandi P, Barbara C, Bocci G, Fioravanti A, Di PA, Del TM, Danesi R. 2005. Idarubicin and idarubicinol effects on breast cancer multicellular spheroids. *Journal of Chemotherapy* 17:663–667.
- Oudar O. 2000. Spheroids: Relation between tumour and endothelial cells. *Critical Reviews in Oncology/Hematology* 36:99–106.
- Owen MR, Byrne HM, Lewis CE. 2004. Mathematical modelling of the use of macrophages as vehicles for drug delivery to hypoxic tumour sites. *Journal of Theoretical Biology* 226:377–391.
- Padron JM, van der Wilt CL, Smid K, Smitskamp-Wilms E, Backus HH, Pizao PE, Giaccone G, Peters GJ. 2000. The multilayered postconfluent cell culture as a model for drug screening. *Critical Reviews in Oncology/Hematology* 36:141–157.
- Parry RL, Chin TW, Epstein J, Hudson PL, Powell DM, Donahoe PK. 1992. Recombinant human mullerian inhibiting substance inhibits human ocular melanoma cell lines *in vitro* and *in vivo*. *Cancer Research* 52:1182–1186.
- Pelkonen O, Myllynen P, Taavitsainen P, Boobis AR, Watts P, Lake BG, Price RJ, Renwick AB, Gomez-Lechon MJ, Castell JV, Ingelman-Sundberg M, Hiderstrand M, Guillouzo A, Corcos L, Goldfarb PS, Lewis DF. 2001. Carbamazepine: A 'blind' assessment of CYP-associated metabolism and interactions in human liver-derived *in vitro* systems. *Xenobiotica* 31:321–343.
- Persaud R, Zhou H, Baker SE, Hei TK, Hall EJ. 2005. Assessment of low linear energy transfer radiation-induced bystander mutagenesis in a three-dimensional culture model. *Cancer Research* 65:9876–9882.
- Pervez S, Kirkland SC, Epenetos AA, Mooi WJ, Evans DJ, Krausz T. 1989. Effect of polarity and differentiation on antibody localization in multicellular tumour spheroid and xenograft models and its potential importance for *in vivo* immunotargeting. *International Journal of Cancer* 44:940–947.
- Porwol T, Merten E, Opitz N, Acker H. 1996. Three-dimensional imaging of rhodamine 123 fluorescence distribution in human melanoma cells by means of confocal laser scanning microscopy. *Acta Anatomica* 157:116–125.
- Pourreau-Schneider N, Berthois Y, Mitre H, Charpin C, Jacquemier J, Martin PM. 1984. Estrogen response of MCF-7 cells grown on diverse substrates and in suspension culture: Promotion of morphological heterogeneity, modulation of progesterone receptor induction; cell-substrate interactions on collagen gels. *Journal of Steroid Biochemistry* 21:763–771.
- Rae JM, Ramus SJ, Waltham M, Armes JE, Campbell IG, Clarke R, Barndt RJ, Johnson MD, Thompson EW. 2004. Common origins of MDA-MB-435 cells from various sources with those shown to have melanoma properties. *Clinical and Experimental Metastasis* 21:543–552.
- Ricci-Vitiani L, Lombardi DG, Pilozzi E, Biffoni M, Todaro M, Peschle C, De MR. 2007. Identification and expansion of human colon-cancer-initiating cells. *Nature* 445:111–115.
- Rieber M, Rieber MS. 2006. Signalling responses linked to betulinic acid-induced apoptosis are antagonized by MEK inhibitor U0126 in adherent or 3D spheroid melanoma irrespective of p53 status. *International Journal of Cancer* 118:1135–1143.
- Rofstad EK, Eide K, Skoyum R, Hystad ME, Lyng H. 1996. Apoptosis, energy metabolism, and fraction of radiobiologically hypoxic cells: A study of human melanoma multicellular spheroids. *International Journal of Radiation Biology* 70:241–249.
- Ronen SM, Stier A, Degani H. 1990. NMR studies of the lipid metabolism of T47D human breast cancer spheroids. *FEBS Letters* 266:147–149.
- Sacks PG, Oke V, Vasey T, Lotan R. 1989. Retinoic acid inhibition of a head and neck multicellular tumor spheroid model. *Head and Neck* 11:219–225.
- Santini MT, Rainaldi G. 1999. Three-dimensional spheroid model in tumor biology. *Pathobiology* 67:148–157.
- Sausville EA. 2000. Overview. Cancer drug discovery: Pathway promise or covalent certainty for drug effect – quo vadis? *Current Opinion in Investigational Drugs* 1:511–513.
- Schuster U, Buttner R, Hofstadter F, Knuchel R. 1994. A heterologous *in vitro* coculture system to study interaction between human bladder cancer cells and fibroblasts. *Journal of Urology* 151:1707–1711.

- Schwachofer JH, Crooijmans RP, van GJ, Hoogenhout J, Jerusalem CR, Kal HB. 1989. Sublethal damage repair and radiosensitivity of human squamous cell carcinoma cells grown with different culture techniques. *British Journal of Radiology* 62:1084–1088.
- Schwachofer JH, Crooijmans RP, Hoogenhout J, Jerusalem CR, Kal HB, Theeuwes AG. 1990. Radiosensitivity of human melanoma spheroids influenced by growth rate. *International Journal of Radiation Oncology, Biology, Physics* 19:1191–1197.
- Schwachofer JH, Crooijmans RP, Hoogenhout J, Kal HB, Theeuwes AG. 1991. Sublethal damage repair in two radio-resistant human tumor cell lines irradiated as multicellular spheroids. *Tumour biology* 12:207–216.
- Seidl P, Huettinger R, Knuechel R, Kunz-Schughart LA. 2002. Three-dimensional fibroblast-tumor cell interaction causes downregulation of RACK1 mRNA expression in breast cancer cells *in vitro*. *International Journal of Cancer* 102:129–136.
- Selden C, Khalil M, Hodgson H. 2000. Three dimensional culture upregulates extracellular matrix protein expression in human liver cell lines – a step towards mimicking the liver *in vivo*? *International Journal of Artificial Organs* 23:774–781.
- Shellman YG, Park YL, Marr DG, Casper K, Xu Y, Fujita M, Swerlick R, Norris DA. 2003. Release of vascular endothelial growth factor from a human melanoma cell line, WM35, is induced by hypoxia but not ultraviolet radiation and is potentiated by activated Ras mutation. *Journal of Investigative Dermatology* 121:910–917.
- Siebert A, Denkert C, Leclerc A, Hauptmann S. 1999. Suppression of the reactive oxygen intermediates production of human macrophages by colorectal adenocarcinoma cell lines. *Immunology* 98:551–556.
- Silzle T, Kreutz M, Dobler MA, Brockhoff G, Knuechel R, Kunz-Schughart LA. 2003. Tumor-associated fibroblasts recruit blood monocytes into tumor tissue. *European Journal of Immunology* 33:1311–1320.
- Simiantonaki N, Jayasinghe C, Kirkpatrick CJ. 2002. Differential endothelial CAM-expression after stimulation with supernatants of LPS- and cytokine-stimulated HT-29 and ST-ML-12 tumor cells growing as monolayer cultures and multicellular spheroids. *Anticancer Research* 22:2641–2649.
- Sommers GM, Alfieri AA. 1998. Multimodality therapy: Radiation and continuous concomitant cis-platinum and PKC inhibition in a cervical carcinoma model. *Cancer Investigation* 16:462–470.
- Song H, David O, Clejan S, Giordano CL, Pappas-Lebeau H, Xu L, O'Connor KC. 2004. Spatial composition of prostate cancer spheroids in mixed and static cultures. *Tissue Engineering* 10:1266–1276.
- Sonoda T, Kobayashi H, Kaku T, Hirakawa T, Nakano H. 2003. Expression of angiogenesis factors in monolayer culture, multicellular spheroid and *in vivo* transplanted tumor by human ovarian cancer cell lines. *Cancer Letters* 196:229–237.
- Soranzo C, Della TG, Ingrosso A. 1986. Formation, growth and morphology of multicellular tumor spheroids from a human colon carcinoma cell line (LoVo). *Tumori* 72:459–467.
- Soranzo C, Ingrosso A. 1988. A comparative study of the effects of anthracycline derivatives on a human adenocarcinoma cell line (LoVo) grown as a monolayer and as spheroids. *Anticancer Research* 8:369–373.
- Spoettl T, Hausmann M, Herlyn M, Gunckel M, Dirmeier A, Falk W, Herfarth H, Schoelmerich J, Rogler G. 2006. Monocyte chemoattractant protein-1 (MCP-1) inhibits the intestinal-like differentiation of monocytes. *Clinical & Experimental Immunology* 145:190–199.
- Spoettl T, Hausmann M, Menzel K, Piberger H, Herfarth H, Schoelmerich J, Bataille F, Rogler G. 2007. Role of soluble factors and three-dimensional culture in *in vitro* differentiation of intestinal macrophages. *World Journal of Gastroenterology* 13:1032–1041.
- Strasberg RM, Anzellotti A, Sanchez-Delgado RA, Rieber M. 2004. Tumor apoptosis induced by ruthenium(II)-ketoconazole is enhanced in nonsusceptible carcinoma by monoclonal antibody to EGF receptor. *International Journal of Cancer* 112:376–384.
- Sturgis EM, Sacks PG, Masui H, Mendelsohn J, Schantz SP. 1994. Effects of antiepidermal growth factor receptor antibody 528 on the proliferation and differentiation of head and neck cancer. *Otolaryngology – Head and Neck Surgery* 111:633–643.
- Stuschke M, Budach V, Budach W, Feldmann HJ, Sack H. 1992. Radioresponsiveness, sublethal damage repair and stem cell rate in spheroids from three human tumor lines: Comparison with xenograft data. *International Journal of Radiation Oncology, Biology, Physics* 24:119–126.
- Stuschke M, Budach V, Stuben G, Streffer C, Sack H. 1995. Heterogeneity in the fractionation sensitivities of human tumor cell lines: Studies in a three-dimensional model system. *International Journal of Radiation Oncology, Biology, Physics* 32:395–408.
- Sutherland RM. 1988. Cell and environment interactions in tumor microregions: The multicell spheroid model. *Science* 240:177–184.
- Sutherland RM, Inch WR, McCredie JA. 1971a. Phytohemagglutinin (PHA)-induced transformation of lymphocytes from patients with cancer. *Cancer* 27:574–578.
- Sutherland RM, Inch WR, McCredie JA, Kruuv J. 1970. A multi-component radiation survival curve using an *in vitro* tumour model. *International Journal of Radiation Biology and Related Studies in Physics, Chemistry, and Medicine* 18:491–495.
- Sutherland RM, McCredie JA, Inch WR. 1971b. Growth of multicell spheroids in tissue culture as a model of nodular carcinomas. *Journal of the National Cancer Institute* 46:113–120.
- Sutherland RM, Sordat B, Bamat J, Gabbert H, Bourrat B, Mueller-Klieser W. 1986. Oxygenation and differentiation in multicellular spheroids of human colon carcinoma. *Cancer Research* 46:5320–5329.
- Sutton JE, Roos IA, Hillcoat BL. 1982. Combined actions of 5-fluorouracil and 1-(2-chloroethyl)-3-(4-methylcyclohexyl)-1-nitrosourea on human colonic carcinoma cells *in vitro*. *Cancer Research* 42:5172–5175.
- Tabatabai M, Williams DK, Bursac Z. 2005. Hyperbolic growth models: theory and application. *Theoretical Biology and Medical Modelling* 2:14–26.
- Talts U, Kuhn U, Roos G, Rauch U. 2000. Modulation of extracellular matrix adhesiveness by neurocan and identification of its molecular basis. *Experimental Cell Research* 259:378–388.
- Tannock IF, Kopelyan I. 1986. Influence of glucose concentration on growth and formation of necrosis in spheroids derived from a human bladder cancer cell line. *Cancer Research* 46:3105–3110.
- Teutsch HF, Goellner A, Mueller-Klieser W. 1995. Glucose levels and succinate and lactate dehydrogenase activity in EMT6/Ro tumor spheroids. *European Journal of Cell Biology* 66:302–307.
- Thielecke H, Mack A, Robitzki A. 2001. A multicellular spheroid-based sensor for anti-cancer therapeutics. *Biosensors and Bioelectronics* 16:261–269.
- Thomsen LL, Baguley BC, Wilson WR. 1992. Nitric oxide: Its production in host-cell-infiltrated EMT6 spheroids and its role in tumour cell killing by flavone-8-acetic acid and 5,6-dimethylxanthene-4-acetic acid. *Cancer Chemotherapy and Pharmacology* 31:151–155.

- Timmins NE, Dietmair S, Nielsen LK. 2004. Hanging-drop multicellular spheroids as a model of tumour angiogenesis. *Angiogenesis* 7:97–103.
- Tofilon PJ, Arundel CM, Deen DF. 1987. Response to BCNU of spheroids grown from mixtures of drug-sensitive and drug-resistant cells. *Cancer Chemotherapy and Pharmacology* 20:89–95.
- Tonary AM, Pezacki JP. 2006. Simultaneous quantitative measurement of luciferase reporter activity and cell number in two- and three-dimensional cultures of hepatitis C virus replicons. *Analytical Biochemistry* 350:239–248.
- Tong JZ, De Lagausie P, Furlan V, Cresteil T, Bernard O, Alvarez F. 1992. Long-term culture of adult rat hepatocyte spheroids. *Experimental Cell Research* 200:326–332.
- Torisawa YS, Takagi A, Shiku H, Yasukawa T, Matsue T. 2005. A multicellular spheroid-based drug sensitivity test by scanning electrochemical microscopy. *Oncology Reports* 13:1107–1112.
- Vamvakidou AP, Mondrinos MJ, Petushi SP, Garcia FU, Lelkes PI, Tozeren A. 2007. Heterogeneous breast tumoroids: An *in vitro* assay for investigating cellular heterogeneity and drug delivery. *Journal of Biomolecular Screening* 12:13–20.
- Vermeulen S, Van MV, Van HL, Van RF, Bracke M, Mareel M. 1996. Regulation of the invasion suppressor function of the cadherin/catenin complex. *Pathology, Research and Practice* 192:694–707.
- Voskoglou-Nomikos T, Pater JL, Seymour L. 2003. Clinical predictive value of the *in vitro* cell line, human xenograft, and mouse allograft preclinical cancer models. *Clinical Cancer Research* 9:4227–4239.
- Walenta S, Dotsch J, Mueller-Klieser W. 1990. ATP concentrations in multicellular tumor spheroids assessed by single photon imaging and quantitative bioluminescence. *European Journal of Cell Biology* 52:389–393.
- Walker GM, Beebe DJ. 2002. A passive pump method for microfluidic devices. *Lab on a chip* 2:131–134.
- Walker J, Martin C, Callaghan R. 2004. Inhibition of P-glycoprotein function by XR9576 in a solid tumour model can restore anticancer drug efficacy. *European Journal of Cancer* 40:594–605.
- Warrick J, Meyvantsson I, Ju J, Beebe DJ. 2007. High-throughput microfluidics: Improved sample treatment and washing over standard wells. *Lab on a chip* 7:316–321.
- Wartenberg M, Donmez F, Ling FC, Acker H, Hescheler J, Sauer H. 2001. Tumor-induced angiogenesis studied in confrontation cultures of multicellular tumor spheroids and embryoid bodies grown from pluripotent embryonic stem cells. *FASEB Journal* 15:995–1005.
- Wartenberg M, Fischer K, Hescheler J, Sauer H. 2002. Modulation of intrinsic P-glycoprotein expression in multicellular prostate tumor spheroids by cell cycle inhibitors. *Biochimica et Biophysica Acta* 1589:49–62.
- Wartenberg M, Schallenberg M, Hescheler J, Sauer H. 2003. Reactive oxygen species-mediated regulation of eNOS and iNOS expression in multicellular prostate tumor spheroids. *International Journal of Cancer* 104:274–282.
- Wartenberg M, Finkensieper A, Hescheler J, Sauer H. 2006. Confrontation cultures of embryonic stem cells with multicellular tumor spheroids to study tumor-induced angiogenesis. *Methods in Molecular Biology* 331:313–328.
- Yuhás JM, Blake S, Weichselbaum RR. 1984. Quantitation of the response of human tumor spheroids to daily radiation exposures. *International Journal of Radiation Oncology, Biology, Physics* 10:2323–2327.
- Zhang X, Wang W, Yu W, Xie Y, Zhang X, Zhang Y, Ma X. 2005. Development of an *in vitro* multicellular tumor spheroid model using microencapsulation and its application in anticancer drug screening and testing. *Biotechnology Progress* 1289–1296.