


**BD**

Helping all people  
live healthy lives

# BD Biosciences Cell Growth and Differentiation



Supplier Partnerships for Customer Solutions



# Table of Contents

Human Embryonic Stem Cells .....4



Endothelial Cells .....6

Hepatocytes ..... 10

Neuronal Cells..... 12

Epithelial Cells..... 14

Tumor Cells..... 17



# Cell Growth and Differentiation

## Enhancing Cell Culture and Accelerating Discovery



The development and normal functioning of cells depends on interactions with molecules in their microenvironment. The major classes of molecules that regulate cellular development and function include growth and differentiation factors, cell adhesion molecules, and the components of the extracellular matrix (ECM).

The ECM, composed of a number of different macromolecules, influences behavior, (adherence, spreading, differentiation, and migration) and the pattern of gene expression of the cells in contact with it. To create physiologically relevant *in vitro* models that support normal cell growth and function, the components of the *in vivo* environment must be incorporated. Use of ECM proteins as coating for tissue culture surfaces permits the development of cell type specific model systems which closely mimic *in vivo* conditions.

Recognizing the increasingly important role the ECM plays in the regulation of fundamental cellular processes BD Biosciences offers a wide range of extracellular matrix proteins and attachment factors for researchers to incorporate into their cell culture systems. For over 20 years, BD has provided the research market with a wide variety of purified proteins. We were the first to offer a unique line of tissue culture vessels coated with a variety of ECM proteins and attachment factors: BD BioCoat Cellware. BD's extensive experience in protein purification, along with rigorous quality assurance testing guarantees high-quality, consistent products.

At BD Biosciences - Discovery Labware we are committed to enhancing cell culture and accelerating discovery worldwide through dedicated customer service, innovative product solutions, and technical expertise. We strive to make cell culture research more efficient and convenient for researchers by offering outstanding quality, consistency, and value.

### Commitment to Quality

We understand the importance of lot-to-lot consistency and the need for reproducible results. Through proprietary manufacturing technology, validated procedures, strict compliance with established protocols, and exacting quality control, we are able to assure the biological performance of our products as well as consistency from lot-to-lot.

### Delivering Choice

The optimal surface for cell attachment, proliferation, and differentiation is dependent on the particular cell type. BD Falcon™, BD BioCoat™, and BD™ ECM proteins provide diverse options for a variety of cells, including but not limited to commonly used cell lines such as HEK-293, primary neuronal cells, and three-dimensional culture.

### Technical Expertise

Our scientists routinely study a broad range of cells to better understand their cellular function. Our team of highly skilled and dedicated Technical Support Specialists are available to assist you in protocol development and troubleshooting.

### Customizable Solutions

We offer a custom product service to meet the unique needs of our customers. Our custom capabilities range from special package sizes and sterilization needs to barcoding and custom coating. Through our custom coating services, we will apply the coating of your choice on BD and alternative cultureware products. If you are not sure which coating you need, our Technical Support Specialists can recommend surfaces for your cell type.

# Cell Culture Surfaces

BD Biosciences offers a wide variety of surface chemistries and attachment factors appropriate for a broad range of applications. The surface of our BD Falcon™ Cultureware is rendered permanently hydrophilic via a unique vacuum-gas plasma tissue culture treatment process. This treatment process is produced in a closed, highly controlled environment ensuring a consistent treatment surface. BD Primaria™ and BD BioCoat™ surface options are ideal for enhanced cell attachment and growth of a variety of primary cells, stem cells, and transformed cell lines in serum-free or serum-containing cultures. A non-treated surface is also available for suspension or non-adherent cell culture and may also be used to study cell-cell or cell-protein interactions in an *in vitro* system.

## BD Falcon Non-treated Polystyrene

- Hydrophobic surface with low to moderate binding properties. Ideal for cell-cell or cell-protein studies.

## BD Falcon Tissue Culture-treated (TC)

- Hydrophilic surface enhances cell attachment, spreading, and cell growth by binding serum proteins to the surface. Highly controlled vacuum-gas plasma treatment creates negatively charged carboxyl groups on the polystyrene surface.
- Tested for confluency of MRC-5 cells and sterilized by gamma-irradiation.

## BD Primaria

- Supports neuronal, primary, endothelial, and tumor cells which may have difficulty attaching to or differentiate poorly on traditional TC surfaces. This surface has a unique mixture of negative and nitrogen containing positive functional groups on the polystyrene surface.
- The surface consistency of each lot is confirmed by electron spectroscopy chemical analysis (ESCA).

## BD BioCoat Poly-D-Lysine (PDL)

- Pre-coated with PDL, which promotes cell attachment of transfected and primary cells (e.g., neuronal).
- Tested for the ability to promote firm attachment of rat cerebellar granule (RCG) cells.
- Stable for six months from date of shipment at 4-30°C. Coverslips, CultureSlides, and Coverslip-Bottom Dishes stable for at least three months from date of shipment at 4°C.

## BD BioCoat Collagen I

- Pre-coated with Collagen I, derived from rat tail tendon.
- Tested for the ability to promote attachment and spreading of HT-1080 human fibrosarcoma cells.
- Stable for at least six months from date of shipment when stored at 4-30°C under dry conditions. Coverslips and CultureSlides are stable for at least three months from date of shipment when stored at 2-8°C.

## BD BioCoat Collagen IV

- Pre-coated with Collagen IV. Useful as a substrate for nerve, epithelial, endothelial, and muscle cells.
- Tested for the ability to promote attachment and spreading of PC12 rat pheochromocytoma cells or to initiate differentiation (neurite outgrowth) of NG-108 rat glioma/mouse neuroblastoma cells.
- Stable for at least three months at 2-8°C. Do not freeze.

## BD BioCoat Gelatin

- Pre-coated with Gelatin, which is commonly used for culture of vascular endothelial cells and F9 teratocarcinoma cells.
- Tested to promote proliferation of Human Umbilical Vein Endothelial Cells (HUVEC).
- Stable for at least three months from date of shipment when stored at 4-30°C under dry conditions.

## BD BioCoat Fibronectin

- Pre-coated with Human Fibronectin (HFN), which promotes cell attachment through integrin binding. HFN promotes cellular migration during wound healing and improves survival of primary cells.
- Tested to promote attachment and spreading of BHK-1 hamster kidney cells.
- Stable for at least three months at 2-8°C. Do not freeze.

## BD BioCoat Laminin

- Pre-coated with Laminin, a major component of the basement membrane used as a substrate to culture and maintain differentiated functions of a variety of cells including neuroblastoma cells and breast cancer cell lines.
- Tested for the ability to initiate neurite outgrowth of NG-108 rat glioma/mouse neuroblastoma cells.
- Stable for at least three months at 2-8°C. Do not freeze.

## BD BioCoat Laminin/Fibronectin

- Pre-coated with a combination of ECMs, which provide superior attachment and growth of glial precursor cells.
- Tested for receptor agonist induced changes in intracellular calcium-using FLUO-3 in primary rat cortical enriched cultures.
- Stable for at least three months at 2-8°C. Do not freeze.

## BD BioCoat Poly-D-Lysine/Laminin (PDL/Laminin)

- Pre-coated with a combination of ECMs, which supports neuronal differentiation of human and mouse stem cells.
- Tested for the ability to promote neurite outgrowth with primary rat cerebellar granule (RCG) cells and NG-108 rat glioma/mouse neuroblastoma cells.
- Stable for at least 3 months at 2-8°C. Do not freeze.

## BD BioCoat Poly-L-Ornithine/Laminin (PLO/Laminin)

- Pre-coated with a combination of ECMs, which support growth of neuroblastoma cells and differentiation of N2a and ScN3a cells.
- Tested for the ability to promote neurite outgrowth with primary rat cerebellar granule (RCG) cells and NG-108 rat glioma/mouse neuroblastoma cells.
- Stable for at least three months at 2-8°C. Do not freeze.

## BD BioCoat Matrigel™ Matrix

- Pre-coated with solubilized basement membrane matrix extracted from Engelbreth-Holm-Swarm (EHS) mouse sarcoma. Rich in ECM proteins, especially laminin, collagen IV, heparin sulphate proteoglycans, and entactin.
- Tested for the ability to promote neurite outgrowth from chick dorsal root ganglia in the absence of Nerve Growth Factor (NGF).
- Stable for at least three months at -20°C. Keep frozen until use.



hES CELLS	HEPATOCYTES	ENDOTHELIAL CELLS	NEURONAL CELLS	EPITHELIAL CELLS	TUMOR CELLS	BD BIOSCIENCES - DISCOVERY LABWARE PRODUCT	
■	■	■	■	■	■	BD Matrigel™ Matrix	Cell Culture Reagents
■					■	Laminin/Entactin Complex High Concentration	
	■	■	■	■	■	Collagen I	
			■		■	Fibronectin	
■			■		■	Laminin	
			■		■	Poly-D-Lysine	
	■		■		■	BD™ PuraMatrix™ Peptide Hydrogel	
■					■	bFGF	
	■					Hepatocyte Culture Media	
	■	■	■	■	■	ITS	
		■				Vascular Endothelial Growth Factor (VEGF)	
		■				Endothelial Cell Growth Supplement (ECGS)	
			■			Nerve Growth Factor (NGF)	
			■			Endothelial Growth Factor (EGF)	
				■		Enterocyte Differentiation Medium	
	■					Intestinal Epithelium Differentiation Media Pack	
				■		MITO+ Serum Extender	
				■		Seeding Basal Medium	
		■				HUVEC-2	
	■	■	■	■	■	Calcein AM	
	■	■	■	■	■	DilC <sub>12</sub> (3)	
■	■	■	■	■	■	Dispase	
■	■	■	■	■	■	Cell Recovery Solution	
■						BD BioCoat™ Matrigel™ Matrix Plates for Embryonic Stem Cell Culture	Cell Culture Tools
	■	■		■	■	BD BioCoat Collagen I Cellware	
	■					BD BioCoat Matrigel Matrix - for hepatocytes	
			■	■	■	BD BioCoat Poly-Lysine Cellware	
■			■		■	BD BioCoat Laminin Cellware	
			■		■	BD BioCoat Poly-L-Ornithine/Laminin Cellware	
			■		■	BD BioCoat Poly-D-Lysine/Laminin Cellware	
■	■	■	■	■	■	BD Falcon™ Tissue Culture-treated Flasks	
■	■	■	■	■	■	BD Falcon CultureSlides	
■	■	■	■	■	■	BD Falcon 96-well Imaging Plates	
■	■	■	■	■	■	BD Primaria™ Cultureware	
	■					Cholyl-lysyl-Fluorescein (CLF)	Hepato-cytes
	■					BD Gentest™ Hepatocytes	
	■					Hepatocyte Differentiation Environment	
		■				Endothelial Cell Growth Environment	Cell Environments
		■				BD BioCoat Angiogenesis System: Endothelial Cell Tube Formation	
		■				BD BioCoat Angiogenesis System: Endothelial Cell Migration	
		■				BD BioCoat Angiogenesis System: Endothelial Cell Invasion	
				■		BD BioCoat Intestinal Epithelium Differentiation Environment	
				■		BD BioCoat HTS Caco-2 Assay System	
					■	BD BioCoat Matrigel Invasion Chamber	
					■	BD BioCoat Tumor Invasion System	
				■		BD BioCoat Fibrillar Collagen Cell Culture Inserts	Membrane Insert Systems
				■		BD BioCoat Fibrillar Collagen 24-Multiwell Insert System	
■	■	■	■	■	■	BD BioCoat and BD Falcon Inserts	

For guideline use only. This is not a complete list of all applications for these products.




## Human Embryonic Stem Cells

Human embryonic stem (hES) cells are pluripotent cells derived from the inner cell mass of a blastocyst. These cells can either self-renew, thereby maintaining their pluripotency, or differentiate into all three germ layers depending upon the culture conditions. Induced pluripotent stem (iPS) cells, which are similar in potential to hES cells, have been generated by infecting adult cells. iPS cells, like hES cells, can form all three germ layers as well as self-renew. Tremendous hope is associated with the potential application of hES and iPS cells in cell therapy and regenerative medicine because of their ability to differentiate into multiple, clinically useful cell types. Defined culture conditions are essential to realizing the potential of hES and iPS cells.




A culture environment for hES cells consisting of both a serum-free, defined medium, and a cell culture surface specifically qualified for hES cells saves researchers time and resources normally spent qualifying reagents. BD Biosciences, StemCell Technologies, and the WiCell™ Research Institute have established a strategic collaboration to develop optimized, feeder-independent cell culture environments for hES cell research, including serum-free, defined media and qualified surfaces. BD Matrigel™ Matrix, coupled with a variety of culture media, has been widely accepted as an alternative substrate to feeder-dependent culture of hES cells<sup>1-4</sup>, and BD Matrigel Matrix has been used to culture iPS cells<sup>5-6</sup>. BD Matrigel Matrix is a reconstituted basement membrane isolated from the Engelbreth-Holm-Swarm (EHS) mouse sarcoma.

FIGURE 1 • HUMAN EMBRYONIC STEM CELLS CULTURED ON BD MATRIGEL hESC-QUALIFIED MATRIX

A.



B.



1A. Phase contrast images of H9 colonies grown on mouse embryonic fibroblast (MEF) feeder layer in hES media (left), BD Matrigel hESC-qualified Matrix in MEF-conditioned media (middle), or mTeSR1™ maintenance media (right). Images were taken at 4x magnification.


1B. Flow cytometry analysis of H9 cells cultured on BD Matrigel hESC-qualified Matrix coated surface in mTeSR1 maintenance media. Cells were probed with the following antibodies: Tra-1-60 PE (Cat. No. 560193), Tra-1-81 PE (Cat. No. 560161), SSEA-4 PE (Cat. No. 560128) and Oct3/4 PE (Cat. No. 560186) compared to isotype control. Percent positive is indicated. Cells were run on a BD FACSCalibur™ system and the data was analyzed with CellQuest software.

StemCell Technologies has commercially developed and optimized WiCell™ Research Institute's mTeSR™1 medium formulation to standardize feeder-independent hES cell culture. mTeSR1 is complete, defined and serum-free, and has been designed to maintain and expand hES cells in an undifferentiated state when used with BD Matrigel™ hESC-qualified Matrix as a substrate (**Figure 1**).


An alternative surface for hES cell culture is BD™ Laminin/Entactin Complex High Concentration (**Figure 2**). BD Laminin/Entactin Complex High Concentration, with a purity greater than or equal to 90%, is a more defined surface that can support undifferentiated hES cell growth. Unlike BD Matrigel hESC-qualified Matrix, this surface is not specifically qualified for maintenance of undifferentiated hES cells.

**FIGURE 2 • BD LAMININ/ENTACTIN COMPLEX HIGH CONCENTRATION FOR HUMAN EMBRYONIC STEM CELL CULTURE**



**A. BD Matrigel hESC-qualified Matrix**




**BD Laminin/Entactin Complex High Concentration**



**B.**



**C.**



2A. Phase contrast images of H9 cells grown on BD Matrigel hESC-qualified Matrix (left) and BD Laminin/Entactin Complex High Concentration (right) in mTeSR1 maintenance media. Images were taken at 4x magnification.

2B. Flow cytometry analysis of H9 cells cultured on BD Laminin/Entactin Complex High Concentration (red line) and BD Matrigel hESC-qualified Matrix coated surface (green line) in mTeSR1 maintenance media. Cells were probed with the following antibodies: SSEA-4 PE (Cat. No. 560128) and Oct3/4 PE (Cat. No. 560186) compared to isotype control (black line). Cells were run on a BD FACSCalibur™ system and the data was analyzed with CellQuest software. Both surfaces supported undifferentiated expansion of hESC, H9.

2C. G banding chromosome analysis. Karyotype analysis of H9 cells grown on BD Laminin/Entactin Complex High Concentration in mTeSR1 media for 26 passages. Cells maintained normal karyotype under these culture conditions.

## Tools for Human Embryonic Stem Cell Culture

Cat. No.	Description	Qty.
----------	-------------	------

### Cell Culture Reagents

#### Extracellular Matrix Proteins

80085-668	Laminin/Entactin Complex High Concentration	10.5 mg
-----------	---	---------

#### Cytokines and Media Additives

47743-574	bFGF, human recombinant	10 µg
-----------	-------------------------	-------

#### Cell Recovery Reagents

47743-724	Dispase	100 ml
47743-696	Cell Recovery Solution	100 ml

### Cell Culture Tools

#### BD BioCoat™ Matrigel™ Matrix Plates for Embryonic Stem Cell Culture

89022-350	6-well Plates	5
-----------	---------------	---

#### BD Falcon™ Multiwell Cell Culture Plates

62406-161	6-well Flat-bottom with lid, Tissue Culture-treated	1
-----------	---	---

For a complete product listing, see page 19.



### DID YOU KNOW?

- BD Biosciences offers a full range of pipets and tubes. Please contact your sales representative for more information.






## Endothelial Cells

Endothelial cells are a specialized type of epithelial cell which forms the inner layer of blood vessels. These cells play a key role in angiogenesis, the development of new blood vessels from pre-existing vessels. Angiogenesis is a multi-step process that is important for both physiological and pathological development. During angiogenesis, endothelial cells are activated and express matrix metalloproteinases (MMPs), which degrade the vascular basement membrane. In response to environmental cues, endothelial cells secrete MMPs and then invade through the basement membrane to form new capillary networks.

Endothelial cells are tested in a variety of assays for functions that contribute to the angiogenesis process. Collagen I coated surfaces are suitable for culturing endothelial cells such as fetal bovine heart endothelial cells (FBHECs) and human umbilical vein endothelial cells (HUVECs) (**Figure 3**). *In vitro* assays of endothelial cell function include cell migration<sup>7</sup>, invasion<sup>8</sup>, and tubule formation<sup>9-15</sup>. Both the BD BioCoat™ Angiogenesis System: Endothelial Cell Invasion and the BD BioCoat Angiogenesis System: Endothelial Cell Migration allow for rapid data collection without multiple handling steps. These quantitative assays utilize BD FluoroBlok™ microporous polyethylene terephthalate (PET) membranes (3 µm pore size) which effectively block the fluorescence signal from labeled cells that have not invaded or migrated through the membrane, respectively, thereby allowing the selective detection of cells that reside on the underside of the membrane (**Figure 4**). To perform fluorescence detection, cells may be pre-labeled or post-labeled with a fluorescent dye (**Figure 5**). The pre-labeling technique enables real-time kinetic measurements of cell migration or invasion. Endothelial cells must be able to migrate and enzymatically degrade the basement membrane in order for angiogenesis to occur. The wells of BD BioCoat Angiogenesis System: Endothelial Cell Invasion are evenly coated with BD Matrigel™ Matrix, which allows researchers to examine the ability of endothelial cells to invade through reconstituted basement membrane in response to chemoattractants, such as VEGF, in the presence or absence of anti-angiogenic agents (**Figure 6**).

FIGURE 3 • EFFECTS OF BD BIOCOAT ENDOTHELIAL CELL GROWTH ENVIRONMENT ON HUVEC




BD BioCoat Endothelial Cell Growth Environment utilizes BD BioCoat Collagen I Cellware and BD™ Endothelial Cell Culture Medium to enhance endothelial attachment and proliferation. HUVECs grown for five days using the BD BioCoat Endothelial Cell Growth Environment form a confluent monolayer and show numerous mitotic cells (A). HUVECs grown for five days in basal medium containing 10% FBS on tissue culture-treated plastic show sparse growth (B).

### DID YOU KNOW?

- The use of BD™ Cell Recovery Solution or BD Dispase is necessary to recover cells cultured on BD Matrigel™ Matrix.




FIGURE 4 • LABELING CELLS POST-INVASION WITH CALCEIN AM





A fluorescence plate reader quantifies cells post-invasion by measuring fluorescence which correlates to cell number. Cells on top of the BD FluoroBlok™ membrane are not detected by a bottom-reading fluorometer.

FIGURE 5 • LABELING METHODS FOR ENDPOINT OR REAL-TIME KINETIC MIGRATION AND INVASION ASSAYS



BD FluoroBlok Inserts can be used for endpoint or real-time kinetic assays. For endpoint assays, the cell migration or invasion assay is performed with unlabeled cells. At the end of the assay the cells are labeled with a fluorescent dye, such as BD™ Calcein AM, and the data is collected using a bottom reading fluorescent plate reader. For real-time kinetic assays, the cells are pre-labeled with a fluorescent dye, such as BD DiIC<sub>1,2</sub>(3). After labeling, the migration or invasion assay is run with data collected over a time course using a bottom reading fluorescent plate reader.

FIGURE 6 • EFFECTS OF TIMP-2 AND 1'10' PHENANTHROLINE IN VEGF-MEDIATED HMVEC INVASION



Human microvascular endothelial cells (HMVECs) were assayed in the BD BioCoat Angiogenesis System: Endothelial Cell Invasion in the presence of VEGF (4 µg/ml) with varying concentrations of (left) TIMP-2 or (right) 1'10' phenanthroline in the bottom chamber. Cells were allowed to invade for 22 ± 1 hour. Cells were labeled post-invasion with BD Calcein AM (4 µg/ml) and then analyzed for invasion through BD Matrigel™ Matrix using an Applied Biosystems CytoFluor® 4000 plate reader [485/540 nm (Ex/Em) wavelengths]. Data represents the mean of n=3 inserts ± S.D.

## Tools for Endothelial Cell Culture

Cat. No.	Description	Qty.
----------	-------------	------

### Cell Culture Reagents

#### Extracellular Matrix Proteins

44743-720	BD Matrigel Basement Membrane Matrix Growth Factor Reduced	10 ml
-----------	--	-------

#### Cell Recovery Reagents

47743-724	Dispase	100ml
47743-696	Cell Recovery Solution	100 ml

#### Fluorescent Dyes

89044-506	DiIC <sub>12</sub> (3)	100 mg
89044-502	Calcein AM	10 x 50 µg

#### HUVEC Cells

BD354151	HUVEC-2 Cells	1 cryovial
----------	---------------	------------

#### Specialty Media

62405-070	Endothelial Cell Culture Media	500 ml
-----------	--------------------------------	--------

#### Cytokines and Media Additives

62405-784	Endothelial Cell Growth Supplement, bovine	15 mg
47743-610	Vascular Endothelial Growth Factor, human recombinant	10 µg

### Cell Culture Tools

#### BD BioCoat Collagen I Cellware

62405-617	100 mm Dish	10
-----------	-------------	----

### Cell Environments

#### BD BioCoat Cell Environment

62405-590	Endothelial Cell Growth Environment	1
-----------	-------------------------------------	---

### Membrane Insert Systems

#### BD BioCoat Angiogenesis System: Endothelial Cell Migration

47743-788	24-Multiwell Insert Plate with lid	1
-----------	------------------------------------	---

#### BD BioCoat Angiogenesis System: Endothelial Cell Invasion

BD354141	24-Multiwell Insert Plate with lid	1
----------	------------------------------------	---

#### BD BioCoat Angiogenesis System: Endothelial Tube Formation

47745-226	96-Multiwell Insert Plate with lid	1
-----------	------------------------------------	---


For a complete product listing, see page 19.

BD BioCoat™ Angiogenesis System: Endothelial Cell Migration consists of BD FluoroBlok™ inserts evenly coated with human fibronectin (**Figure 7**). Studies conducted using the post-labeling technique demonstrated that BD™ HUVEC-2 cells migrate towards VEGF in a concentration dependent manner (**Figure 8**).

During angiogenesis, endothelial cells form capillaries once they have invaded through the basement membrane. The correct culture surface is critical for successful endothelial cell tube formation *in vitro*.


\* BD BioCoat Angiogenesis System: Endothelial Cell Tube Formation offers a standardized and robust assay for studying endothelial cell tubulogenesis. For customers interested in establishing an assay for tube formation using vial BD Matrigel Matrix, we recommend pre-testing lots to ensure optimal performance.

FIGURE 7 • HUVEC MIGRATION ON UNCOATED AND HUMAN FIBRONECTIN-COATED INSERTS



Migration assays were conducted using HUVECs in the BD BioCoat Angiogenesis System: Endothelial Cell Migration and compared with uncoated BD FluoroBlok 24-Multiwell Inserts using both FBS (5%) and VEGF (10 µg/ml) as chemoattractants. The cells were allowed to migrate for 22 ± 1 hour. Cells were labeled post-migration with Calcein AM (4 µg/ml) and measured by detecting the fluorescence of the cells that migrated through the BD FluoroBlok membrane using an Applied Biosystems CytoFluor® 4000 plate reader [485/530 nm (Ex/Em) wavelengths]. The results indicate a marked increase in migration in response to VEGF when the assay was performed on the fibronectin-coated inserts included in the system. Data represents the mean of n=3 inserts ± S.D.


FIGURE 8 • BD HUVEC-2 CELLS EXHIBIT CONCENTRATION-DEPENDENT MIGRATION TOWARDS VEGF



BD HUVEC-2 cells assayed in the BD BioCoat Angiogenesis System: Endothelial Cell Migration (96-Multiwell format) in response to increasing concentrations of VEGF. Samples were incubated for 22 hours. Cells were labeled post-migration with BD Calcein AM and measured by detecting the fluorescence of cells that migrated through the fibronectin-coated BD FluoroBlok membrane with the Victor2™ plate reader (PerkinElmer) at 485 nm emission. Data represents the mean of n=4 inserts ± S.D.


Both primary endothelial cells and endothelial cell lines have been demonstrated to form tubules on the BD BioCoat™ Angiogenesis System: Endothelial Cell Tube Formation (**Figures 9-11**) which is comprised of a 3D gel of BD Matrigel Matrix. The BD BioCoat Angiogenesis Systems are available in 24- and 96-Multiwell formats, which can be used for moderate to high throughput compound screening. BD Matrigel™ Matrix has also been extensively used to study *in vivo* angiogenesis<sup>10-11, 16-18</sup> as a less technically challenging alternative to the corneal implantation model. A "plug" of material is placed subcutaneously, followed by histological quantification 7-10 days later. These *in vitro* and *in vivo* assays give researchers multiple options for exploring endothelial cell functions that are essential during angiogenesis.

FIGURE 9 • HUMAN ENDOTHELIAL CELL TYPES EXHIBIT TUBE FORMATION




HUVEC, HMVEC, and the human endothelial cell line HMEC-1 exhibit tube formation on BD BioCoat Angiogenesis System: Endothelial Cell Tube Formation. For this study, 20,000 cells of each cell type were added to wells containing pre-solidified BD Matrigel Matrix. The assay was incubated for 18 hours. Each bar represents the mean of n=32 wells ± S.D.

FIGURE 10 • CONFOCAL IMAGE OF BD™ HUVEC-2 CELL TUBE FORMATION



BD HUVEC-2 cells were assayed using the BD BioCoat Angiogenesis System: Endothelial Cell Tube Formation. Cells were stained using BD Calcein AM. Confocal images were captured using the BD Pathway™ Bioimager in confocal mode using the 4x objective (NA 0.13) for quantification of tubule formation.

FIGURE 11 • SURAMIN INHIBITS HMEC-1 TUBE FORMATION



HMEC-1 cells (40,000 cells/ml) were treated with Suramin at concentrations ranging from 0-40 μM and then analyzed for tube formation using BD BioCoat Angiogenesis System: Endothelial Cell Tube Formation. 50 μl of cells plus compound were added to wells containing pre-solidified BD Matrigel Matrix. Samples were incubated at 37°C, 5% CO<sub>2</sub> for 18 hours before staining with BD Calcein AM. Images were acquired with a 2x objective lens and the total tube length was measured using MetaMorph® (Universal Imaging Corporation™). Each bar represents the mean of n=8 wells ± S.D.




Pseudo-colored image for illustrative purposes only.

## Hepatocytes

Hepatocytes are liver epithelial cells used for both basic research and drug metabolism studies. Fresh and cryopreserved primary hepatocytes contain all the major enzyme pathways for drug and xenobiotic biotransformation. These include the major phase I drug metabolism enzyme family (P450) and phase II enzymes (UGT, SULT, GST and NAT). Hepatocytes also contain all the gene regulation pathways for P450 induction. Appropriate culture conditions are required to maintain hepatic P450 activity.

Hepatocytes can be cultured on Collagen I<sup>19-22</sup>, BD Matrigel™ Matrix<sup>23-27</sup> or BD™ PuraMatrix™<sup>28-29</sup>. BD BioCoat™ Collagen I Cellware is a commonly used surface for cultures of both fresh and cryopreserved hepatocytes<sup>30-31</sup> (**Figure 12**). Cells cultured on this surface maintain their biological activity, as shown by P450 induction (**Figure 13**). Sandwich cultures, such as hepatocytes grown on BD BioCoat Collagen I with BD Matrigel Matrix overlay, are used to assess bile canaliculi formation<sup>32</sup>. Choly-lysyl-fluorescein (CLF) is a fluorescein-labeled bile acid that is secreted into bile canaliculi by bile salt export pump (BSEP) which can be used to visualize bile canaliculi (**Figure 14**). BD Matrigel Matrix has been shown to suppress cell growth and prevent growth-associated dedifferentiation<sup>23</sup>, as well as maintain liver-specific functions *in vitro* longer than most collagen-based systems<sup>24-26</sup>. Hepatocytes cultured on BD Matrigel Matrix also have a more differentiated morphology than hepatocytes cultured on collagen I (**Figure 15**). Both BD Collagen I and BD Matrigel Matrix are animal-derived products; BD PuraMatrix, a synthetic peptide hydrogel, is a suitable alternative for assays that require a xeno-free culture environment. Therefore the appropriate culture surface depends on the experimental goals (e.g., drug metabolism, bile canaliculi formation or xeno-free environment).

FIGURE 12 • BD INDUCIBLE CRYOPRESERVED HUMAN HEPATOCYTES CULTURED ON BD BIOCOAT COLLAGEN I



BD Gentest™ Inducible-qualified Human CryoHepatocytes were isolated using the BD Gentest CryoHepatocyte Purification Kit and resuspended in freshly prepared ISOMs seeding media at a concentration of  $1 \times 10^6$  cells/ml. Cells were plated onto BD BioCoat Collagen I 24-well plates and incubated for approximately 2 hours, after which plating media was removed and replaced with supplemented BD Hepatocyte Culture Media.


### \* DID YOU KNOW?

- BD Biosciences offers a custom barcoding service. This service provides high-quality barcode labels affixed to any side of a microplate.




FIGURE 13 • INDUCTION OF BD GENTEST™ INDUCIBLE-QUALIFIED HUMAN CRYOHEPATOCYTES

## A. CYP3A4




## B. CYP1A2



BD Gentest Inducible-qualified Human CryoHepatocytes were isolated using the BD Gentest CryoHepatocyte Purification Kit and resuspended into freshly prepared ISOMs seeding media at a concentration of  $1 \times 10^6$  cells/ml. Cells were plated onto BD BioCoat™ Collagen I 24-well Multiwell Plates and incubated for approximately 2 hours, after which plating media was removed and replaced with supplemented BD™ Hepatocyte Culture Media. Cells were monitored for degree of attachment at 18-24 hours after plating and daily during the experiment. Cells were induced with either 20  $\mu$ M Rafampicin (A) or 20  $\mu$ M  $\beta$ -Naphthoflavone (B) over a 3-day period. Controls were treated with the appropriate solvent control. Metabolic activity was determined on day 5 of the experiment using 200  $\mu$ M Testosterone as a substrate to measure CYP3A4 activity and 100  $\mu$ M Phenacetin as a substrate for CYP1A2. Assays were run for 30 minutes and 60 minutes, respectively. Analysis was performed by HPLC and activity expressed per mg of protein.


FIGURE 14 • BD GENTEST CHOLY-LYSYL-FLUORESCIN SEQUESTERED IN BILE CANALICULI




CLF sequestered in the bile canaliculi of BD Gentest Inducible-qualified Human CryoHepatocytes cultured on BD BioCoat Collagen I overlaid with BD Matrigel Matrix.

FIGURE 15 • EFFECTS OF ECM ON CELL MORPHOLOGY: MICROGRAPHS OF HEPATOCYTES CULTURED ON VARIOUS CULTURE SUBSTRATA


## A.



## B.



## C.



Scanning electron micrographs of primary rat hepatocytes cultured for two days on collagen I (A), collagen I gel (B), or BD Matrigel Matrix (C). Note the clusters of spherical cells for hepatocytes cultured on BD Matrigel Matrix, typical of differentiated cells.

## Tools for Hepatocyte Cell Culture

Cat. No.	Description	Qty.
----------	-------------	------

## Cell Culture Reagents

## Hepatocyte Culture Media Kit

62405-070	Hepatocyte Culture Media Kit	500 ml
-----------	------------------------------	--------

## Extracellular Matrix Proteins

47743-656	Collagen I, rat tail	100 mg
47743-716	BD Matrigel Matrix, phenol red-free	10 ml

## Cell Recovery Reagents

47743-724	Dispase	100 ml
47743-696	Cell Recovery Solution	100 ml

## Cell Culture Tools

## BD BioCoat™ Collagen I

62405-601	6-well plates	5
-----------	---------------	---

## BD BioCoat Matrigel Matrix Multiwell Plates for Hepatocytes


62405-144	6-well plates	5
-----------	---------------	---

## Cell Environments

## Hepatocyte Differentiation Environment

62405-096	6-well plate	1
-----------	--------------	---

For a complete product listing, see page 19.




## Neuronal Cells

Neuroscience is a rapidly evolving field that encompasses a variety of cell types, including neurons and neuronal stem cells. *In vitro* culture of these diverse cell types requires appropriate culture surfaces for attachment and proliferation/differentiation, as detailed in the examples below. NG-108 rat glioma/mouse neuroblastoma cells and PC-12 cells, two neuronal cell lines, require different surfaces for attachment. NG-108 cells attach loosely to tissue culture-treated cellware, but when they are cultured on BD BioCoat™ Laminin Cellware they exhibit a more typical neuronal morphology (**Figure 16**). PC-12 cells, derived from a transplantable rat pheochromocytoma, develop neurites in response to NGF when they are cultured on collagen I (**Figure 17**). Other surfaces, including BD BioCoat Poly-D-Lysine Cellware<sup>33</sup> and BD BioCoat Poly-D-Lysine/Laminin<sup>34</sup>, can also be used to culture PC-12 cells. Primary neuronal cells utilize different attachment surfaces depending on their origin and the composition of the media used during culture. Primary mouse cortical neurons and primary mouse basal forebrain cholinergic neurons have been cultured on BD BioCoat Poly-L-Lysine Cellware<sup>35</sup> and BD BioCoat Poly-D-Lysine/Laminin Cellware<sup>36</sup>, respectively. Primary human neural stem cells have been grown under serum-containing conditions in tissue culture-treated BD Falcon Cell Culture Flasks<sup>37</sup>. Using serum-free conditions, Thonhoff, et al., showed that neuronal stem cells maintain their capacity to differentiate into both Tuj1<sup>+</sup> neuronal cells and GFAP<sup>+</sup> astroglial cells on BD™ PuraMatrix™ while differentiation of neuronal stem cells grown on BD Matrigel™ Matrix was skewed toward GFAP<sup>+</sup> astroglial cells<sup>38</sup>. Both BD PuraMatrix<sup>38-40</sup> and BD Primaria™<sup>41</sup> are defined, xeno-free surfaces for 3D and 2D culture, respectively, which are compatible with neuronal cells. BD Primaria Cultureware enhances neuronal cell attachment as compared to tissue culture-treated cellware, as shown with chick embryo spinal cord neurons (**Figure 18**). These examples\* illustrate the need for an appropriate growth surface which is determined by the cell type and whether a xeno-free surface with defined media is required by the experimental model.

\*Other examples available in references 42-44.

FIGURE 16 • EFFECTS OF BD BIOCOAT LAMININ CELLWARE ON NG-108 NEUROBLASTOMA CELLS




NG-108 rat glioma/mouse neuroblastoma cell morphology is surface dependent. Cells cultured on tissue culture plastic are loosely adhered and remain rounded (A). Cells cultured on BD BioCoat Laminin cellware exhibit a spindle-shaped morphology and dendritic processes (B).



### DID YOU KNOW?

- BD Biosciences offers a full range of 96-, 384-, and 1536-well Microplates. Custom packaging, labeling (e.g., barcoding), and custom coatings are also available. Please contact your sales representative for more information.


FIGURE 17 • PC12 NEURITE OUTGROWTH, CULTURED ON BD™ COLLAGEN I




PC12 cells were maintained in DMEM with 10% FBS, 5% horse serum and 1% penicillin/streptomycin. For neurite generation, approximately 15,000 cells/well were plated in BD Falcon™ 96-well Imaging Plates that were coated with BD Collagen I, rat tail using 1.8 µg collagen per well. After 24 hours, the medium was replaced with differentiation medium (DMEM with 0.1% FBS, 0.05% horse serum, 100 ng/ml NGF). The medium was replenished every third day for 10 days. For imaging, cells were fixed with 3.7% paraformaldehyde for 20 minutes and permeabilized with 0.1% Triton-X-100 for 5 minutes. Neurites were stained with a primary mouse anti-β-tubulin antibody (Cat. No. 556321) using 0.125 µg antibody/well followed by AlexaFluor® 488 goat anti-mouse IgM at a concentration of 0.25 µg/well. Hoechst 33342 was used at 0.1 µg/well to stain the nuclei. To prevent the dissociation and fracture of fragile neuronal networks, the number of washes in the fixation and processing steps were minimized and extra care was taken in aspirating and dispensing liquids in wells. Images were acquired on a BD Pathway™ as a 4x4 montage using a 20x objective (0.75 NA).

FIGURE 18 • CHICK EMBRYO SPINAL CORD NEURONS CULTURED ON BD PRIMARIA™ CULTUREWARE

A.



B.



When chick embryo spinal cord neurons are cultured on BD Primaria Cultureware, growth is enhanced and extensive neurite development occurs. In this experiment, cells clumped and detached from traditional TC plates after 20 days in culture (A) but remained viable and differentiated on BD Primaria Cultureware (B).

## Tools for Neuronal Cell Culture

Cat. No.	Description	Qty.
----------	-------------	------

### Cell Culture Reagents

#### Extracellular Matrix Proteins

47743-656	Collagen I, rat tail	100 mg
47743-728	Fibronectin, human	1 mg
47743-734	Laminin, mouse	1 mg
47743-715	BD Matrigel Matrix	10 ml
47743-736	Poly-D-Lysine	20 mg

#### Cytokines and Media Additives

47743-594	7S Nerve Growth Factor, mouse, natural	100 µg
47743-588	2.5S Nerve Growth Factor, mouse, natural	10 µg
47743-566	Endothelial Growth Factor, human recombinant	100 µg

#### Cell Recovery Reagents

47743-724	Dispase	100 ml
47743-696	Cell Recovery Solution	100 ml

### Cell Culture Tools

#### BD BioCoat™ Laminin Cellware

62405-706	6-well plates	5
-----------	---------------	---

#### BD BioCoat Poly-L-Ornithine/Laminin Cellware

62405-747	96-well plates	5
-----------	----------------	---

#### BD BioCoat Poly-D-Lysine/Laminin Cellware

62405-111	24-well plates	5
-----------	----------------	---

#### BD BioCoat Poly-D-Lysine Cellware

62405-747	6-well	5
-----------	--------	---

#### BD Primaria Cultureware

25382-687	60 x 15 mm Dish with lid	200
-----------	--------------------------	-----

#### BD Falcon™ CultureSlides


62405-178	8-well	96
-----------	--------	----

#### BD Falcon 96-well Imaging Plate

89022-352	Black/Clear, with lid	32
-----------	-----------------------	----

For a complete product listing, see page 19.






## Epithelial Cells

Epithelial cells are found throughout the body, from skin to glandular formations within tissues. *In vivo* these cells are attached to a three dimensional basement membrane matrix. The interactions between the epithelial cell and matrix proteins effect cell morphology and function. Two highly specified epithelial cell types have been discussed in the hepatocyte and endothelial cell sections, utilizing both 2-dimensional (2D) and three-dimensional (3D) culture systems. *In vitro*, 2D and 3D culture systems can be used to study different aspects of cell growth and differentiation. 2D culture systems are used for cell attachment and proliferation. 3D environments are utilized in studies requiring a more *in vivo*-like setting, such as mammary acini formation.

The BD BioCoat Cellware provides a range of 2D surfaces for cell growth. Both keratinocytes<sup>45-46</sup> and HEK-293<sup>47-49</sup> cells are examples of epithelial cells that can be studied in 2D culture environments. Keratinocytes are a major component of the epidermis; BD BioCoat Collagen I supports growth of human neonatal keratinocytes (**Figure 19**). HEK-293 cells are a human epithelial kidney cell line which exhibit enhanced attachment to poly-lysine coated surfaces as compared to tissue culture-treated surfaces. This is particularly important if the cells need to remain attached during subsequent washes (**Figure 20**). The appropriate 2D surface is determined by the cell type.

**FIGURE 19 • PROLIFERATION OF HUMAN NEONATAL KERATINOCYTES ON BD BIOCOAT™ COLLAGEN I**



Human neonatal keratinocytes cultured on BD BioCoat Collagen I.

Three-dimensional growth substrates can support certain cellular behaviors that are not observed when cells are cultured on a planar two-dimensional surface, as exemplified by mammary epithelial<sup>50-54</sup> and Caco-2<sup>55-56</sup> cells. *In vivo*, mammary epithelial cells form polarized acini. When tumorigenic human mammary carcinoma cells (T4-2) are cultured on a 3D substrate comprised of reconstituted basement membrane (Growth Factor Reduced BD Matrigel Matrix) they form large disorganized colonies, as shown with the T4-vector control in a study from Dr. Bissell's laboratory<sup>51</sup> (**Figure 21**). Epidermal growth factor receptor (EGFR) had previously been shown to be elevated in T4-2 cells, and downregulation of this signaling pathway in T4-2 cells cultured in 3D BD Matrigel™ Matrix is known to lead to phenotypic reversion to polarized acini. These cells exhibit polarized acinar architecture in the presence of the EGFR inhibitor AG1478 or when stably expressing dominant negative Rap1 (T4-DN-Rap1); reversion to a normal phenotype is shown by proper localization of  $\alpha 6$ -integrin (basal marker),  $\beta$ -catenin (basolateral marker) and GM130 (apical marker). These data show that three-dimensional BD Matrigel Matrix culture conditions are conducive to studying signaling pathways involved in regulating mammary acinar architecture.

Another example of the effect of 3D growth substrates on cellular phenotypes is the use of BD BioCoat Fibrillar Collagen Inserts in Caco-2 assays. Caco-2 cells are an epithelial cell line derived from a colorectal adenocarcinoma commonly used to measure compound permeability. The gold standard for modeling drug permeability across the intestinal epithelium *in vitro* is measuring permeability across differentiated Caco-2 cells, where the cells have been cultured for 21 days on cell culture inserts. BD BioCoat HTS Caco-2 Assay System and BD BioCoat Intestinal Epithelium Differentiation Environment utilize BD BioCoat Fibrillar Collagen Inserts and a specialized media to enhance the rate of Caco-2 differentiation from 21 to 3 days (**Figures 22-23**), thereby reducing the time and labor required for the analysis of compound permeability.

The 2D and 3D cell culture systems available from BD Biosciences provide multiple options to researchers studying epithelial cells *in vitro*.




### DID YOU KNOW?

- BD Biosciences offers custom coatings. Please contact your sales representative for more information.



FIGURE 20 • ADHERANCE OF HEK-293 CELLS TO  
BD BIOCOAT™ POLY-D-LYSINE CULTUREWARE



HEK-293 cells have enhanced attachment to BD BioCoat Poly-D-Lysine Cultureware as compared to BD Falcon Tissue Culture-treated Cultureware. An equal number of cells were plated on BD BioCoat Poly-D-Lysine 384-well black/clear (right) and BD Falcon Tissue Culture-treated 384-well Black/Clear Plates (left) and grown under serum-free conditions. Before washing (top), there were a similar number of cells in the BD BioCoat Poly-D-Lysine coated wells and the BD Falcon Tissue Culture-treated wells. After washing, using a Skatron Washer (Molecular Devices) (middle), the cells remained attached to the BD BioCoat Poly-D-Lysine wells while few cells remained attached to the BD Falcon Tissue Culture-treated wells. Post-wash, the cells were visualized using Calcein AM (bottom).

## Tools for Epithelial Cell Culture

Cat. No.	Description	Qty.
----------	-------------	------

### Cell Culture Reagents

#### Extracellular Matrix Proteins

47743-712	Collagen I, rat tail	10 x 100 mg
47743-706	BD Matrigel Matrix	5 ml

#### Cell Recovery Reagents

47743-724	Dispase	100 ml
47743-696	Cell Recovery Solution	100 ml

### Cell Culture Tools

#### BD BioCoat™ Collagen I Cellware

12777-074	75 cm <sup>2</sup> vented-cap Flasks	5
-----------	--------------------------------------	---

#### BD BioCoat Poly-D-Lysine Cellware

62406-048	100 mm Dishes	10
-----------	---------------	----

### Cell Environments

#### Intestinal Epithelium

##### Differentiation Environment

62405-098	Intestinal Epithelium Differentiation Environment	1
-----------	---	---

#### BD BioCoat HTS Caco-2 Assay Systems

12777-040	BD BioCoat Fibrillar Collagen 24-Multiwell Insert System plus media to perform 24 individual three-day Caco-2 assays	1
62405-446	Differentiation Medium	2 x 250 ml
62405-442	Intestinal Epithelium Differentiation Media Pack	1 kit
47743-630	MITO+ Serum Extender	5 ml
62405-444	Seeding Basal Medium	2 x 250 ml
12777-044	Five BD BioCoat Fibrillar Collagen 1 24-Multiwell Insert System plus media to perform 24 individual three-day Caco-2 assays	1


### Membrane Insert Systems

#### BD BioCoat Fibrillar Collagen Cell Culture Inserts

62405-119	1 $\mu$ m inserts in four 6-well plates	24
-----------	---	----


For a complete product listing, see page 19.

FIGURE 21 • EFFECT OF RAP1 ACTIVITY ON T4-2 CELL POLARITY IN 3D GROWTH FACTOR REDUCED BD MATRIGEL™ MATRIX CULTURE




Growth Factor Reduced BD Matrigel Matrix supports mammary acini formation *in vitro*. Malignant T4-2 cells were grown in three-dimensional culture on Growth Factor Reduced BD Matrigel Matrix. Cells were stably transfected with control (T4-vector) or dominant negative-Rap1 (T4-DN-Rap1). Inhibition of EGFR with AG1478 was used as a positive control for reversion of T4-2 to normal mammary acinar architecture. Indirect immunofluorescence was used to analyze cell polarity markers for basal ( $\alpha 6$ -interin), basolateral ( $\beta$ -catenin) and apical (GM130) membrane domains. Bar, 5  $\mu$ m. Images kindly provided by Dr. Masahiko Itoh and Dr. Mina Bissell, originally published in Cancer Research 67(10):4759-4766<sup>51</sup>. Reproduced with permission.

FIGURE 22 • PERMEABILITY OF MANNITOL AND ANTIBIOTICS THROUGH CACO-2 MONOLAYERS



Barrier formation occurs three days post-seeding in the BD BioCoat Intestinal Epithelium Differentiation Environment and two to four weeks with conventional methods. Monolayers formed using either the BD BioCoat Intestinal Epithelium Differentiation Environment or conventional methods are equally permeable for each of the three compounds tested.


FIGURE 23 • P-GLYCOPROTEIN (P-GP) FUNCTION IN CACO-2 CELLS



Caco-2 cells were cultured using the three-day BD BioCoat HTS Caco-2 Assay System supplemented with MITO+ Serum Extender (A) or the traditional 21-day system (B). P-gp function was assessed by adding 10 nM <sup>3</sup>H-labeled vinblastine in PBS to either the apical or basal side of the insert. Samples were withdrawn from the non-labeled side of the insert and counted by scintillation counting. To inhibit the P-gp with verapamil, 100  $\mu$ M verapamil was added to the insert chambers.


# Tumor Cells

Cancerous cells have altered cellular functions as compared to the normally functioning, non-malignant cells from which they are derived. Cell morphology and signaling pathway studies *in vitro* that incorporate the use of 3D culture systems can give insights into the effects of mis-regulated or mis-expressed proteins, as exemplified by human mammary carcinoma cells (T4-2)<sup>51</sup> (**Figure 20**). The hallmark of metastatic cells is their ability to invade through the basement membrane and migrate to other parts of the body. Cell migration can be studied using either BD Falcon™ Cell Culture Inserts or BD FluoroBlok™ Cell Culture Inserts for moderate to high-throughput screening (**Figure 24**). Cells must be able to both secrete proteases that break down the basement membrane as well as migrate in order to be invasive. Invasion through BD Matrigel™ Matrix-coated Cell Culture Inserts has become the gold standard for quantitative and qualitative measurement of the metastatic potential of a cell<sup>10, 57-63</sup>. This matrix provides a true barrier to non-invasive cells while presenting the appropriate protein structure for penetration of invading cells.



Pseudo-colored image for illustrative purposes only.

FIGURE 24 • HT-1080 MIGRATION



Migration of Calcein AM (A) and DiIC<sub>12</sub>(3) (B) labeled human fibrosarcoma cells (HT-1080) through BD Falcon FluoroBlok 96-Multiwell Inserts, 8 µm pore size. DMEM with 5% FCS was used as a chemoattractant in the lower wells, while DMEM/0.1% BSA was added to the control wells. The plates were incubated for four hours at 37°C, after which fluorescence of cells which had migrated through the microporous membrane was measured on the Applied Biosystems CytoFluor® 4000 and PerkinElmer HTS 7000 Plus fluorescent plate readers using excitation/emission wavelengths of 485/530 nm for Calcein AM or 530/590 nm for DiIC<sub>12</sub>(3). Values represent the mean of 8 wells ± S.D. Migration from as few as 4,000 input cells can be detected.

## Tools for Tumor Cell Culture

Cat. No.	Description	Qty.
----------	-------------	------

### Cell Culture Reagents

#### Extracellular Matrix Proteins

47747-220	BD Matrigel Matrix, High Concentration	10 ml
-----------	--	-------

#### Cell Recovery Reagents

47743-696	Cell Recovery Solution	100 ml
47743-724	Dispase	100 ml

#### Fluorescent Dyes

89044-502	Calcein AM	10 x 50 µg
89044-506	DiIC <sub>12</sub> (3)	100 mg

### Membrane Insert Systems

#### BD BioCoat Matrigel Invasion Chambers

62405-744	8.0 µm inserts in two 24-well plates	24
-----------	--------------------------------------	----

#### BD BioCoat Tumor Invasion System

47733-148	One insert plate with 24-well plate and lid	1
-----------	---	---

#### BD BioCoat Cell Culture Inserts

62405-722	Laminin, 3 µm inserts in two 24-well plates	24
-----------	---	----

#### BD Falcon Cell Culture Inserts

62406-226	3 µm pore size with 24-well plate and lid	1
-----------	---	---

For a complete product listing, see page 19.




### DID YOU KNOW?

- BD Biosciences offers a full range of dishes and flasks. Please contact your sales representative for more information.


The BD BioCoat™ Matrigel Invasion Chambers and BD BioCoat Tumor Invasion Systems are optimized systems that utilize standardized coating procedures to ensure even coating of BD Matrigel™ Matrix for reproducible results (**Figure 25**). The BD BioCoat Tumor Invasion System provides a unique, quantitative platform that can be used to determine the effects of anti-metastatic compounds on invasive cell types (**Figure 26**). For *in vivo* studies, BD Matrigel Matrix can be used to help support tumor cell engraftment in mice<sup>64-66</sup>. These tools allow researchers to dissect various areas of tumor biology, from analysis of signaling pathways *in vitro* to *in vivo* tumor formation.

FIGURE 25 • COMPARISON OF MEAN PERCENT INVASION

#### A. BD BioCoat 96-Multiwell Tumor Invasion System




#### B. BD BioCoat 24-Multiwell Tumor Invasion System



Multiple lots of the BD BioCoat 96-Multiwell Tumor Invasion System and BD BioCoat 24-Multiwell Tumor Invasion System were assayed to show reproducibility with these systems. Multiple lots of BD BioCoat 96-Multiwell Tumor Invasion System (A) and BD BioCoat 24-Multiwell Tumor Invasion System (B) were assayed. Fluorescently labeled cells residing on the bottom of the insert membrane were measured post-invasion with either a Victor2™ plate reader (BD BioCoat 96-Multiwell Tumor Invasion System) or a CytoFluor® plate reader (BD BioCoat 24-Multiwell Tumor Invasion System). Mean percent invasion of NIH-3T3 and HT-1080 cells were compared. Cells were labeled post-invasion using BD Calcein AM.

FIGURE 26 • INHIBITION OF PC3 MIGRATION AND INVASION BY DOXYCYCLINE



PC3 invasion is inhibited by doxycycline. PC3 cell invasion was measured using BD BioCoat 24-Multiwell Tumor Invasion System, which is based on the fluorescence blocking BD FluoroBlok™ PET microporous membrane, and migration was measured using BD Falcon HTS FluoroBlok 24-Multiwell Insert System. At the end of the assay, cells were stained with BD Calcein AM.



## Cell Culture Reagents

### Extracellular Matrix Proteins

	DESCRIPTION	QTY./CASE	CAT. NO.
BD Matrigel™ Basement Membrane Matrix	BD Matrigel Matrix	5 ml	<a href="#">47743-706</a>
	BD Matrigel Matrix	10 ml	<a href="#">47743-715</a>
	BD Matrigel Matrix (50 ml)	5 x 10 ml	<a href="#">47743-710</a>
	BD Matrigel Matrix High Concentration (HC)	10 ml	<a href="#">47747-220</a>
	BD Matrigel Matrix Phenol Red-Free	10 ml	<a href="#">47743-716</a>
	BD Matrigel Matrix HC Phenol Red-free	10 ml	<a href="#">80094-328</a>
	BD Matrigel Matrix Growth Factor Reduced (GFR)	5 ml	<a href="#">47743-718</a>
	BD Matrigel Matrix GFR	10 ml	<a href="#">47743-720</a>
	BD Matrigel Matrix HC GFR	10 ml	<a href="#">80094-330</a>
	BD Matrigel Matrix Phenol Red-free GFR	10 ml	<a href="#">47743-722</a>
Fibronectin	Fibronectin, human	1 mg	<a href="#">47743-728</a>
	Fibronectin, human	5 mg	<a href="#">47743-654</a>
	Fibronectin, human (25 mg)	5 x 5 mg	<a href="#">47743-730</a>
Collagen I	Collagen I, bovine	30 mg	<a href="#">47743-676</a>
	Collagen I, human	0.25 mg	<a href="#">47743-688</a>
	Collagen I, rat tail	100 mg	<a href="#">47743-656</a>
	Collagen I, rat tail (1 g)	10 x 100 mg	<a href="#">47743-712</a>
	Collagen I, human recombinant	250 ug	<a href="#">12000-612</a>
Laminin	Laminin, mouse	1 mg	<a href="#">47743-734</a>
	Ultra-pure Laminin, mouse	1 mg	<a href="#">47743-682</a>
	Laminin/Entactin Complex High Concentration	10.5 mg	<a href="#">80085-668</a>
Poly-D-Lysine	Poly-D-Lysine, synthetic	20 mg	<a href="#">47743-736</a>

### Cytokines and Media Additives

	DESCRIPTION	QTY./CASE	CAT. NO.
Epidermal Growth Factor (EGF)	Mouse, natural (culture grade)	100 µg	<a href="#">62405-786</a>
	Mouse, natural (culture grade) (10 x 100 µg)	1 mg	<a href="#">47743-756</a>
	Mouse, natural (receptor grade)	100 µg	<a href="#">47743-562</a>
	Mouse, natural (receptor grade) (5 x 100 µg)	500 µg	<a href="#">47743-564</a>
	Human recombinant	100 µg	<a href="#">47743-566</a>
	Human recombinant (10 x 100 µg)	1 mg	<a href="#">47743-568</a>
Basic Fibroblast Growth Factor (bFGF)	bFGF, bovine natural	10 µg	<a href="#">47743-572</a>
	bFGF, human recombinant	10 µg	<a href="#">47743-574</a>
	bFGF, human recombinant (50 µg)	5 x 10 µg	<a href="#">47743-670</a>
	bFGF, human recombinant (100 µg)	10 x 10 µg	<a href="#">47743-672</a>
ITS Universal Culture Supplement Premix	5 liter equivalent	5 ml	<a href="#">47743-626</a>
	20 liter equivalent	20 ml	<a href="#">47743-624</a>
Nerve Growth Factor (NGF)	2.5S NGF, mouse natural	10 µg	<a href="#">47743-588</a>
	2.5S NGF, mouse natural	100 µg	<a href="#">47743-590</a>
	2.5S NGF, mouse natural (1 mg)	2 x 500 µg	<a href="#">47743-592</a>
	7S NGF, mouse natural	100 µg	<a href="#">47743-594</a>
Vascular Endothelial Growth Factor (VEGF)	Human recombinant	10 µg	<a href="#">47743-610</a>
MITO+ Serum Extender	5 liter equivalent	5 ml	<a href="#">47743-630</a>
Endothelial Cell Growth Supplement (ECGS)	Bovine	15 mg	<a href="#">62405-784</a>
	Bovine	100 mg	<a href="#">47743-652</a>
Specialty Media	E-STIM Endothelial Cell Culture Medium	500 ml	<a href="#">62405-070</a>
	Hepatocyte Culture Media	500 ml	<a href="#">62405-898</a>
	Intestinal Differentiation Media Pack	1 pack	<a href="#">62405-442</a>
	Seeding Basal Medium	2 x 250 ml	<a href="#">62405-444</a>
	Enterocyte Differentiation Medium	2 x 250 ml	<a href="#">62405-446</a>
HUVEC-2 Cells	HUVEC-2 Cells	1 cryovial	<a href="#">BD354151</a>

### BD™ Cell Recovery/Detachment Reagents

Cell Recovery Reagents	Dispase	100 ml	<a href="#">47743-724</a>
	Cell Recovery Solution	100 ml	<a href="#">47743-696</a>

### BD Fluorescent Dyes

Fluorescent Dyes	Calcein AM Fluorescent Dye	10 x 50 µg	<a href="#">89044-502</a>
	Calcein AM Fluorescent Dye	1 mg	<a href="#">89044-504</a>
	DiIC <sub>12</sub> (3) Fluorescent Dye	100 mg	<a href="#">89044-506</a>

## Cell Culture Tools

### BD BioCoat™ Collagen I Cellware

DESCRIPTION	QTY./CASE	CAT. NO
6-well plates	5	<a href="#">62405-601</a>
6-well plates (10 sleeves of 5)	50	<a href="#">62405-413</a>
12-well plates	5	<a href="#">62405-603</a>
12-well plates (10 sleeves of 5)	50	<a href="#">62405-415</a>
24-well plates	5	<a href="#">62405-607</a>
24-well plates (10 sleeves of 5)	50	<a href="#">62405-417</a>
48-well plates	5	<a href="#">62405-609</a>
48-well plates (10 sleeves of 5)	50	<a href="#">62406-020</a>
96-well plates	5	<a href="#">62405-611</a>
96-well plates (10 sleeves of 5)	50	<a href="#">62405-419</a>
96-well plates	80	<a href="#">47748-278</a>
96-well black/clear plates	5	<a href="#">62406-290</a>
96-well black/clear plates (10 sleeves of 5)	50	<a href="#">62406-266</a>
96-well black/clear plates	80	<a href="#">47748-282</a>
96-well white/clear plates	5	<a href="#">62406-388</a>
96-well white/clear plates (10 sleeves of 5)	50	<a href="#">62406-386</a>
96-well white plates	5	<a href="#">62406-288</a>
96-well white plates (10 sleeves of 5)	50	<a href="#">62406-286</a>
96-well white plates	80	<a href="#">47748-280</a>
96-well white/clear plates	80	<a href="#">47748-284</a>
35 mm culture dishes	20	<a href="#">62405-613</a>
35 mm culture dishes (5 sleeves of 20)	100	<a href="#">62405-421</a>
60 mm culture dishes	20	<a href="#">62405-615</a>
60 mm culture dishes (5 sleeves of 20)	100	<a href="#">62405-423</a>
100 mm culture dishes	10	<a href="#">62405-617</a>
100 mm culture dishes (4 sleeves of 10)	40	<a href="#">62405-425</a>
150 mm culture dishes	5	<a href="#">62405-619</a>
25 cm <sup>2</sup> plug-seal flasks	10	<a href="#">62405-539</a>
25 cm <sup>2</sup> plug-seal flasks (5 sleeves of 10)	50	<a href="#">62405-427</a>
25 cm <sup>2</sup> vented-cap flasks	10	<a href="#">62405-025</a>
25 cm <sup>2</sup> vented-cap flasks (5 sleeves of 10)	50	<a href="#">12777-072</a>
75 cm <sup>2</sup> plug-seal flasks	5	<a href="#">62405-621</a>
75 cm <sup>2</sup> plug-seal flasks (10 sleeves of 5)	50	<a href="#">62405-429</a>
75 cm <sup>2</sup> vented-cap flasks	5	<a href="#">12777-074</a>
75 cm <sup>2</sup> vented-cap flasks (10 sleeves of 5)	50	<a href="#">12777-076</a>
150 cm <sup>2</sup> plug-seal flasks	5	<a href="#">62405-875</a>
150 cm <sup>2</sup> plug-seal flasks (8 sleeves of 5)	40	<a href="#">62406-022</a>
150 cm <sup>2</sup> vented-cap flasks	5	<a href="#">12777-078</a>
150 cm <sup>2</sup> vented-cap flasks (8 sleeves of 5)	40	<a href="#">12777-080</a>
Coverslips 22 mm round No.1 German glass	60	<a href="#">BD354089</a>
1-well CultureSlides	12	<a href="#">62406-704</a>
2-well CultureSlides	12	<a href="#">62405-876</a>
4-well CultureSlides	12	<a href="#">62406-716</a>
8-well CultureSlides	12	<a href="#">62405-878</a>

### BD BioCoat™ Poly-D-Lysine Cellware

DESCRIPTION	QTY./CASE	CAT. NO
6-well plates	5	<a href="#">62405-747</a>
6-well plates (10 sleeves of 5)	50	<a href="#">62406-024</a>
12-well plates	5	<a href="#">62406-026</a>
12-well plates (10 sleeves of 5)	50	<a href="#">62406-028</a>
24-well plates	5	<a href="#">62405-749</a>
24-well plates (10 sleeves of 5)	50	<a href="#">62406-030</a>
48-well plates	5	<a href="#">62406-718</a>
48-well plates (10 sleeves of 5)	50	<a href="#">62406-700</a>
96-well plates	5	<a href="#">62405-027</a>
96-well plates (10 sleeves of 5)	50	<a href="#">62406-032</a>
96-well plates	80	<a href="#">47748-262</a>
96-well black/clear plates	5	<a href="#">62406-034</a>
96-well black/clear plates (10 sleeves of 5)	50	<a href="#">62406-036</a>
96-well black/clear plates	80	<a href="#">47748-266</a>
96-well white/clear plates	5	<a href="#">62406-368</a>
96-well white/clear plates (10 sleeves of 5)	50	<a href="#">62406-366</a>
96-well white/clear plates	80	<a href="#">47748-268</a>
96-well white plates	5	<a href="#">62406-284</a>
96-well white plates (10 sleeves of 5)	50	<a href="#">62406-282</a>
96-well white/opaque plates	80	<a href="#">47748-264</a>
35 mm culture dishes	20	<a href="#">62406-038</a>
35 mm culture dishes (5 sleeves of 20)	100	<a href="#">62406-040</a>
60 mm culture dishes	20	<a href="#">62406-042</a>
60 mm culture dishes (5 sleeves of 20)	100	<a href="#">62406-044</a>
100 mm culture dishes	10	<a href="#">62406-084</a>
100 mm culture dishes (4 sleeves of 10)	40	<a href="#">62406-050</a>
150 mm culture dishes	5	<a href="#">62406-702</a>
25 cm <sup>2</sup> plug-seal flasks	10	<a href="#">62405-752</a>
25 cm <sup>2</sup> plug-seal flasks (5 sleeves of 10)	50	<a href="#">62406-052</a>
25 cm <sup>2</sup> vented-cap flasks	10	<a href="#">12777-088</a>
25 cm <sup>2</sup> vented-cap flasks (5 sleeves of 10)	50	<a href="#">12777-090</a>
75 cm <sup>2</sup> plug-seal flasks	5	<a href="#">62405-755</a>
75 cm <sup>2</sup> plug-seal flasks (10 sleeves of 5)	50	<a href="#">62406-054</a>
75 cm <sup>2</sup> vented-cap flasks	5	<a href="#">12777-092</a>
75 cm <sup>2</sup> vented-cap flasks (10 sleeves of 5)	50	<a href="#">12777-094</a>
150 cm <sup>2</sup> plug-seal flasks	5	<a href="#">62405-887</a>
150 cm <sup>2</sup> plug-seal flasks (8 sleeves of 5)	40	<a href="#">62406-056</a>
150 cm <sup>2</sup> vented-cap flasks	5	<a href="#">12777-084</a>
150 cm <sup>2</sup> vented-cap flasks (8 sleeves of 5)	40	<a href="#">12777-096</a>
Coverslips 12 mm round No.1 German glass	80	<a href="#">BD354086</a>
35 mm Coverslip-bottom dishes	20	<a href="#">BD354077</a>
1-well CultureSlides	12	<a href="#">62406-709</a>
2-well CultureSlides	12	<a href="#">62405-888</a>
4-well CultureSlides	12	<a href="#">62406-714</a>
8-well CultureSlides	12	<a href="#">62405-890</a>

**BD BioCoat™ Poly-L-Lysine Cellware**

DESCRIPTION	QTY./CASE	CAT. NO
6-well plates	5	<a href="#">62406-280</a>
6-well plates (10 sleeves of 5)	50	<a href="#">62406-278</a>
96-well plates	5	<a href="#">62406-276</a>
96-well plates (10 sleeves of 5)	50	<a href="#">62406-274</a>
35 mm culture dishes	20	<a href="#">62406-272</a>
35 mm culture dishes (5 sleeves of 20)	100	<a href="#">62406-270</a>
60 mm culture dishes	20	<a href="#">62406-268</a>
60 mm culture dishes (5 sleeves of 20)	100	<a href="#">62406-292</a>
Coverslips 12 mm round No.1 German glass	80	BD354085

**BD BioCoat™ Laminin Cellware**

DESCRIPTION	QTY./CASE	CAT. NO
6-well plates	5	<a href="#">62405-694</a>
12-well plates	5	<a href="#">62405-696</a>
24-well plates	5	<a href="#">62405-698</a>
48-well plates	5	<a href="#">62405-700</a>
96-well plates	5	<a href="#">62405-702</a>
35 mm culture dishes	20	<a href="#">62405-704</a>
60 mm culture dishes	20	<a href="#">62405-706</a>
100 mm culture dishes	10	<a href="#">62405-708</a>
150 mm culture dishes	5	<a href="#">62405-710</a>
25 cm <sup>2</sup> plug-seal flasks	10	<a href="#">62405-048</a>
75 cm <sup>2</sup> plug-seal flasks	10	<a href="#">62405-712</a>

**BD BioCoat™ Matrigel™ Matrix – for Hepatocytes**

DESCRIPTION	QTY./CASE	CAT. NO
6-well plates	5	<a href="#">62405-144</a>

**BD BioCoat™ Matrigel™ Matrix Plates for Embryonic Stem Cell Culture**

DESCRIPTION	QTY./CASE	CAT. NO
6-well plates	5	<a href="#">89022-350</a>

**BD BioCoat™ Poly-D-Lysine/Laminin Cellware**

DESCRIPTION	QTY./CASE	CAT. NO
6-well plates	5	<a href="#">62405-109</a>
24-well plates	5	<a href="#">62405-111</a>
96-well plates	5	<a href="#">62405-113</a>
100 mm culture dishes	10	<a href="#">62405-115</a>
Coverslips 12 mm round No.1 German glass	80	<a href="#">BD354807</a>
2-well CultureSlides	12	<a href="#">BD354687</a>
8-well CultureSlides	12	<a href="#">BD354688</a>

**BD BioCoat™ Poly-L-Ornithine/Laminin Cellware**

DESCRIPTION	QTY./CASE	CAT. NO
6-well plates	5	<a href="#">62405-434</a>
24-well plates	5	<a href="#">62405-436</a>
96-well plates	5	<a href="#">62405-438</a>

**BD Falcon™ Cultureware**

	DESCRIPTION	QTY./CASE	CAT. NO.
1-well CultureSlides	8.6 cm <sup>2</sup> growth surface area per well	96	<a href="#">62405-172</a>
		24	<a href="#">53106-300</a>
2-well CultureSlides	4.0 cm <sup>2</sup> growth surface area per well	96	<a href="#">62405-174</a>
		24	<a href="#">53106-302</a>
4-well CultureSlides	1.7 cm <sup>2</sup> growth surface area per well	96	<a href="#">62405-176</a>
		24	<a href="#">53106-304</a>
8-well CultureSlides	0.7 cm <sup>2</sup> growth surface area per well	96	<a href="#">62405-178</a>
		24	<a href="#">53106-306</a>
BD Primaria™ Cell Culture Dishes with lid	35x10 mm style	200	<a href="#">25382-654</a>
	60x15 mm style	200	<a href="#">25382-687</a>
	100x20 mm style	200	<a href="#">25382-701</a>
BD Primaria™ Cell Culture Flasks with plug-seal screw cap	25 cm <sup>2</sup> growth area, 50 ml, canted neck	200	<a href="#">29184-801</a>
	75 cm <sup>2</sup> growth area, 250 ml, straight neck	100	<a href="#">29184-845</a>
BD Primaria™ Cell Culture Flasks with 0.2 µm membrane vented screw cap	25 cm <sup>2</sup> growth area, 50 ml, canted neck	100	<a href="#">15705-068</a>
	75 cm <sup>2</sup> growth area, 250 ml, straight neck	100	<a href="#">15705-070</a>
BD Primaria™ Cell Culture Plates, flat-bottom with lid	6-well	50	<a href="#">62406-455</a>
	24-well	50	<a href="#">62406-456</a>
	96-well	50	<a href="#">62406-503</a>
96-well Imaging Plate	Black/Clear, with lid	32	<a href="#">89022-352</a>

## Cell Environments

	DESCRIPTION	QTY./CASE	CAT. NO
BD BioCoat™ Tumor Invasion System	One insert plate with one 24-well plate and lid	1	<a href="#">47733-148</a>
	Five insert plates with five 24-well plates and lids	5	<a href="#">47733-146</a>
BD BioCoat Angiogenesis System: Endothelial Cell Invasion	One insert plate with one 24-well plate and lid	1	<a href="#">BD354141</a>
	Five insert plates with five 24-well plates and lids	5	<a href="#">BD354142</a>
BD BioCoat Angiogenesis System: Endothelial Cell Migration	One insert plate with one 24-well plate and lid	1	<a href="#">47743-788</a>
	Five insert plates with five 24-well plates and lids	5	<a href="#">47743-790</a>
	One insert plate with one 96-well plate and lid	1	<a href="#">47747-300</a>
	Five insert plates with five 96-well plates and lids	5	<a href="#">47747-302</a>
BD BioCoat Angiogenesis System: Endothelial Tube Formation	96-well Black/Clear BD Optilux™ Microplate	1	<a href="#">47745-226</a>
	96-well Black/Clear Optilux Microplate	5	<a href="#">47745-228</a>
BD BioCoat BD Matrigel™ Invasion Chambers	8 µm inserts in four 6-well plates	24	<a href="#">62405-742</a>
	8 µm inserts in two 24-well plates	24	<a href="#">62405-744</a>
BD BioCoat GFR Matrigel Invasion Chambers	8 µm inserts in two 24-well plates	24	<a href="#">62405-158</a>
BD BioCoat Endothelial Cells	Endothelial Cell Growth Environment	1	<a href="#">62405-590</a>

	DESCRIPTION	QTY./CASE	CAT. NO
BD BioCoat Hepatocyte Differentiation	Hepatocyte Differentiation Environment	1	<a href="#">62405-096</a>
BD BioCoat Intestinal Epithelial Differentiation Environment	Intestinal Epithelium Differentiation Environment	1	<a href="#">62405-098</a>
BD BioCoat HTS Caco-2 Assay Systems	1 µm inserts in one 24-Multiwell plate with feeder tray and lid	1	<a href="#">12777-040</a>
	1 µm inserts in one 24-Multiwell plate with feeder tray and lid	5	<a href="#">12777-042</a>
BD BioCoat Fibrillar Collagen 24-Multiwell Insert Systems	1 µm inserts in one 24-Multiwell plate with feeder tray and lid	1	<a href="#">12777-044</a>
	1 µm inserts in one 24-Multiwell plate with feeder tray and lid	5	<a href="#">12777-046</a>



## Membrane Insert Systems

For use with BD Falcon™ Cell Culture Insert Companion Plates

	DESCRIPTION	QTY./CASE	CAT. NO.
0.4 µm, Transparent PET membrane	for 6-well plates	48	<a href="#">62406-163</a>
	for 12-well plates	48	<a href="#">62406-174</a>
	for 24-well plates	48	<a href="#">62404-550</a>
1 µm, Transparent PET membrane	for 6-well plates	48	<a href="#">62406-171</a>
	for 12-well plates	48	<a href="#">62406-172</a>
	for 24-well plates	48	<a href="#">62406-173</a>
3 µm, Transparent PET membrane	for 6-well plates	48	<a href="#">62406-164</a>
	for 12-well plates	48	<a href="#">62406-174</a>
	for 24-well plates	48	<a href="#">62406-169</a>
0.4 µm, HD inserts Translucent PET membrane	for 6-well plates	48	<a href="#">62406-179</a>
	for 12-well plates	48	<a href="#">62406-180</a>
	for 24-well plates	48	<a href="#">62406-181</a>
3 µm HD Inserts, Translucent PET membrane	for 6-well plates	48	<a href="#">62406-170</a>
	for 12-well plates	48	<a href="#">62406-177</a>
	for 24-well plates	48	<a href="#">62406-178</a>
8 µm Translucent PET membrane	for 6-well plates	48	<a href="#">62406-196</a>
	for 12-well plates	48	<a href="#">62406-176</a>
	for 24-well plates	48	<a href="#">62406-198</a>
BD Falcon Cell Culture Insert Companion Plates	6-well plate	50	<a href="#">62406-185</a>
	12-well plate	50	<a href="#">62406-187</a>
	24-well plate	50	<a href="#">62406-189</a>
BD Falcon 24-Multiwell Insert Systems	1 µm PET membrane	1	<a href="#">62406-222</a>
	1 µm PET membrane	5	<a href="#">47727-004</a>
	3 µm PET membrane	1	<a href="#">62406-226</a>
	3 µm PET membrane	5	<a href="#">BD351183</a>
	8 µm PET membrane	1	<a href="#">62406-230</a>
	8 µm PET membrane	5	<a href="#">BD351185</a>
BD Falcon 24-Multiwell Insert Systems	Feeder tray with lid	5	<a href="#">BD351186</a>
BD Falcon FluoroBlok™ 96-Multiwell Insert Systems	3 µm, One insert plate with 96-well plate and lid	1	<a href="#">BD351161</a>
	8 µm, Five insert plates with 96-well plates and lids	5	<a href="#">BD351164</a>
BD Falcon 96-Square Well, Flat-Bottom Plate and Lid	96-square well, flat- bottom plate and lid	5	<a href="#">BD353928</a>

	DESCRIPTION	QTY./CASE	CAT. NO.
BD BioCoat™ Collagen I Cell Culture Inserts	0.4 µm inserts in four 6-well plates	24	<a href="#">62405-626</a>
	0.4 µm inserts in four 12-well plates	24	<a href="#">62405-540</a>
	0.4 µm inserts in two 24-well plates	24	<a href="#">62405-628</a>
	1 µm inserts in four 6-well plates	24	<a href="#">62405-542</a>
	1 µm inserts in four 12-well plates	24	<a href="#">62405-544</a>
	1 µm inserts in two 24-well plates	24	<a href="#">62405-546</a>
	3 µm inserts in four 6-well plates	24	<a href="#">62405-630</a>
	3 µm inserts in four 12-well plates	24	<a href="#">62405-634</a>
	3 µm inserts in two 24-well plates	24	<a href="#">62405-632</a>
BD BioCoat Collagen IV Cell Culture Inserts	1 µm inserts in two 24-well plates	24	<a href="#">62405-554</a>
	3 µm inserts in four 6-well plates	24	<a href="#">62405-654</a>
	3 µm inserts in two 24-well plates	24	<a href="#">62405-656</a>
BD BioCoat Fibrillar Collagen Cell Culture Inserts	1 µm inserts in four 6-well plates	24	<a href="#">62405-119</a>
	1 µm inserts in four 12-well plates	24	<a href="#">62405-121</a>
	1 µm inserts in two 24-well plates	24	<a href="#">62405-050</a>
BD BioCoat Fibronectin Cell Culture Inserts	0.4 µm inserts in four 6-well plates	24	<a href="#">62405-686</a>
	0.4 µm inserts in two 24-well plates	24	<a href="#">62405-688</a>
	1 µm inserts in two 24-well plates	24	<a href="#">62405-566</a>
	3 µm inserts in four 12-well plates	24	<a href="#">62405-568</a>
	3 µm inserts in two 24-well plates	24	<a href="#">62405-655</a>
BD BioCoat FluoroBlok Fibronectin Cell Culture Inserts	3 µm inserts in two 24-well plates	24	<a href="#">BD354597</a>
BD BioCoat Collagen I 24-Multiwell Insert System	3 µm insert plate with 24-well plate and lid	1	<a href="#">BD354598</a>
BD BioCoat™ Fibronectin 24-Multiwell Insert System	3 µm insert plate with 24-well plate and lid	1	<a href="#">BD354599</a>

## Membrane Insert Systems (continued)

	DESCRIPTION	QTY./CASE	CAT. NO
BD BioCoat™ Collagen I Cell Culture Inserts	0.4 µm inserts in four 6-well plates	24	<a href="#">62405-626</a>
	0.4 µm inserts in four 12-well plates	24	<a href="#">62405-540</a>
	0.4 µm inserts in two 24-well plates	24	<a href="#">62405-628</a>
	1 µm inserts in four 6-well plates	24	<a href="#">62405-542</a>
	1 µm inserts in four 12-well plates	24	<a href="#">62405-544</a>
	1 µm inserts in two 24-well plates	24	<a href="#">62405-546</a>
	3 µm inserts in four 6-well plates	24	<a href="#">62405-630</a>
	3 µm inserts in four 12-well plates	24	<a href="#">62405-634</a>
	3 µm inserts in two 24-well plates	24	<a href="#">62405-632</a>
BD BioCoat Collagen IV Cell Culture Inserts	1 µm inserts in two 24-well plates	24	<a href="#">62405-554</a>
	3 µm inserts in four 6-well plates	24	<a href="#">62405-654</a>
	3 µm inserts in two 24-well plates	24	<a href="#">62405-656</a>
BD BioCoat Fibrillar Collagen Cell Culture Inserts	1 µm inserts in four 6-well plates	24	<a href="#">62405-119</a>
	1 µm inserts in four 12-well plates	24	<a href="#">62405-121</a>
	1 µm inserts in two 24-well plates	24	<a href="#">62405-050</a>
BD BioCoat Fibronectin Cell Culture Inserts	0.4 µm inserts in four 6-well plates	24	<a href="#">62405-686</a>
BD BioCoat FluoroBlok™ Fibronectin Cell Culture Inserts	3.0 µm inserts in two 24-well plates	24	BD354597
BD BioCoat Collagen I 24-Multiwell Insert System	3.0 µm insert plate with 24-well plate and lid	1	BD354598
BD BioCoat Fibronectin 24-Multiwell Insert System	3.0 µm insert plate with 24-well plate and lid	1	BD354599
BD BioCoat Laminin Cell Culture Inserts	0.4 µm inserts in two 24-well plates	24	62405-718
	3 µm inserts in two 24-well plates	24	<a href="#">62405-722</a>

	DESCRIPTION	QTY./CASE	CAT. NO
BD BioCoat Matrigel™ Matrix Cell Culture Inserts	0.4 µm inserts in four 6-well plates	24	<a href="#">62405-734</a>
	0.4 µm inserts in two 24-well plates	24	<a href="#">62405-736</a>
BD BioCoat Control Cell Culture Inserts	0.4 µm inserts in four 6-well plates	24	<a href="#">62405-020</a>
	0.4 µm inserts in four 12-well plates	24	<a href="#">62405-018</a>
	0.4 µm inserts in two 24-well plates	24	<a href="#">62405-016</a>
	1 µm inserts in four 6-well plates	24	<a href="#">62405-014</a>
	1 µm inserts in four 12-well plates	24	<a href="#">62405-012</a>
	1 µm inserts in two 24-well plates	24	<a href="#">62405-010</a>
	3 µm inserts in four 6-well plates	24	<a href="#">62405-008</a>
	3 µm inserts in four 12-well plates	24	<a href="#">62405-006</a>
	3 µm inserts in two 24-well plates	24	<a href="#">62405-004</a>
	8 µm inserts in four 6-well plates	24	<a href="#">62405-002</a>
	8 µm inserts in two 24-well plates	24	<a href="#">62405-000</a>
BD Falcon™ FluoroBlok Cell Culture Inserts For use with BD Falcon 24-well Cell Culture Insert Companion Plates (Cat. No. 353504)	1 µm inserts	48	<a href="#">62406-498</a>
	3 µm inserts	48	<a href="#">62406-500</a>
	8 µm inserts	48	<a href="#">62406-504</a>
	1 µm insert system in one 24-well plate	5	<a href="#">BD351154</a>
	3 µm insert system in one 24-well plate	1	<a href="#">BD351155</a>
	3 µm insert system in one 24-well plate	5	<a href="#">BD351156</a>
	8 µm insert system in one 24-well plate	1	<a href="#">BD351157</a>
	8 µm insert system in one 24-well plate	5	<a href="#">BD351158</a>
BD BioCoat Deep-Well Plates For use with BD BioCoat Cell Culture Inserts	6-well Deep-Well Plates	4	<a href="#">62405-150</a>

# References

## Human Embryonic Stem Cells

1. Xu C, Inokuma MS, Denham J, Golds K, Kundu P, Gold JD, Carpenter MK. (2001) Feeder-free growth of undifferentiated human embryonic stem cells. *Nat Biotechnol.* 19:971.
2. Amit M, Shariki C, Margulets V, Itskovitz-Eldor J. (2004) Feeder layer- and serum-free culture of human embryonic stem cells. *Biol Reprod.* 70:837.
3. Ludwig TE, Bergendahl V, Levenstein ME, Yu J, Probasco MD, Thomson JA. (2006) Feeder-independent culture of human embryonic stem cells. *Nat Methods.* 3(8):637.
4. Ludwig TE, Levenstein ME, Jones JM, Berggren WT, Mitchen ER, Frane JL, Crandall LJ, Daigh CA, Conard KR, Piekarczyk MS, Llanas RA, Thomson JA. (2006) Derivation of human embryonic stem cells in defined conditions. *Nat Biotechnol.* 24(2):185.
5. Takahashi K, Tanabe K, Ohnuki M, Narita M, Ichisaka T, Tomoda K, Yamanaka S. (2007) Induction of pluripotent stem cells from adult human fibroblasts by defined factors. *Cell.* 131:1.
6. Yu J, Vodyanik MA, Smuga-Otto K, Antosiewicz-Bourget J, Frane JL, Tian S, Nie J, Jonsdottir GA, Ruotti V, Stewart R, Slukvin II, Thomson JA. (2007) Induced pluripotent stem cell lines derived from human somatic cells. *Science.* 318(5858):1917.

## Endothelial Cells

7. Nakamura K, Taguchi E, Miura T, Yamamoto A, Takahashi K, Bichat F, Guilbaud N, Hasegawa K, Kubo K, Fujiwara Y, Suzuki R, Kubo K, Shibuya M, Ise T. (2006) KRN951, a highly potent inhibitor of vascular endothelial growth factor receptor tyrosine kinases, has antitumor activities and affects functional vascular properties. *Cancer Res.* 66(18):9134.
8. Steinle JJ, Booz GW, Meininger CJ, Day JNE, Granger HJ. (2003)  $\beta_2$ -adrenergic receptors regulate retinal endothelial cell migration and proliferation. *J Biol Chem.* 278(23):20681.
9. Di Simone N, De Santis M, Tamburrini E, Di Nicuolo F, Lucia MB, Riccardi P, D'Ipollito S, Cauda R, Caruso A. (2007) Effects of antiretroviral therapy on tube-like network formation of human endothelial cells. *Biol Pharm Bull.* 30(5):982.
10. Kong D, Li Y, Wang Z, Banerjee S, Sarkar FH. (2007) Inhibition of angiogenesis and invasion by 3,3'-diindolylmethane is mediated by the NF- $\kappa$ B downstream target genes *MMP-9* and *uPA* that regulated bioavailability of VEGF in prostate cancer. *Cancer Res.* 67(7):3310.
11. Michaud-Levesque J, Demeule M, Beliveau R. (2007) *In vivo* inhibition of angiogenesis by a soluble form of melanotransferrin. *Carcinogenesis.* 28(2):280.
12. Nishiyama K, Takaji K, Uchijima Y, Kurihara Y, Asano T, Yoshimura M, Ogawa H, Kurihara H. (2007) Protein kinase A-regulated nucleocytoplasmic shuttling of Id1 during angiogenesis. *J Biol Chem.* 282(23):17200.
13. Takeda Y, Kazarov AR, Butterfield CE, Hopkins BD, Benjamin LE, Kaipainen A, Hemler ME. (2007) Deletion of tetraspanin *Cd51* results in decreased pathologic angiogenesis *in vivo* and *in vitro*. *Blood.* 109(4):1524.
14. Di Simone N, Di Nicuolo F, Sanguinetti M, Castellani R, D'Asta M, Caforio L, Caruso A. (2006) Resistin regulates human choriocarcinoma cell invasive behaviour and endothelial cell angiogenic processes. *J Endocrinol.* 189:691.
15. Folkman J and Haudenschild C. (1980) Angiogenesis *in vitro*. *Nature.* 288(5791):551.
16. Birdsey GM, Dryden NH, Amsellem V, Gebhardt F, Sahnan K, Haskard DO, Dejana E, Mason JC, Rand AM. (2008) Transcription factor Erg regulates angiogenesis and endothelial apoptosis through VE-cadherin. *Blood.* 111(7):3498.
17. Murphy EZ, Majeti BK, Barnes LA, Makale M, Weis SM, Lutu-Fuga K, Wrasidlo W, Cheresch DA. (2008) Nanoparticle-mediated drug delivery to tumor vasculature suppresses metastasis. *Proc Natl Acad Sci.* 105(27):9343.
18. Kisuck J, Butterfield CE, Duda DG, Eichenberger SC, Saffaripour S, Ware J, Ruggeri ZM, Jain RK, Folkman J, Wagner DD. (2006) Platelets and platelet adhesion support angiogenesis while preventing excessive hemorrhage. *Proc Natl Acad Sci.* 103(4):855.

## Hepatocytes

19. Fahmi OA, Boldt S, Kish M, Obach RS, Tremaine LM. (2008) Prediction of drug-drug interactions from *in vitro* induction data: application of the relative induction score approach using cryopreserved human hepatocytes. *Drug Metab Dispos.* 36(9):1971.
20. Healan-Greenberg C, Waring JF, Kempf DJ, Blomme EA, Tirona RG, Kim RB. (2008) A human immunodeficiency virus protease inhibitor is a novel functional inhibitor of human pregnane X receptor. *Drug Metab Dispos.* 36(3):500.
21. Lee P, Peng H, Gelbart T, Beutler E. (2004) The IL-6- and lipopolysaccharide-induced transcription of hepcidin in HFE-, transferrin receptor 2-, and  $\beta_2$  microglobulin-deficient hepatocytes. *Proc Natl Acad Sci.* 101(25):9263.
22. DiPersio CM, Jackson DA, Zaret KS. (1991) The extracellular matrix coordinately modulates liver transcription factors and hepatocyte morphology. *Mol Cell Biol.* 11(9):4405.

23. Rana B, Mischoulon D, Xie Y, Bucher NL, Farmer SR. (1994) Cell-extracellular matrix interactions can regulate the switch between growth and differentiation in rat hepatocytes: reciprocal expression of C/EBP alpha and immediate-early growth response transcription factors. *Mol Cell Biol.* 14(9):5858.
24. Schuetz EG, Li D, Omiecinski CJ, Muller-Eberhard U, Kleinman HK, Elswick B, Guzelian PS. (1988) Regulation of gene expression in adult rat hepatocytes cultured on a basement membrane matrix. *J Cell Physiol.* 134:309.
25. Schuetz JD and Schuetz EG. (1993) Extracellular matrix regulation of multidrug resistance in primary monolayer cultures of adult rat hepatocytes. *Cell Growth and Diff.* 4:31.
26. Mann DJ, Strain AJ, Bailey E. (1992) Hormonal induction of malic enzyme in rat hepatocytes cultured on laminin-rich gels. *J Mol Endocrinol.* 8(3):235.
27. Kane RE, Tector J, Brems JJ, Li A, Kaminski D. (1991) Sulfation and glucuronidation of acetaminophen by cultured hepatocytes reproducing *in vivo* sex-differences in conjugation on Matrigel and type 1 collagen. *In Vitro Cell Dev Biol.* 27A:953.
28. Wang S, Nagrath D, Chen PC, Berthiaume F, Yarmush ML. (2008) Three-dimensional primary hepatocyte culture in synthetic self-assembling peptide hydrogel. *Tissue Eng.* 14(2):227.
29. Semino CE, Merok JR, Crane GG, Panagiotakos G, Zhang S. (2003) Functional differentiation of hepatocyte-like spheroid structures from putative liver progenitor cells in three-dimensional peptide scaffolds. *Differentiation.* 71(4-5):262.
30. Dike LE, Haiyan X, Snodgrass BR, Patten CJ. (2006) Characteristics of replatable and inducible cryopreserved hepatocytes. Poster presented at the 14th North American ISSX Meeting, Rio Grande, PR, Poster No. 162.
31. Weng Y, Stresser DM, Zhang JG. (2005) Characterization of CYP1A2, 2B6 and 3A4 induction in primary cultures of human hepatocytes by RT-PCR, Enzyme Activity and Western Blot Poster presented at the 8th International ISSX Meeting, Sendai, Japan.
32. Bi YA, Kazilias D, Duignan DB. (2006) Use of cryopreserved human hepatocytes in sandwich culture to measure hepatobiliary transport. *Drug Metab Dispos.* 34(9):1658.

## Neuronal Cells

33. Willard MD, Willard FS, Li X, Cappell SD, Snider WD, Siderovski DP. (2007) Selective role for RGS12 as a Ras/Raf/MEK scaffold in nerve growth factor-mediated differentiation. *EMBO J.* 26:2029.
34. Rosario M, Franke R, Bednarski C, Birchmeier W. (2007) The neurite outgrowth multidaptor RhoGAP, NOMA-GAP, regulates neurite extension through SHP2 and Cdc42. *J Cell Biol.* 178(3):503.
35. Wetzel M, Li L, Harms KM, Roitbak T, Ventura PB, Rosenberg GA, Khokha R, Cunningham LA. (2008) Tissue inhibitor of metalloproteinases-3 facilitates Fas-mediated neuronal cell death following mild ischemia. *Cell Death Diff.* 15:143-151.
36. Schnitzler AC, Lopez-Coviella I, Blusztajn JK. (2008) Purification and culture of nerve growth factor receptor (p75)-expressing basal forebrain cholinergic neurons. *Nat Protocols.* 3(1):34.
37. Redmond Jr DE, Bjurgstad KB, Teng YD, Ourednik V, Ourednik J, Wakeman DR, Parsons XH, Gonzalez R, Blanchard BC, Kim SU, Gu Z, Lipton SA, Markakis EA, Roth RH, Elsworth JD, Sladek Jr JR, Sidman RL, Snyder EY. (2007) Behavioral improvement in a primate Parkinson's model is associated with multiple homeostatic effects of human neural stem cells. *Proc Natl Acad Sci.* 104(29):12175.
38. Thonhoff JR, Lou DI, Jordan PM, Zhao X, Wu P. (2008) Compatibility of human fetal neural stem cells with hydrogel biomaterials *in vitro*. *Brain Res.* 1187:42.
39. Gelain F, Bottai D, Vecsoki A, Zhang S. (2006) Designer self-assembling peptide nanofiber scaffolds for adult mouse neural stem cell 3-dimensional cultures. *PLoS ONE.* 1(1):e119.
40. Aguirre A, Rizvi TA, Ratner N, Gallo V. (2005) Overexpression of the epidermal growth factor receptor confers migratory properties to nonmigratory postnatal neural progenitors. *J Neurosci.* 25(48):11092.
41. Holgado-Madruga M, Moscatello DK, Emlet DR, Dieterich R, Wong AJ. (1997) Grb2-associated binder-1 mediates phosphatidylinositol 3-kinase activation and the promotion of cell survival by nerve growth factor. *Proc Natl Acad Sci.* 94:12419.
42. Dimos JT, Rodolfo KT, Nakiak KK, Weisenhalt LM, Mitumoto H, Chung W, Croft GF, Saphier G, Leibel R, Goland R, Wichterle H, Henderson CE, Eggan K. (2008) Induced pluripotent stem cells generated from patients with ALS can be differentiated into motor neurons. *Science.* 321:1218.
43. Watanabe K, Ueno M, Kamiya D, Nishiyama A, Matsumura M, Wataya T, Takahashi JB, Nishikawa S, Nishikawa SI, Mugiura K, Sasaki Y. (2007) A ROCK inhibitor permits survival of dissociated human embryonic stem cells. *Nat Biotechnol.* 25:681.
44. Flanagan LA, Rebaza LM, Derzic S, Schwartz PH, Monuki ES. (2006) Regulation of human neural precursor cells by laminin and integrins. *J Neurosci Res.* 83(5):845.



## Epithelial Cells

45. Tscharrnke M, Pofahl R, Chrostek-Grashoff A, Smyth N, Niessen C, Niemann C, Hartwig B, Herzog V, Klein HW, Krieg T, Brakebusch C, Haase I. (2007) Impaired epidermal wound healing *in vivo* upon inhibition or deletion of Rac1. *J Cell Sci.* 120:1480.

46. Grose R, Hutter C, Bloch W, Thorey I, Watt FM, Fassler R, Brakebusch, Werner S. (2002) A crucial role of  $\beta 1$  integrins for keratinocyte migration *in vitro* and during cutaneous wound repair. *Development.* 129:2303.
47. Shenoy SK, Xizo K, Venkataramanan V, Snyder PM, Freedman NJ, Weissman AM. (2008) Nedd4 mediates agonist-dependent ubiquitination, lysosomal targeting, and degradation of the  $\beta_2$ -adrenergic receptor. *J Biol Chem.* 283(3):22166.
48. Madras BK, Xie Z, Lin Z, Jassen A, Panas H, Lynch L, Johnson R, Livni E, Spencer TJ, Bonab AA, Miller GM, Fischman AJ. (2006) Modafinil occupies dopamine and norepinephrine transporters *in vivo* and modulates the morphologies and trace amine activity *in vitro*. *J Pharmacol Exp Ther.* 319:561.
49. Shenoy SK, Drake MT, Nelson CD, Houtz DA, Xiao K, Madabushi S, Reiter E, Premont RT, Lichtarge O, Lefkowitz RJ. (2006)  $\beta$ -arrestin-dependent, G protein-independent ERK1/2 activation by the  $\beta_2$  adrenergic receptor. *J Biol Chem.* 281(2):1261.
50. Nelson CM, Inman JL, Bissell MJ. (2008) Three-dimensional lithographically defined organotypic tissue arrays for quantitative analysis of morphogenesis and neoplastic progression. *Nat Protocol.* 3(4):674.
51. Itoh M, Nelson CM, Myers CA, Bissell MJ. (2007) Rap1 integrates tissue polarity, lumen formation, and tumorigenic potential in human breast epithelial cells. *Cancer Res.* 67(10):4759.
52. Kenny PA, Lee GY, Myers CA, Neve RM, Semeiks JR, Spellman PT, Lorenz K, Lee EH, Barcellos-Hoff MH, Petersen OW, Gray JW, Bissell MJ. (2007) The morphologies of breast cancer cell lines in three-dimensional assays correlate with their profiles of gene expression. *Mol Oncol.* 1(1):84.
53. Friedland JC, Lakins JN, Kazanietz MG, Chernoff J, Boettiger D, Weaver VM. (2007)  $\alpha 6 \beta 4$  integrin activates Rac-dependent p21-activated kinase 1 to drive NF- $\kappa$ B-dependent resistance to apoptosis in 3D mammary acini. *J Cell Sci.* 120: 3700.
54. Debnath J, Muthuswamy SK, Brugge JS. (2003) Morphogenesis and oncogenesis of MCF-10A mammary epithelial acini grown in three-dimensional basement membrane cultures. *Methods.* 30(3):256.
55. Shirasaka Y, Kawasaki M, Sakane T, Omatsu H, Moriya Y, Nakamura T, Sakaeda T, Okumura K, Langguth P, Yamashita S. (2006) Induction of human p-glycoprotein in Caco-2 cells: development of a highly sensitive assay system for p-glycoprotein-mediated drug transport. *Drug Metab. Pharmacokinet.* 21(5):414.
56. Yamashita S, Konishi K, Yamazaki Y, Taki Y, Sakane T, Sezaki, H, Furuyama Y. (2002) New and better protocols for a short-term Caco-2 cell culture system. *J Pharma Sci.* 91(3):669.

## Tumor Cells

57. Albini A and Benelli R. (2007) The chemoinvasion assay: a method to assess tumor and endothelial cell invasion and its modulation. *Nat Protocols.* 2(3):505.
58. Oxelmark E, Roth JM, Brooks PC, Braunstein SE, Schneider RJ, Garabedian MJ. (2006) The coxapone p23 differentially regulates estrogen receptor target genes and promotes tumor cell adhesion and invasion. *Mol Cell Biol.* 26(14):5205.
59. Duxbury MA, Ito H, Zinner MJ, Ashley SW, Whang EE. (2004) EphA2: a determinant of malignant cellular behavior and a potential therapeutic target in pancreatic adenocarcinoma. *Oncogene.* 23:1448.
60. Seton-Regers SE, Lu Y, Hines LM, Koundinya M, LaBaer J, Muthuswamy SK, Brugge JS. (2004) Cooperation of the ErbB2 receptor and transforming growth factor  $\beta$  in induction of migration and invasion in mammary epithelial cells. *Proc Natl Acad Sci.* 101(5):1257.
61. Singh A, Singh UP, Grizzle WE, Lillard Jr JW. (2004) CXCL12-CXCR4 interactions modulate prostate cancer cell migration, metalloproteinase expression and invasion. *Lab Invest.* 84:1666.
62. Takada Y, Kobayashi Y, Aggarwal BB. (2005) Evodiamine abolishes constitutive and inducible NF- $\kappa$ B activation by inhibiting I $\kappa$ B $\alpha$  kinase activation, thereby suppressing NF- $\kappa$ B-regulated antiapoptotic and metastatic gene expression, up-regulating apoptosis, and inhibiting invasion. *J Biol Chem.* 280(17):17203.
63. Ichikawa H, Takada Y, Murakami A, Aggarwal BB. (2005) Identification of a novel blocker of I kappa B alpha kinase that enhances cellular apoptosis and inhibits cellular invasion through suppression of NF-kappa B-regulated gene products. *J Immunol.* 174(11):7383.
64. Cho RW, Wang X, Diehn M, Shedden K, Chen GY, Sherlock G, Gurney A, Lewicki J, Clarke MF. (2008) Isolation and molecular characterization of cancer stem cells in MMTV-Wnt-1 murine breast tumors. *Stem Cells.* 26(2):364.
65. Purhonen S, Palm J, Rossi D, Kaskenpaa N, Rajantie J, Yla-Herttuala S, Alitalo K, Weissman IL, Salven P. (2008) Bone marrow-derived circulating endothelial precursors do not contribute to vascular endothelium and are not needed for tumor growth. *Proc Natl Acad Sci.* 105(18):6620.
66. Feldmann G, Dhara S, Fendrich V, Bedja D, Beatty R, Mullendore M, Karikari C, Alvarez H, Iacobuzio-Donahue C, Jimeno A, Gabrielson KL, Matsui VW, Matra A. (2007) Blockade of hedgehog signaling inhibits pancreatic cancer invasion and metastases: a new paradigm for combination therapy in solid cancers. *Cancer Res.* 67:2187.



Supplier Partnerships for Customer Solutions

1.800.932.5000

**vwr.com**

Prices, product appearance and specifications are current at the time of printing, subject to change without notice. Availability for certain products may be limited by federal, state, provincial or local licensing requirements. All prices are in U.S. dollars unless otherwise noted. Offers valid in USA and Canada, void where prohibited by law or company policy, while supplies last. Visit [vwr.com](http://vwr.com) to view our privacy policy and additional disclaimers.

VWR, forms of VWR and the VWR logo and/or design are either registered trademarks ®, trademarks ™, or service marks ™ of VWR International in the United States and/or other countries. All other marks referenced herein are registered trademarks, trademarks or service marks of their respective owner(s). For a complete list of trademark owners, please visit [vwr.com](http://vwr.com)

©2008 VWR International, LLC. All rights reserved.

AlexaFluor is a trademark of Invitrogen Corporation.  
 CytoFluor is a trademark of Applied Biosystems.  
 MetaMorph is a registered trademark of Universal Imaging Corporation (UIC).  
 mlesR1 and ViCell is the property of ViCell Research Institute.  
 PuraMatrix is a registered trademark of 3DM, Inc.  
 Victor2 is a trademark of PerkinElmer, Inc.  
 VWR logo is the property of VWR International.  
 BD, BD logo, and all other trademarks are the property of Becton, Dickinson and Company. ©2008  
 B08G148

1208 5M Lit. No. 02088