

Review

Autophagy and Its Possible Roles in Nervous System Diseases, Damage and Repair

David C. Rubinsztein^{1,*}

Marian DiFiglia²

Nathaniel Heintz³

Ralph A. Nixon^{4,5}

Zheng-Hong Qin²

Brinda Ravikumar¹

Leonidas Stefanis⁶

Aviva Tolkovsky⁷

¹Departments of Medical Genetics; University of Cambridge; Cambridge Institute for Medical Research; Wellcome Trust/MRC Building; Addenbrooke's Hospital; Hills Road, Cambridge UK

²Laboratory of Cellular Neurobiology; Massachusetts General Hospital and Harvard Medical School; Charlestown, Massachusetts USA

³Howard Hughes Medical Institute; Laboratory of Molecular Biology; The Rockefeller University; New York, New York USA

⁴Center for Dementia Research; Nathan S. Kline Institute; Orangeburg, New York USA

⁵Departments of Psychiatry and Cell Biology; New York University; New York, New York USA

⁶Neurobiology Laboratory; Foundation of Biomedical Research; Academy of Athens; Athens Greece

⁷Department of Biochemistry; University of Cambridge; Cambridge UK

*Correspondence to: David C. Rubinsztein; Departments of Medical Genetics; University of Cambridge; Cambridge Institute for Medical Research; Wellcome Trust/MRC Building; Addenbrooke's Hospital; Hills Road, Cambridge CB2 2XY UK; Tel.: +44.1223.762608; Fax: +44.1223.331206; Email: dcr1000@cus.cam.ac.uk

Received 12/15/04; Accepted 12/30/04

Previously published online as an *Autophagy* E-publication:
<http://www.landesbioscience.com/journals/autophagy/abstract.php?id=1513>

KEY WORDS

autophagy, Alzheimer's disease, Parkinson's disease, Huntington's disease, prion, α -synuclein, amyloid, apoptosis

ACKNOWLEDGEMENTS

See page 19.

ABSTRACT

Increased numbers of autophagosomes/autophagic vacuoles are seen in a variety of physiological and pathological states in the nervous system. In many cases, it is unclear if this phenomenon is the result of increased autophagic activity or decreased autophagosome-lysosome fusion. The functional significance of autophagy and its relationship to cell death in the nervous system is also poorly understood. In this review, we have considered these issues in the context of acute neuronal injury and a range of chronic neurodegenerative conditions, including the Lurcher mouse, Alzheimer's, Parkinson's, Huntington's and prion diseases. While many issues remain unresolved, these conditions raise the possibility that autophagy can have either deleterious or protective effects depending on the specific situation and stage in the pathological process.

INTRODUCTION

The ubiquitin-proteasome and autophagy-lysosome pathways are the two major routes for protein and organelle clearance in eukaryotic cells. Proteasomes predominantly degrade short-lived nuclear and cytosolic proteins. Poly-ubiquitination generally serves as a signal for proteasomal substrate recognition.¹ Since substrates need to be unfolded to pass through the narrow pore of the proteasome barrel, this precludes clearance of large membrane proteins and protein complexes (including oligomers and aggregates). Mammalian lysosomes, on the other hand can degrade substrates like protein complexes and organelles.² The bulk degradation of cytoplasmic proteins or organelles is largely mediated by macroautophagy, generally referred to as autophagy. It involves the formation of double membrane structures called autophagosomes/autophagic vacuoles (AVs), which fuse with lysosomes to form autolysosomes (also called autophagolysosomes) where their contents are then degraded by acidic lysosomal hydrolases.³ Autophagosomes are generated by elongation of small membrane structures known as autophagosome precursors. However the precise origin(s) of these structures has yet to be elucidated (Fig. 1).⁴ Autophagy can be induced under physiological stress conditions such as starvation. Several protein kinases regulate autophagy, the best characterized being the mammalian target of rapamycin (mTOR), which negatively regulates the pathway.⁵ While the focus of this review is on macroautophagy, lysosomes can also directly engulf cytoplasm by invagination, protrusion and/or septation of the lysosomal limiting membrane, a process called microautophagy. Another pathway for cytosolic protein clearance via lysosomes is chaperone-mediated autophagy. This pathway is selective for specific cytosolic proteins that contain a pentapeptide motif. This motif is recognized by the chaperone heat shock cognate 70 (Hsc70), which transfers protein substrates to the lysosomal membrane, where, through binding to the receptor lysosome-associated membrane protein 2a (lamp2a), they are translocated into the lysosomal lumen and degraded.⁶

Increased numbers of AVs are a feature of a number of neurological diseases, certain neuronal cell death pathways and after neuronal injury. In many cases it is unclear if this is due to increased autophagic activity (i.e., induction of autophagosome formation) or decreased autophagosome-lysosome fusion (resulting in decreased removal of autophagosomes) (Fig. 2). The role of autophagy when it occurs coincidentally with cell death is also unclear in many cases. If the increased autophagosomes are due to increased autophagic activity, is this directly contributing to the cell death, protecting against cell death, or occurring as an epiphenomenon driven by a pathway perturbed or contributing to the cell death (Fig. 3)?

Here we aim to put some of these questions into perspective in reviewing the current understanding of autophagy and its roles in the nervous system.

AUTOPHAGY AND INJURY

The phenomenon of autophagy in injured neurons was first described many years ago, being especially noted following axotomy or excitotoxicity. Autophagy was also observed after drug-induced block in nerve conduction or axonal transport, to some extent models of axotomy due to paralysis of axonal function (reviewed in ref. 7). Axotomy may cause neuronal cell death, especially in embryonic or neonatal animals, as this is the period when neurons are dependent on their targets for the supply of survival factors. However, in mature animals, where survival factors are derived locally or via autocrine loops, axotomy of peripheral neurons and motoneurons can lead to a robust regenerative response without any neuronal death.⁸⁻¹⁴ In both cases, autophagy has been reported to be markedly enhanced, thus leading to two opposing theories: (1) that autophagy is a mechanism that clears the way for neuronal regeneration or (2) that it is a medium for cell destruction. The notion that autophagy is beneficial to survival is consistent with the evolutionary role of autophagy as a cell-autonomous nutritive mechanism,¹⁵ while the role of autophagy in cell death is consistent with the ability of autophagy to dismantle cellular organelles, especially after stress or injury (a kind of protection as well).¹⁶⁻¹⁸ There is no reason why both roles cannot be performed depending on cell context. Even caspases, normally causal in apoptosis, may be important for cell survival under certain conditions where death by autophagy may prevail.^{19,20}

AUTOPHAGY IN LIVING NEURONS

The increase in autophagic activity after axotomy in living neurons is concomitant with chromatolysis, a stereotypic response where an area of the cytoplasm becomes largely devoid of organelles, especially ribosomes, and fills with numerous vesicles of various types, some of which have been described as autophagosomes.²¹ The autophagic response occurs rapidly. For example, axotomy of superior cervical ganglion (SCG) neurons in adult rats results in increased autophagic activity in the cell body within two hours.¹² In this case, the response peaks within two days, concomitant with the very beginnings of regenerative responses. The contents of the vesicles in the cell bodies include mitochondria and Golgi/ER, classical substrates of autophagy, but many contain dense core vesicles associated with neurotransmitter synthesis. Thus, it was suggested that in the cell body, the autophagic activity clears away those aspects of neuronal function that are not concerned with the production of material for axon re-growth. A surge in vesicle formation, including autophagosomes, was also observed in the proximal stub of the axon that remains connected to the cell body, from whence regeneration




Figure 1. Macroautophagy in mammalian cells. A portion of the cytosol is enclosed by elongation of small membrane structures to form the autophagosomes/autophagic vacuoles (AVs). The outer membranes of AVs then fuse with lysosomes to form autolysosomes where their contents are degraded. Atg12-Atg5-Atg16L localize to the pre-autophagosome membranes, however dissociate upon completion of autophagosome formation. LC3-II (Atg8 homolog) is recruited onto autophagosome membrane throughout the process.




Figure 2. Autophagosome number is a function of synthesis and autophagosome-lysosome fusion/degradation. Increased numbers of autophagosomes can result from induction of autophagy but also from failure of autophagosome fusion with lysosomes.




Figure 3. Possible theoretical relationships between autophagy and cell death, when they occur coincidentally.

will arise.¹³ As mitochondria were found inside many autophagosomes, it was proposed that these need to be cleared before remodeling of the axon could begin. Remarkably, the number of vesicles of various morphologies remained significantly elevated in the neurons too up to 143 days, long after regeneration was nominally complete. The role of this extended reaction, and whether it finally abates, is not clear. In chromatolysis, it was noted that the response was lessened the more distant the insult occurred from the cell. In addition, nerve block was less efficacious at inducing this response compared to nerve cut.²² Hence, the signals that induce autophagy may be linked to the severity of damage, though what these signals are is not known. Because excitotoxicity can under some circumstances also provoke an

increase in autophagy, as detailed below and elsewhere,²³ damage due to pathological effects of calcium may be a common factor. However, though calcium has been implicated in some cases of autophagy^{24,25} no direct causal relationships between calcium-induced signals and injury-induced autophagy have been described, though a calcium-regulated involvement in sequestration of synuclein by autophagic vesicles has been recently inferred²⁶ (see below).

AUTOPHAGY AND CELL DEATH

Newly axotomized neurons from the SCG of neonatal rats are dependent on nerve growth factor (NGF) for their survival. Despite a strong regenerative response, little autophagy was observed in the cell bodies during the first week of culture in the presence of NGF. However, culture without NGF induced a 20–30-fold increase in vesicular content, some of which are autophagosomes,^{27,28} concomitantly with the induction of apoptosis. The vesicle reaction was prolonged if the neurons were not allowed to die by apoptosis (using a pan-caspase inhibitor). This reaction was not due to withdrawal of trophic support (i.e., starvation) since it occurred to similar extents in cell bodies of neurons induced to undergo apoptosis in the presence of NGF (through use of cytosine arabinoside that activates a dominant p53-dependent death pathway).²⁹ Given that loss of mitochondria was observed after a few days of incubation in the absence of NGF,³⁰ and these mitochondria no longer maintained a membrane potential, it was hypothesized that autophagy is responsible for disappearance of functionally-failed mitochondria, as proposed previously.³¹ This idea was strengthened by the observation that bafilomycin A1, a V-type H⁺ ATPase inhibitor that alkalinizes the lysosomal lumen, prevented the loss of mitochondria. However, the mechanisms and origins of these vesicles is still unclear, as rapamycin, the inhibitor of mTOR and an inducer of autophagy in many systems,^{32–34} causes no changes in autophagic response in these neurons in the presence of NGF (though it blocks S6 kinase activity and S6 phosphorylation, Goemans C, unpublished observations) while the commonly used autophagic inhibitor 3-methyladenine, that blocked mitochondrial removal, was found to inhibit not only PI-3 Kinase activity important for autophagosome formation,³⁵ but also the cell death signalling pathway mediated via JNK and p38, and mitochondrial dysfunction.²⁷ Moreover, the mammalian Atg8 homologue MAP-LC3 II, a molecular marker of autophagosomes,³⁶ failed to segregate with any of the vesicles. Thus, whether autophagy plays a causal role in cell death in this model of axotomy is not clear. It may be that GABARAP (gamma 2-subunit of GABA-A receptor associated protein) and GATE-16 (Golgi-associated ATPase Enhancer of 16 kDa), the two other Atg8 homologues, play a role in autophagosome formation in neurons, as at least GABARAP has been described to associate within intracellular membranes such as the Golgi apparatus and postsynaptic cisternae,³⁷ though evidence for its association with autophagic vesicles is still lacking. Interestingly, GABARAP and GATE-16 were recently described as binding partners for ULK1 (Unc-51-like kinase involved in neurite extension) and GABARAP was associated with ULK1 in vesicles in many different classes of neurons.³⁸ The *c. elegans* homologue of ULK-1, Unc-51, which participates in vesicle transport, has high homology to Atg1p, a yeast serine/threonine kinase that is crucial for initiation of autophagic vacuole formation. Whether ULK-1 plays a role in autophagy is unknown.

An interesting model, also invoking autophagy, emerged in a study reporting on the mechanism of death of cultured SCG neurons habituated to survival dependent on glial-derived neurotrophic factor

(GDNF).³⁹ When SCG neurons were isolated from neonatal rats and allowed to regenerate with NGF, they died by apoptosis upon NGF-deprivation. However, when these neurons were regenerated in the presence of GDNF, the neurons die non-apoptotically upon withdrawal of GDNF. Numerous AVs, typically with a double membrane structure, or with multilamellar membranes possibly indicating autolysosomes, appeared in the cytoplasm of neurons concomitantly with death. Interestingly, the mitochondria appeared to remain fully functional, thus divorcing the link between mitochondrial damage and activation of autophagy. Use of dominant negative forms of various caspases implicated caspase 2 and 7 in this death, so there may be a link between caspases and autophagy, though the relationship is not clear.

Clarke has been the strongest proponent of autophagy being involved in developmental cell death of certain neuron populations—defined as “type 2” cell death to distinguish it from “type 1” apoptotic death.⁴⁰ Hornung et al.⁴¹ showed that injection of colchicine to the retina of embryonic chicks (to mimic target deprivation during development, but which also causes direct axonal injury) resulted in the accumulation of numerous vesicles in the dying isthmo-optic nucleus neurons that project to the retina. A range of organelles were found within the vesicles strengthening the notion that these are autophagic. Further recent work from this group has demonstrated autophagic arousal after administration of the glutamate mimetic N-methyl-D-aspartate (NMDA) to rodent hippocampal slices in vitro.⁴² Within 2 hours, numerous vesicles were found in cells that already appear significantly degenerated. The region demarcated by enhanced autophagy was reminiscent of the area “cleared” in chromatolytic neurons and in other neuronal models of excitotoxicity.⁴³ However, only one example of an autophagosome was provided (by EM) that appears as a single membraned structure with cytoplasmic contents. The associated double membraned structure appears to be derived from the adjacent Golgi stack and does not appear to be part of this vesicle. A novel aspect of this work is its link to signalling pathways implicated in cell death. Thus, administration of a JNK-signal blocking peptide derived from the JNK binding domain of JIP1,⁴⁴ a sequence used previously to block neuronal apoptosis in response to NGF-deprivation,⁴⁵ inhibited both the increase in nuclear P-cJun associated with NMDA-induced toxicity, and the subsequent pathological changes including accumulation of AVs. In this model, as well in the model studied previously,⁴¹ death was associated with a massive increase in endocytosis. However, Borsello et al.⁴⁶ have shown that increased endocytic activity—which might have been interpreted as a mechanism for matching the increase in autophagosome number to enable vesicle maturation into an active lysosome is not necessarily part of autophagic death and is also prevalent in a necrotic excitotoxic cell death model induced by kainic acid, which does not involve an increase in autophagy. Perhaps the rapidity of death in this necrotic model precludes activation of autophagy, just as necrosis precludes execution of apoptosis. In any case, the arguments for autophagy being a mediator of death in these models are still indirect. Clark has argued strongly for a type of death executed by autophagy,⁴⁰ and against lysosomal involvement by release of enzymes into the cytoplasm. Studies are now emerging where autophagic genes are shown to participate in apoptotic cell death.^{47,48} However, there are also examples where lysosomal enzymes released from lysosomes participate in controlled forms of cell death.^{49,50} Vacuolisation is also observed in the cytoplasm under both conditions but whether autophagy is involved in this type of death is still not clear. Extensive neuronal vacuolisation is observed,

for example, in the CNS neurons of a mouse in which neuropathy target esterase/swisscheese (normally inhibited by organophosphates) has been deleted but these are unlikely to be autophagic.⁵¹ There is no reason why both mechanisms cannot participate in cell death depending on time and stimulus.

Another form of neuronal injury in which autophagy has been implicated is that induced by metamphetamine.⁵² This type of injury is inflicted specifically on dopaminergic midbrain neurons as the drug forces the intracytoplasmic accumulation of dopamine, a potentially oxidative and toxic catechol neurotransmitter. Treatment results in the accumulation of numerous vesicles described as autophagic which are correlated with loss of mitochondria and neuronal death. Typically, most of the profiles are multilamellar vesicles that could arise from lysosomes without an intermediate step of autophagy. Similar multilamellar profiles were displayed in *C. elegans* neurons dying in response to the hyperactivation of a mechanosensory ion channel, a kind of excitotoxic death.⁵³ Recently this death was found to be inhibited by a series of proteases, including some cysteine proteases typically found in lysosomes.⁵⁴ Whether these become activated as a result of autophagic activity is readily testable, as autophagy has already been dissected genetically in relation to longevity in *C. elegans*.⁵⁵

It is possible that activation of autophagy is important for production of novel forms of hybrid vesicles that then perform specific functions. The importance of interactions of autophagic vesicles with other types of vesicles, in addition to endosomes,^{56,57} should not be dismissed. An interesting potential cross over point is provided by the protein Alix, which is enriched in regions of neurodegeneration induced by the mitochondrial toxin 3-nitropropionic acid and the seizure-inducing excitotoxin kainic acid,^{58,59} and is possibly causal in neuronal cell death.⁶⁰ Alix induces vacuolisation which, together with other factors, creates vesicles displaying multivesicular (that is, several non-concentric vesicles) or multilamellar (concentric vesicle) organisation.⁶¹ Its link with autophagosomes is worth exploring. AVs containing active caspases have also been described in granulo-vacuolar forms of degeneration⁶² but again the identity of these vacuoles as autophagosomes has not been substantiated.

All the occurrences of autophagy in neurons to date have been based on electron microscopy or on use of drugs that are not highly specific as autophagy inducers and inhibitors. The distinction between autophagosomes, lysosomes, endosomes, amphisomes, and their intermediates is not unequivocal. Thus, the importance of autophagy in neuronal death, survival and regeneration awaits application of molecular dissectors that can interfere with this process and distinguish between AVs and other forms of vesicles.

INSIGHTS FROM LURCHER MICE: AUTOPHAGY, NEURONAL FUNCTION AND DYSFUNCTION

One of the strongest associations between autophagy and neurodegeneration is seen in the *Lurcher* mouse, where heterozygotes with an activating mutation in the orphan glutamate receptor (GluR δ 2^{Lc}) develop cerebellar ataxia in the first four postnatal weeks, due to degeneration of the cerebellar cortex. This mutation proved that a molecular alteration in a neurotransmitter receptor can cause degeneration in vivo, and provided the first genetic model in which mechanisms of excitotoxic neuronal death could be intensively investigated in situ.⁶³ Subsequent studies revealed that *lurcher* Purkinje cells activate apoptotic responses in response to this mutation,⁶⁴⁻⁶⁶ although attempts to genetically rescue Purkinje cell death in *lurcher*

mice by blocking the apoptotic pathway were unsuccessful.^{67,68} It is interesting to note that secondary death of granule cells in response to Purkinje cell loss was rescued. These data clearly indicated that, while apoptotic responses may contribute to *Lurcher* pathology, other mechanisms must also come into play.

To try to identify these mechanisms, Yue et al.,²³ characterized a protein complex bound to the C-terminus of the GluR δ 2 receptor, identifying nPIST and Beclin1 as components of the complex. Although Beclin1 was first identified as binding to Bcl2, an inhibitor of apoptosis, it was also immediately recognized as the mammalian orthologue of Atg6 (also known as Vps30), an essential component of the yeast autophagy pathway.⁶⁹ This raised two exciting possibilities: that autophagy was the missing pathway contributing to Purkinje cell death in *lurcher* mice, and that the coupling of autophagy to neurotransmitter receptors plays an important role in regulating neuronal function. To investigate autophagy in the context of Purkinje cell death in *lurcher* mice, Yue et al.,²³ first showed that a fusion of Beclin1 and green fluorescent protein (Beclin1-GFP) could be used as a convenient visual assay for autophagy. They then used this assay to demonstrate that nPIST and Beclin1 could synergize to induce autophagy, that autophagy is induced in heterologous cells transfected with GluR δ 2^{Lc} but not wild type GluR δ 2, and that the death of cells transfected with the mutant receptor could be partially rescued by pharmacologic inhibition of autophagy with 3-methyladenine. However, there is the caveat mentioned earlier that this compound also inhibits death signalling via JNK and p38, and mitochondrial dysfunction.²⁷ Electron microscopic examination of dying *lurcher* Purkinje cells in vivo revealed massive induction of autophagy. These data demonstrated a biochemical link between the neurotransmitter receptor GluR δ 2 and autophagy, and suggested that constitutive activation of autophagy in response to the mutant receptor could lead to cell death. The model that emerged from these studies is that both apoptosis and autophagy can contribute to excitotoxic cell death in vivo, and that Beclin1 is the key molecule orchestrating the relative contributions of these two pathways.

The activation of autophagy in response to signalling through the GluR δ 2^{Lc} receptor raises important issues for models of excitotoxic cell death. For example, one would like to know the relative contributions of ion influx and depolarization of the cell membrane, and excess signalling through receptor-bound complexes to excitotoxic cell death. Insight into this issue in *lurcher* mice has been obtained through studies of the GluR δ 2^{Lc} allele bred onto a hotfoot (a null allele of GluR δ 2) background.⁷⁰ In these mice, Purkinje cell death is accelerated, occurring prior to the stage at which depolarization can be measured in electrophysiologic recordings. Nevertheless, electron microscopic analysis of these cells clearly demonstrates that autophagy is activated. These data provide strong evidence that massive ion flux into the cell and the consequent depolarization of the cell membrane are not critical for GluR δ 2^{Lc} mediated cell death in vivo. The evidence collected from studies of *lurcher* mice, therefore, strongly suggest that constitutive activation of receptor-mediated cell signalling, and the consequent induction of downstream pathways including autophagy, are the critical events that underlie neurodegeneration. While these data strongly couple autophagy to cell death, a direct causal role for autophagy in the cell death execution has yet to be definitively proven in vivo.⁷¹ The fact that the coupling of autophagy to GluR δ 2 (which is expressed principally in cerebellar Purkinje cells) occurs through nPIST and Beclin1 (which are very widely expressed in the nervous system) suggests that this pathway is regulated by a variety of other neurotransmitter receptors and their

ligands. Since synaptic abnormalities have been associated with a wide variety of neurological disorders, the activation of autophagy in these diseases might result from aberrant synaptic signalling events. Although the studies of *lurcher* mice strongly suggest that excess activation of autophagy can contribute to neurodegeneration in response to excitotoxic stimuli, the activation of autophagy via synaptic stress may in many cases first serve a neuro-protective role. Our understanding of the positive and negative consequences of autophagy in the context of neurological disease will be greatly advanced by genetic dissection of this pathway in other animal models of neurological disease.

The most important discovery in these studies of *lurcher* mice is the simple fact that autophagy is directly regulated by a neurotransmitter receptor. Although this has important implications in a disease context and in the context of excitotoxicity, the linking of autophagy and neurotransmitter activity is also likely to be of fundamental importance to normal synaptic function. For example, the GABA receptor associated protein GABARAP⁷² has been shown to be important for trafficking of GABA receptors within the cells.⁷³ Given this result, and the coupling of autophagy to the GluR δ 2 receptor, one might imagine a more general role for autophagy in trafficking and turnover of neurotransmitter receptors at the synapse. This is consistent with studies of Purkinje cell synapses in mice carrying null mutations of GluR δ 2 receptor, where a variety of synaptic abnormalities have been observed.⁷⁴ The contributions of autophagic mechanisms to fundamental mechanisms of synaptic function, including protein trafficking and turnover at the synapse and synaptic remodeling are important areas for future investigation.

ALZHEIMER'S DISEASE AND AUTOPHAGY

Alzheimer's disease (AD), the most prevalent human neurodegenerative disorder, involves a progressive loss of neurons leading to profoundly impaired memory and declines in other intellectual functions. AD is defined by two hallmark neuropathological lesions, the intra-neuronal neurofibrillary tangle containing aggregated forms of the microtubule-associated protein, tau, and the neuritic plaque, mainly composed of extracellular aggregates of β -amyloid peptide (A β) in association with degenerating dendrites or axons.⁷⁵ Preceding cell death, the synaptic terminals, axons and dendrites of neurons undergo a process of degeneration and abortive regeneration over many years,⁷⁶⁻⁸⁰ which creates a highly characteristic pattern of grossly enlarged, "dystrophic" neurites.⁸¹⁻⁸³ Beginning at the earliest stages of this process, the lysosomal system of neurons in affected brain regions is mobilized⁸⁴: lysosomes proliferate, the expression of lysosome-related components increases, and the trafficking of cathepsins to early endosomes is accentuated.⁸⁴⁻⁸⁸ As neurons become more metabolically compromised, lysosomes and other hydrolase-containing compartments progressively accumulate. Indeed, cathepsin D mRNA production remains highly elevated in single neurofibrillary tangle-bearing neurons even though most transcripts unrelated to the lysosomal system decline or remain unchanged.⁸⁹ Increased or altered activity of the endocytic pathway has been considered one possible stimulus for lysosomal system mobilization in AD^{85,90} but macroautophagy is emerging as a key driving force for this and other features of AD neuropathology.

Although rarely in evidence in the normal mature brain, macroautophagy is induced in vulnerable populations of neurons at early stages of AD and in a transgenic mouse model of AD (PS/APP), in which human mutant presenilin-1 (PS-1) and the Swedish variant of

the amyloid precursor protein (APP) are overexpressed.⁹¹⁻⁹³ Mutations in PS-1 and APP cause autosomal dominant, early-onset AD. In both conditions, autophagosomes and other AVs proliferate in dendrites and the levels of LC3-II, a specific marker of autophagosome formation, are elevated.⁹¹⁻⁹³ LC3 in affected brain regions of PS/APP mice translocates to vesicles, reflecting the conversion of LC3-I to LC3-II, and both LC3 isoforms redistribute to dendrites from a mainly perikaryal location. This process begins at young ages before β -amyloid pathology develops in these mice indicating that macroautophagy induction is an early response in the disease and not solely a consequence of amyloid deposition. Autophagy-related pathology in AD and PS/APP mouse brains progresses as autophagosomes and other AV subtypes, together with hydrolase-positive dense bodies,⁹⁴⁻⁹⁶ accumulate in large numbers in grossly distended dystrophic neurites, becoming the principal organelle in these structures. Although AV accumulations are not specific to the degenerative phenomena of AD,^{23,97-99} autophagic-lysosomal pathology in the brain is considerably more widespread and robust in AD than in other adult-onset neurodegenerative diseases.¹⁰⁰ Specifically, the extensive neuritic dystrophy^{101,102} and the characteristic gross distension of these neurites in AD are not typical in other neurodegenerative diseases lacking β -amyloid.¹⁰³ Moreover, the near replacement of cytoplasmic contents in the dystrophic neurites, together with the high incidence of AVs in less obviously affected neurites, constitute a uniquely large "burden" of autophagy-related compartments in the AD brain.

Macroautophagy is normally active within the growing (regenerating) neurites of cultured neurons.¹⁰⁴⁻¹⁰⁶ An early induction of macroautophagy in AD might, therefore, be expected to support the organelle and protein turnover associated with cycles of neurite degeneration and regeneration¹⁰⁷⁻¹⁰⁹ in affected neurons, which also revert to a more immature developmental state in AD.^{110,111} Macroautophagy may also protect neurons in other ways.^{92,54} Activated caspases are detected in the AVs of PS/APP mice and in the granulovacuolar degeneration (GVDs), which are characteristic vacuoles of suspected autophagic origin in hippocampal neurons of AD brain,⁶² raising the possibility that macroautophagy might protect neurons by turning over apoptosis-inducing factors, including damaged mitochondria.¹¹²⁻¹¹⁵

Ultimately, macroautophagy function in dendrites seems gradually to become impaired during the course of AD. In normal growing (regenerating) neurites of cultured neurons, immature AVs are transported retrogradely toward the cell body where they encounter, and fuse with, acidified hydrolase-containing compartments.^{104,105} AV maturation to autophagolysosomes is efficient under these conditions, resulting in the rapid degradation of AV contents, with minimal evidence of intermediate autophagic compartments. The enormous buildup of both early and late AVs within dystrophic/degenerating neurites in AD brain, however, suggests that the transport of AVs may be impeded or that autophagy locally within the dystrophic segments is particularly robust. In addition, the coexistence of both immature and acid-hydrolase-containing compartments in substantial numbers in the dystrophic neurites suggests that immature AVs have access to hydrolase-containing compartments but may less efficiently carry out the final stages of degradation. This possibility is further suggested by observations on the forms of early-onset familial AD (FAD) caused by mutations of presenilin-1 (PS1).¹¹⁶⁻¹¹⁸ PS1 mutations not only induce more severe amyloid pathology and neurodegeneration in PS-FAD and in mouse models but also potentiate lysosomal system pathology.¹¹⁹ Fibroblasts from patients with PS-FAD

also abnormally accumulate AVs, especially under conditions that stimulate autophagy. Surprisingly, however, the turnover of long-lived proteins by autophagy in these fibroblasts is markedly reduced. A direct role for mutant PS1 in this defect is supported by observations that autophagy-mediated protein degradation is nearly eliminated in blastocysts from mice lacking PS1 and PS2 genes but rescued by reintroducing PS1.¹²⁰ Combined with evidence that PS1 and the γ -secretase complex are enriched in AVs,^{93,121} these studies support the hypothesis that PS plays a key role in autophagy and that FAD-linked PS1 mutations have a loss of function effect on macroautophagy leading to AV accumulation and impaired protein turnover.

Because proteasome activity declines in aging and is further impaired in AD,^{122,123} a loss of autophagy function, if it develops in AD, leaves neurons with no competent mechanism to remove abnormal and potentially toxic proteins. Autophagy does not prevent, and may even promote, the overexpression of cathepsins, which continue to accumulate progressively in AD. The accumulated hydrolase-containing AVs in dystrophic neurites represent a large local reservoir of proteases that, if released, have the potential to initiate or mediate aspects of apoptotic and necrotic cell death in various pathological settings.^{84,112,115,124-127}

The relationship of macroautophagy to AD pathogenesis is not limited to neurodegeneration: an intimate connection to β -amyloidogenesis also seems likely. AVs isolated from several different sources are enriched in components required for A β generation and macroautophagy is a significant pathway for A β peptide generation.^{93,121} AVs contain immunoreactive A β peptide and are enriched in its immediate precursor. They are also highly enriched in components of the γ -secretase complex and exhibit high presenilin-dependent γ -secretase activity.^{93,121} Senile plaques in AD and PS/APP mouse brains are also strongly immunoreactive for these components and presenilin-1 localizes prominently to AVs in dystrophic neurites.⁹³ Presenilin and nicastrin, another component of the γ -secretase complex, have been reported to be enriched in lysosomes¹²⁸ although AVs, which were not distinguished from lysosomes in these earlier studies, are a more concentrated localization of these γ -secretase components. Macroautophagy-mediated A β generation has been demonstrated by evidence that A β production rises when autophagy and AV proliferation are stimulated and falls when autophagy is inhibited by blocking either of two independent signalling pathways converging on mTOR kinase.⁹³ Interestingly, inclusion body myositis, the only known condition in which β -amyloid deposits outside the nervous system,¹²⁹ involves the accumulation of autophagy-related "rimmed" vacuoles containing APP, A β , BACE and presenilin.¹³⁰

A β peptide is believed to be generated at several sites within neurons, including endosomes, Golgi, and ER^{86,88,128,131} and each of these organelles could contribute to the involvement of autophagy in A β generation. Endocytic and autophagic pathways communicate extensively and both Golgi and ER are turned over by autophagy.¹³² In normal brain, autophagy may play a relatively minor role in constitutive A β generation; however, in AD, where AVs accumulate in large numbers and persist in dystrophic neurites, autophagy can contribute substantially to overall β -amyloidogenesis, a process promoted in the mildly acidic pH environment of lysosome-related organelles.¹³³ In theory, AV accumulation associated with any significant neuritic injury could stimulate local A β production, as seen for example, in traumatic brain injury.¹³⁴ The chronic and extensive remodeling of neurites in AD, however, represents a particularly favorable condition for A β generation by the autophagic pathway.

As a pathogenically important pathway for A β generation and a mediator of both cell survival and degenerative phenomena,^{40,84,135} macroautophagy, therefore, represents a new direction for investigations into the pathogenesis and possible therapy of AD.

AUTOPHAGY AND PARKINSON'S DISEASE

Parkinson's disease (PD) is characterized neuropathologically by selective neuronal degeneration, primarily in dopaminergic neurons of substantia nigra pars compacta (SNpc) and other brainstem nuclei, and the accumulation of filamentous intracytoplasmic inclusions, termed Lewy bodies (LBs). Ubiquitin and α -synuclein constitute major components of LBs, and recent studies using α -synuclein antibodies have revealed that the extent of neuropathology in PD is quite widespread.^{136,137} α -Synuclein is a protein of unclear function that localizes predominantly to presynaptic terminals in the mature CNS.¹³⁸ While the genetic factors underlying most forms of late-onset, apparently sporadic PD are poorly understood, missense or duplication/triplication mutations in α -synuclein lead to PD in rare familial cases that follow a dominant pattern of inheritance, indicating a gain-of-function effect.¹³⁸ Other genetic defects have also been detected in rare families with familial PD,¹³⁹ although the majority of cases of PD are sporadic, with no clear genetic basis.

There have been few pathological studies addressing the issue of autophagy in Parkinson's disease. Forno and Norville,¹⁴⁰ when assessing sympathetic ganglion LBs in PD, remarked on the rather vesicular nature of these inclusions, suggesting the participation of the autophagic/lysosomal pathway in the formation or dissolution of these inclusions. Anglade et al.⁹⁷ reported autophagosome-like structures in degenerating neurons in the SNpc of PD patients. In fact, the images that were shown made a more convincing case for autophagy compared to apoptosis.⁹⁷ More recently, AVs have been found to engulf mitochondria in PD substantia nigra.¹⁴¹ Given the abundant genetic, biochemical and pathological data implicating mitochondrial dysfunction in PD,¹³⁹ such dysfunction may serve as the trigger for the activation of macroautophagy, as has been shown in other systems.³¹

Neuromelanin is a pigment that accumulates with age selectively within certain populations of catecholaminergic neurons, including those of the SNpc. Sulzer et al.¹⁴² reported that neuromelanin granules are actually autophagosomes, as they are delimited by double membranes. They have hypothesized that such granules sequester dopamine and its quinone metabolites and thus prevent their potentially lethal accumulation in the cytosol. Consistent with this idea, manipulations that led to a decrease of intracellular dopamine prevented the formation of such granules in an experimental neuronal cell culture system, where granules were induced by application of L-Dopa, the precursor of dopamine.¹⁴² It appears therefore that macroautophagy is quite active even in normal SNpc, presumably serving a protective function. The possibility that this attempt at sequestration may ultimately lead to problems of its own was suggested by the finding that PC12 cells bearing such neuromelanin granules showed a less robust response to Nerve Growth Factor (NGF) application.¹⁴² The reason why neuromelanin granules arrest at the phase of autophagosomes and do not convert to mature lysosomes is unclear.

There is evidence for activation of macroautophagy in some cellular models related to PD. Overexpression of murine α -synuclein in a neuronal cell line induced autophagy, without cell death.¹⁴³ Stefanis et al.¹⁴⁴ showed that stable overexpression of mutant A53T, but not WT, α -synuclein in PC12 cells, led to the marked induction of macroautophagy and cell death without the morphological or

biochemical features of apoptosis. Cells expressing the mutant form expressed a wide range of other abnormalities, including proteasomal and lysosomal dysfunction. Stefanis et al.¹⁴⁴ hypothesized that such abnormalities in the main cellular degradation systems may be responsible for the induction of macroautophagy. Recent evidence provides further support for this hypothesis. PC12 cells expressing mutant α -synucleins are deficient in the degradation of protein substrates through the chaperone-mediated autophagy (CMA) pathway.¹⁴⁵ The upregulation of macroautophagy that occurs in PC12 cells expressing mutant α -synucleins appears therefore to be a compensatory response, following the primary defect at the level of CMA.¹⁴⁵ The primary aberrant effect appears to be an inappropriately tight binding of mutant α -synucleins to lamp2a, the receptor component of CMA. In this fashion, mutant α -synucleins do not permit the binding and uptake of the normal endogenous substrates of CMA.¹⁴⁵ Thus, in this cellular model, macroautophagy is a compensatory response of the cell attempting to cope with constraints imposed on its ability to degrade proteins, and not a primary mechanism of cell death induction. The molecular underpinnings of such compensatory responses are not yet understood. Notably, and despite macroautophagy activation, overall levels of lysosomal degradation were diminished in the lines expressing mutant α -synucleins. This is presumably due to the fact that CMA makes a major contribution to lysosomal degradation in the cells examined, especially under conditions of serum depletion.¹⁴⁵ When CMA is inhibited in the mutant α -synuclein lines, the compensatory activation of macroautophagy is presumably not enough to restore normal levels of lysosome-dependent protein degradation.

It has also been hypothesized that PD may be caused, at least in part, by dysfunction of the other major intracellular degradation system, the ubiquitin-dependent proteasomal system.¹⁴⁶ It has recently been demonstrated that neuronal cell lines treated with low doses of proteasomal inhibitors activate macroautophagy as a compensatory response.¹⁴⁷ Similar findings have been reported earlier in HeLa cells acutely treated with proteasomal inhibitors.¹⁴⁸ In the first such study in primary neurons, Rideout et al.¹⁴⁹ observed marked induction of macroautophagy and activation of the lysosomal pathway in cultured cortical neurons treated with proteasomal inhibitors. Furthermore, pharmacological modulation of macroautophagy led to alterations in the number of LB-like ubiquitinated inclusions formed in this model, suggesting that macroautophagy may be responsible for inclusion dissolution.¹⁴⁹

A related issue that is quite controversial is the degradation of α -synuclein itself. Initial studies showed that α -synuclein, in some cases in a polyubiquitinated form, accumulated in cells upon proteasomal inhibition, suggesting that the proteasome was responsible for α -synuclein degradation through the ubiquitin-dependent pathway.¹⁵⁰⁻¹⁵² A subsequent study, based on a purified system, showed that α -synuclein, as would be predicted from its natively unfolded state, does not need to be ubiquitinated to be degraded by the proteasome.¹⁵³ To complicate matters, other studies have failed to replicate the finding of accumulation of endogenous or overexpressed α -synuclein with proteasomal inhibition.¹⁵⁴⁻¹⁵⁷ Some of these studies in particular were performed with endogenous α -synuclein in PC12 cells or cultured cortical neurons,^{155,156} and thus may reflect more closely the normal turnover of the protein. While some of the earlier studies may have suffered from limitations due to the artificial transient nature of α -synuclein overexpression, the presence of epitope tags, and the expression in a non-neuronal environment, other studies have observed that proteasome inhibition impairs α -synuclein clearance

in both differentiated and undifferentiated PC12 cells.¹⁵⁸ One reason for these apparent discrepancies is that α -synuclein may be processed via a diversity of pathways and that a small change in turnover resulting from proteasome inhibition (which may not be a major degradation route) may not rapidly or overtly translate into alterations in steady-state levels. There is now considerable evidence that α -synuclein levels can accumulate with lysosomal inhibition, although the exact circumstances (cellular milieu, species of α -synuclein, class of lysosomal inhibitor) differ across studies.^{145,157-160} In the most physiological system examined so far, that of cultured post-natal rat dopaminergic neurons, endogenous α -synuclein turnover was significantly increased only by the general lysosomal inhibitor ammonium chloride, but not by the macroautophagy inhibitor 3-methyladenine. Consistent with a minor role of the proteasomes in α -synuclein degradation, application of epoxomicin, the selective proteasomal inhibitor, provided only a very small increase in the half-life of α -synuclein in this setting. In fact, α -synuclein contains a pentapeptide motif that targets it to the pathway of CMA. Analysis in an in vitro system of purified lysosomes has confirmed that α -synuclein can be degraded by CMA. Coupled with the cellular data, which indicate that α -synuclein is degraded by a lysosomal pathway that is not macroautophagy, CMA appears to be the major rate-limiting pathway used for wild-type α -synuclein degradation in a neuronal cell context.¹⁴⁵ Furthermore, the mutants A53T and A30P are deficient in CMA-dependent degradation, a fact that leads to their accumulation, but also to the aberrant effects on the CMA pathway mentioned earlier.¹⁴⁵ Interestingly, the A53T and A30P mutants show a significant dependency on macroautophagy for their clearance.^{145,158} The clearance of these proteins is delayed with inhibitors of macroautophagy and enhanced by rapamycin (which induces macroautophagy). By contrast, as mentioned, wild-type α -synuclein clearance is not obviously affected by macroautophagy inhibitors.^{145,158} Interestingly, the clearance of wild-type α -synuclein is enhanced by rapamycin in both differentiated and undifferentiated PC12 cells.¹⁵⁸ While macroautophagy under normal circumstances is not rate-limiting for wild-type α -synuclein clearance (probably because CMA and maybe the proteasome are the major pathways under basal conditions), it is possible that when macroautophagy is upregulated, it can provide an additional significant clearance route for this protein. Consistent with this, endogenous α -synuclein has been localized to autophagosomes in certain situations, for example in brains of mice that lack PS1.²⁶

Thus, the pathways used for α -synuclein degradation may vary depending on a number of factors, such as the neuronal cell type and the metabolic state. In addition to the proteasome and the lysosomes, other proteolytic systems, such as calpains¹⁶¹ or callikrein-6 (neurosin)¹⁶² may also play a role in α -synuclein cleavage and degradation. It would also be interesting to study the role of macroautophagy in α -synuclein clearance.

In conclusion, macroautophagy appears to occur under physiologic conditions in SNpc, and to be further induced in PD. It is likely to represent a compensatory response to the dysfunction of other intracellular degradation pathways, mitochondrial dysfunction or the accumulation of cytosolic dopamine. The attempt of the neurons to activate macroautophagy may eventually backfire, leading to further cellular dysfunction and autophagic death. Degradation of α -synuclein in lysosomes may be closely related to its aberrant effects. In particular, disruption of CMA may be a primary pathogenic effect of mutant α -synucleins. While mutant forms of α -synuclein are very rare, it may be important to consider if there is dysfunction of CMA in sporadic PD.

PRION DISEASES AND AUTOPHAGY

Human transmissible spongiform encephalopathies, also known as prion diseases, include scrapie in sheep, bovine spongiform encephalopathy (BSE) in cattle, Creutzfeldt-Jakob disease (CJD), Gerstmann-Sträussler-Scheinker disease (GSS), fatal familial insomnia (FFI), kuru and most recently variant CJD (vCJD) in humans. The molecular basis of prion disease is the post-translational conversion of a normal host-encoded, glycosylphosphatidylinositol (GPI)-anchored glycoprotein, the cellular prion protein (PrP^C), to an abnormal isoform, designated PrP^{Sc}. This transition confers PrP^{Sc} with a partial resistance to proteolytic degradation and detergent insolubility.¹⁶³ Consistent with the idea of an altered conformation is the finding that β -sheet content in PrP^{Sc} is higher than in PrP^C.¹⁶⁴

The neuropathology of prion diseases includes the accumulation of an abnormal protein PrP^{Sc} in the brain, neuronal vacuolation (spongiosis), neuronal loss, proliferation of astrocytes and microglia, and deposition of amyloid plaques.¹⁶⁵⁻¹⁶⁷ Studies indicate a role for autophagy in prion disease pathogenesis. In an experimental model of CJD in mice, giant AVs formed in cortical neurons.¹⁶⁰ The AV appeared at the end of the incubation period (4–6 months postinoculation), together with spongiform changes and clinical symptoms.¹⁶⁸ The formation of giant AV was also found in neurons in experimental scrapie in hamsters.¹⁶⁹ The ultrastructural features of autophagy in experimental scrapie and CJD in rodent brain are similar¹⁷⁰ and include membrane sequestration of neuronal cytoplasm, membrane proliferation, and the formation of AVs of various sizes. An important observation in the animal models has been the presence of AVs in all parts of the neuron including synaptic endings.¹⁷¹ Recent electron microscopic analysis of human brain biopsies from patients with CJD, GSS, and FFI revealed predominant changes in axon terminals that included the presence of AVs and features consistent with axonal degeneration.¹⁷² These findings suggest that synaptic dysfunction associated with autophagic dysregulation could precede other clinical and pathological features of these disorders. While there is extensive evidence for increased autophagosome numbers in prion disease brain tissue, it is unclear if the altered PrP conformation enhances autophagic activity or delays autophagosome clearance and if this is in any way related to the ensuing neuronal death.

POLYGLUTAMINE EXPANSION DISEASES AND AUTOPHAGY

Polyglutamine diseases are a group of inherited neurodegenerative disorders caused by CAG trinucleotide repeat expansion mutations that are translated into abnormally long polyglutamine tracts in the mutant proteins. Huntington's disease (HD) is the most common of these diseases, which also include spinocerebellar ataxias types (SCA) 1, 2, 3, 6, 7 and 17, spinobulbar muscular atrophy (SBMA) and dentatorubral-pallidolysian atrophy (DRPLA). Studies in humans and in mouse models suggest that polyglutamine expansion mutations are predominantly associated with a deleterious gain-of-function. One of the pathological hallmarks of this class of diseases is the formation of intracellular aggregates (also known as inclusions), which are formed by the mutant protein and also include other proteins. The roles of these inclusions are controversial (reviewed in ref. 173).

Another feature of certain polyglutamine diseases is the presence of increased numbers of autophagosomes in cell models and in vivo. In the brains of HD patients, immunoreactive huntingtin (HD gene product; htt) was found in cytoplasmic granules that were identified at the electron microscopic level to be multivesicular bodies. These

organelles are involved in protein degradation.¹⁷⁴ This observation supported the idea that the endosomal/lysosomal pathway might be involved in htt degradation. Indeed, exogenous expression of full-length or truncated wild-type or mutant htt in clonal striatal cells results in the association of htt with tubulovesicular organelles and with autophagosomes.⁹⁸ The ultrastructure of the htt-associated autophagosomes seen in vitro is diverse and includes both early autophagosomes with double-membranes and late autophagosomes. Deposition of htt occurs within the lumen and on the limiting membranes of early and late autophagosomes. Autophagosomes bound by truncated or full-length htt that are distributed throughout the cytoplasm coalesce in perinuclear regions to form htt bodies.⁹⁹ The mutant htt bodies contain fibrillar and soluble oligomeric forms of mutant htt. The polyglutamine expansion in htt significantly increases the frequency of cells with htt-associated autophagosomes. Primary striatal neurons from HD transgenic mice expressing truncated mutant htt showed a marked increase in autophagosome/lysosomal bodies following dopamine-stimulated oxidative stress.¹⁷⁵ In addition, HD patient lymphoblasts appear to form more autophagosomes compared to control lymphoblasts.¹⁷⁶ There is some evidence for enhanced autophagy occurring with expression of other polyglutamine disease proteins. Mice expressing mutant ataxin 1 (mutated in SCA1) show formation of degradative vacuoles in degenerating Purkinje cell dendrites.¹⁷⁷ HEK-293 cells exogenously expressing truncated mutant androgen receptor (with a polyglutamine expansion as in SBMA) show patches of immunoreactive mutant protein associated with large-double membraned structures and with lysosomes.¹⁷⁸

One key question is whether the increased autophagosome numbers in HD and related diseases are due to increased autophagosome synthesis, or decreased autophagosome-lysosome fusion. It is possible that similar mechanisms may exist between different polyglutamine expansion diseases, given their shared mutations.¹⁷⁹ Another important issue is if this phenomenon is a protective response, a necessary part of a death pathway, or an epiphenomenon.

In cell models of HD and related polyglutamine diseases (like SCA1 and DRPLA), increased autophagy (assessed with LC3-II blots and LC3 immunocytochemistry) appeared to be specific to cells expressing mutant constructs that also had aggregates. The increased autophagic activity in these contexts appeared to be due to the sequestration and functional inactivation of mTOR (a negative regulator of autophagy) by the polyglutamine aggregates. mTOR inactivation in cells with aggregates was also consistent with data from the brains of an HD mouse model. This may be a mechanism accounting for enhanced autophagy in diseases other than HD as mTOR sequestration was seen in the aggregates in brains of patients with a number of different polyglutamine expansion diseases.¹⁷⁹ Thus, increased autophagy in polyglutamine diseases may be due to increased autophagosome production, due to inhibition of mTOR, which negatively regulates an early stage in this pathway.

Since polyglutamine mutations act primarily by conferring toxic novel properties on the mutant protein, it is important to understand how they are cleared in cells. This is particularly relevant for post-mitotic neurons, where the proteins cannot be diluted by cell division. While the proteasome can process soluble monomeric forms of huntingtin, its narrow pore precludes proteasomal clearance of oligomeric and aggregated species. Furthermore, the proteasome cannot cleave within the polyglutamine tract.¹⁸⁰ Thus, the proteasomes effectively remove the sequences flanking the polyglutamine stretch and leave an expanded polyglutamine stretch with almost no flanking

sequences. This may enhance toxicity/misfolding—previous studies have shown that toxicity/aggregation is reduced when more sequence flanks the polyglutamine expansion (e.g., in huntingtin).¹⁸¹

Autophagy represents an alternative route for clearance of these aggregate-prone polyglutamine-expanded proteins. Chemical inhibition of the autophagy-lysosomal pathway at the autophagosome formation or autophagosome-lysosome fusion stages inhibited the clearance of mutant huntingtin fragment and was associated with increased soluble mutant protein levels, and enhanced aggregate formation and toxicity in cell models.¹⁸² The dependence of huntingtin clearance on autophagy was not only a feature of short fragments of the mutant protein but also for longer products with the expanded repeats.⁹⁹ Consistent with these data, mutant proteins with polyglutamine expansions (from Huntington's disease, spinocerebellar ataxia type 3 and dentatorubral pallidolysian atrophy) are seen in lysosomes in patient brains.¹⁸³ These data suggest that the dependence on autophagy is greatest for the most aggregate-prone species. In other words, the wild-type fragments of huntingtin show minimal accumulation when autophagy is blocked, in contrast to the mutant constructs. This phenomenon is likely to be because the aggregate-prone species are less efficiently cleared by the proteasome—autophagy is a “default” pathway. While the size of typical mammalian autophagosomes (~1.5 µm) suggests that they cannot clear large aggregates visible by light microscopy, they may be able to clear monomeric forms or small microaggregates/oligomers.

Enhancing the clearance of toxic, aggregate-prone, mutant proteins by autophagy may be a potential therapeutic avenue for diseases like HD. Rapamycin, which activates this pathway, enhanced the clearance of mutant huntingtin fragment, and reduced aggregate formation and toxicity in cell models.¹⁸² Rapamycin also protected against neurodegeneration in a *Drosophila* HD model and a rapamycin analogue, CCI-779, improved various HD-associated behavioral tasks and decreased aggregate formation in a transgenic HD mouse model.¹⁷⁹

In summary, enhanced autophagy is seen in HD and related polyglutamine diseases. This phenomenon appears to be at least in part due to the ability of aggregated huntingtin to sequester and inactivate mTOR. While aggregates may result in both deleterious and protective consequences, the sequestration of mTOR appears to represent a protective response, as enhancing autophagy with rapamycin protects against polyglutamine toxicity in cell and animal models.

CONCLUSION

For most scenarios, the key questions outlined in the introduction remain as uncertainties. To reiterate: Are increased autophagosome numbers due to increased autophagy or decreased autophagosome-lysosome clearance and is increased autophagy good, bad or an epiphenomenon? A related question is how autophagy interacts with and contributes to different cell death pathways. While data in cell and animal models suggest that autophagy induction may be beneficial in HD, this needs further development and refinement before embarking on long-term human trials. Also, it is worth considering if this strategy may be beneficial to other diseases caused by toxic, aggregate-prone proteins. It is likely that the answers to these questions vary in different contexts. It is also interesting to consider that the roles of autophagy and even possibly the machinery regulating the process may be different in response to different stimuli, or in the central nervous system compared to peripheral tissues (like the liver).

While there has been great progress in understanding the biology of autophagy, particularly in yeast, many aspects of mammalian autophagy regulation remain poorly understood, limiting the tools available to address these issues. Thus, a major challenge remains to define and characterize the machinery regulating mammalian autophagy.

Acknowledgements

The work in D.C.R.'s lab is funded by the Wellcome Trust, MRC, EUROSCA (Framework VI), and Wyeth. D.C.R. is a Wellcome Trust Senior Clinical Research Fellow. The work in Dr. Nixon's laboratory is supported by the National Institute on Aging and the Alzheimer's Association. The work in L.S.'s lab is funded by the Parkinson's Disease Foundation and the National Parkinson Foundation. M.D.F. is grateful for grant support from NIH grants NS16367 and NS035711 and a grant from the Huntington's Disease Society of America. The work in A.M.T.'s lab is funded by grants from the Wellcome Trust, MRC, BBSRC and the Huntington's Disease Society of America.

References

- Glickman MH, Ciechanover A. The ubiquitin-proteasome proteolytic pathway: destruction for the sake of construction. *Physiol Rev* 2002; 82:373-428.
- Huang WP, Klionsky DJ. Autophagy in yeast: a review of the molecular machinery. *Cell Struct Funct* 2002; 27:409-20.
- Klionsky DJ, Ohsumi Y. Vacuolar import of proteins and organelles from the cytoplasm. *Annu Rev Cell Dev Biol* 1999; 15:1-32.
- Mizushima N, Yamamoto A, Hatano M, Kobayashi Y, Kabeya Y, Suzuki K, Tokuhisa T, Ohsumi Y, Yoshimori T. Dissection of autophagosome formation using Apg5-deficient mouse embryonic stem cells. *J Cell Biol* 2001; 152:657-68.
- Kamada Y, Funakoshi T, Shintani T, Nagano K, Ohsumi M, Ohsumi Y. Tor-mediated induction of autophagy via an Apg1 protein kinase complex. *J Cell Biol* 2000; 150:1507-13.
- Cuervo AM, Dice JF. Lysosomes, a meeting point of proteins, chaperones, and proteases. *J Mol Med* 1998; 76:6-12.
- Tolkovsky AM. Autophagy, neural function and neuronal death. *Autophagy*, ed. D.J. Klionsky. 2003; Eurekah 259-66.
- Arikuni T. Functional changes of the lateral geniculate neurons during chromatolysis. *Med J Osaka Univ* 1965; 16:161-79.
- Dixon JS. 'Phagocytic' lysosomes in chromatolytic neurons. *Nature* 1967; 215:657-8.
- Holtzman E, Novikoff AB, Villaverde H. Lysosomes and GERL in normal and chromatolytic neurons of the rat ganglion nodosum. *J Cell Biol* 1967; 33:419-35.
- Kirpatrick JB. Chromatolysis in the hypoglossal nucleus of the rat: an electron microscopic analysis. *J Comp Neurol* 1968; 132:189-212.
- Matthews MR, Raisman G. A light and electron microscopic study of the cellular response to axonal injury in the superior cervical ganglion of the rat. *Proc R Soc Lond B Biol Sci* 1972; 181:43-79.
- Matthews MR. An ultrastructural study of axonal changes following constriction of post-ganglionic branches of the superior cervical ganglion in the rat. *Philos Trans R Soc Lond B Biol Sci* 1973; 264:479-505.
- Holtzman E. Lysosomes. Plenum Press, New York 1989.
- Reggioli F, Klionsky DJ. Autophagy in the eukaryotic cell. *Eukaryot Cell* 2002; 1:11-21.
- Ashford TP, Porter KR. Cytoplasmic components in hepatic cell lysosomes. *J Cell Biol* 1962; 12:198-202.
- Lemasters JJ, Nieminen AL, Qian T, Trost LC, Elmore SP, Nishimura Y, Crowe RA, Cascio WE, Bradham CA, Brenner DA. The mitochondrial permeability transition in cell death: a common mechanism in necrosis, apoptosis and autophagy. *Biochim Biophys Acta* 1998; 1366:177-96.
- Larsen KE, Sulzer D. Autophagy in neurons: a review. *Histol Histopathol* 2002; 17:897-908.
- Yu L, Lenardo MJ, Baehrecke EH. Autophagy and Caspases: A New Cell Death Program. *Cell Cycle* 2004; 3:1124-6.
- Yu L, Alva A, Su H, Dutt P, Freundt E, Welsh S, Baehrecke EH, Lenardo MJ. Regulation of an ATG7-beclin 1 program of autophagic cell death by caspase-8. *Science* 2004; 304:1500-2.
- Nissl F. Ueber die Veränderungen der Ganglionzellen am Facialiskern des Kaninchens nach ausreissung der Nerven. *Allg Z Psychiat* 1892; 48:197-8.
- Cragg BG. What is the signal for chromatolysis? *Brain Res* 1970; 23:1-21.
- Yue Z, Horton A, Bravin M, DeJager PL, Selimi F, Heintz N. A novel protein complex linking the delta 2 glutamate receptor and autophagy: implications for neurodegeneration in lurcher mice. *Neuron* 2002; 35:921-33.
- Gordon PB, Holen I, Fosse M, Rotnes JS, Seglen PO. Dependence of hepatocytic autophagy on intracellularly sequestered calcium. *J Biol Chem* 1993; 268:26107-12.

25. Kalamidas SA, Kotoulas OB, Hann AC. Studies on glycogen autophagy: effect of phorbol myristate acetate, ionophore A23187, or phentolamine. *Microsc Res Tech* 2002; 57:507-11.
26. Wilson CA, Murphy DD, Giasson BI, Zhang B, Trojanowski JQ, Lee VM. Degradative organelles containing mislocalized alpha- and beta-synuclein proliferate in presenilin-1 null neurons. *J Cell Biol* 2004; 165:335-46.
27. Xue L, Fletcher GC, Tolkskovsky AM. Autophagy is activated by apoptotic signalling in sympathetic neurons: an alternative mechanism of death execution. *Mol Cell Neurosci* 1999; 14:180-98.
28. Fletcher GC, Xue L, Passingham SK, Tolkskovsky AM. Death commitment point is advanced by axotomy in sympathetic neurons. *J Cell Biol* 2000; 150:741-54.
29. Anderson CN, Tolkskovsky AM. A role for MAPK/ERK in sympathetic neuron survival: protection against a p53-dependent, JNK-independent induction of apoptosis by cytosine arabinoside. *J Neurosci* 1999; 19:664-73.
30. Xue L, Fletcher GC, Tolkskovsky AM. Mitochondria are selectively eliminated from eukaryotic cells after blockade of caspases during apoptosis. *Curr Biol* 2001; 11:361-5.
31. Elmore SR, Qian T, Grissom SF, Lemasters JJ. The mitochondrial permeability transition initiates autophagy in rat hepatocytes. *Faseb J* 2001; 15:2286-7.
32. Blommaert EF, Luiken JJ, Blommaert PJ, van Woerkom GM, Meijer AJ. Phosphorylation of ribosomal protein S6 is inhibitory for autophagy in isolated rat hepatocytes. *J Biol Chem* 1995; 270:2320-6.
33. Noda T, Ohsumi Y, Tor, a phosphatidylinositol kinase homologue, controls autophagy in yeast. *J Biol Chem* 1998; 273:3963-6.
34. Edinger AL, Linardic CM, Chiang GG, Thompson CB, Abraham RT. Differential effects of rapamycin on mammalian target of rapamycin signaling functions in mammalian cells. *Cancer Res* 2003; 63:8451-60.
35. Caro LH, Plomp PJ, Wolvetang EJ, Kerkhof C, Meijer AJ. 3-Methyladenine, an inhibitor of autophagy, has multiple effects on metabolism. *Eur J Biochem* 1988; 175:325-9.
36. Mizushima N. Methods for monitoring autophagy. *Int J Biochem Cell Biol* 2004; 36:2491-502.
37. Kittler JT, Rostaing P, Schiavo G, Fritschy JM, Olsen R, Triller A, Moss SJ. The subcellular distribution of GABARAP and its ability to interact with NSF suggest a role for this protein in the intracellular transport of GABA(A) receptors. *Mol Cell Neurosci* 2001; 18:13-25.
38. Okazaki N, Yan J, Yuasa S, Ueno T, Kominami E, Masuho Y, Koga H, Muramatsu M. Interaction of the Unc-51-like kinase and microtubule-associated protein light chain 3 related proteins in the brain: possible role of vesicular transport in axonal elongation. *Brain Res Mol Brain Res* 2000; 85:1-12.
39. Yu LY, Jokitalo E, Sun YF, Mehlen P, D. L., Saarna M, Arumae U. GDNF-deprived sympathetic neurons die via a novel nonmitochondrial pathway. *J Cell Biol* 2003; 163:987-97.
40. Clarke PG. Developmental cell death: morphological diversity and multiple mechanisms. *Anat Embryol* 1990; 181:195-213.
41. Hornung JP, Koppel H, Clarke PG. Endocytosis and autophagy in dying neurons: an ultrastructural study in chick embryos. *J Comp Neurol* 1989; 283:425-37.
42. Borsello T, Croquelois K, Hornung JP, Clarke PG. N-methyl-D-aspartate-triggered neuronal death in organotypic hippocampal cultures is endocytic, autophagic and mediated by the c-Jun N-terminal kinase pathway. *Eur J Neurosci* 2003; 18:473-85.
43. Tarabal O, Caldero J, Llado J, Oppenheim RW, Esquerda JE. Long-lasting aberrant tubulovesicular membrane inclusions accumulate in developing motoneurons after a sublethal excitotoxic insult: a possible model for neuronal pathology in neurodegenerative disease. *J Neurosci* 2001; 21:8072-81.
44. Dickens M, Rogers JS, Cavanagh J, Raitano A, Xia Z, Halpern JR, Greenberg ME, Sawyers CL, Davis RJ. A cytoplasmic inhibitor of the JNK signal transduction pathway. *Science* 1997; 277:693-6.
45. Harding TC, Xue L, Bienemann A, Haywood D, Dickens M, Tolkskovsky AM, Uney JB. Inhibition of JNK by overexpression of the JNK binding domain of JIP-1 prevents apoptosis in sympathetic neurons. *J Biol Chem* 2001; 276:4531-4.
46. Borsello T, Bressoud R, Mottier V, Gonzalez N, Gomez G, Clarke PG. Kainate-induced endocytosis in retinal amacrine cells. *J Comp Neurol* 2003; 465:286-95.
47. Yu L, Alva A, Su H, Dutt P, Freundt E, Welsh S, Bachrecke EH, Lenardo MJ. Regulation of an ATG7-beclin 1 program of autophagic cell death by caspase-8. *Science* 2004; 304:1500-2.
48. Shimizu S, Kanaseki T, Mizushima N, Mizuta T, Arakawa-Kobayashi S, Thompson CB, Tsujimoto Y. Role of Bcl-2 family proteins in a non-apoptotic programmed cell death dependent on autophagy genes. *Nat Cell Biol* 2004; 6:1221-8.
49. Boya P, Andreau K, Poncet D, Zamzami N, Perfettini JL, Metivier D, Ojcius DM, Jaattela M, Kroemer G. Lysosomal membrane permeabilization induces cell death in a mitochondrion-dependent fashion. *J Exp Med* 2003; 197:1323-34.
50. Foghsgaard L, Wissing D, Mauch D, Lademann U, Bastholm L, Boes M, Elling F, Leist M, Jaattela M. Cathepsin B acts as a dominant execution protease in tumor cell apoptosis induced by tumor necrosis factor. *J Cell Biol* 2001; 153:999-1010.
51. Akassoglou K, Malester B, Xu J, Tessarollo L, Rosenbluth J, Chao MV. Brain-specific deletion of neurotrophin target esterase/swisscheese results in neurodegeneration. *Proc Natl Acad Sci USA* 2004; 101:5075-80.
52. Larsen KE, Fon EA, Hastings TG, Edwards RH, Sulzer D. Methamphetamine-induced degeneration of dopaminergic neurons involves autophagy and upregulation of dopamine synthesis. *J Neurosci* 2002; 22:8951-60.
53. Hall DH, Gu G, Garcia-Anoveros J, Gong L, Chalfie M, Driscoll M. Neuropathology of degenerative cell death in *Caenorhabditis elegans*. *J Neurosci* 1997; 17:1033-45.
54. Syntichaki P, Xu K, Driscoll M, Tavernarakis N. Specific aspartyl and calpain proteases are required for neurodegeneration in *C. elegans*. *Nature* 2002; 419:939-44.
55. Melendez A, Tallozy Z, Seaman M, Eskelinen EL, Hall DH, Levine B. Autophagy genes are essential for dauer development and life-span extension in *C. elegans*. *Science* 2003; 301:1387-91.
56. Gutierrez MG, Munafò DB, Beron W, Colombo MI. Rab7 is required for the normal progression of the autophagic pathway in mammalian cells. *J Cell Sci* 2004; 117:2687-97.
57. Jager S, Bucci C, Tanida I, Ueno T, Kominami E, Saftig P, Eskelinen EL. Role for Rab7 in maturation of late autophagic vacuoles. *J Cell Sci* 2004; 117:4837-48.
58. Blum D, Hemming FJ, Galas MC, Torch S, Cuvelier L, Schiffmann SN, Sadoul R. Increased Alix (apoptosis-linked gene-2 interacting protein X) immunoreactivity in the degenerating striatum of rats chronically treated by 3-nitropropionic acid. *Neurosci Lett* 2004; 368:309-13.
59. Hemming FJ, Fraboulet S, Blot B, Sadoul R. Early increase of apoptosis-linked gene-2 interacting protein X in areas of kainate-induced neurodegeneration. *Neuroscience* 2004; 123:887-95.
60. Trioulier Y, Torch S, Blot B, Cristina N, Chatellard-Causse C, Verna JM, Sadoul R. Alix, a protein regulating endosomal trafficking, is involved in neuronal death. *J Biol Chem* 2004; 279:2046-52.
61. Matsuo H, Chevallier J, Mayran N, Le Blanc I, Ferguson C, Faure J, Blanc NS, Matile S, Dubochet J, Sadoul R. Role of LBPA and Alix in multivesicular liposome formation and endosome organization. *Science* 2004; 303:531-4.
62. Su JH, Kessler JB, Head E, Cotman CW. Caspase-cleaved amyloid precursor protein and activated caspase-3 are co-localized in the granules of granulovacuolar degeneration in Alzheimer's disease and Down's syndrome brain. *Acta Neuropathol (Berl)* 2002; 104:1-6.
63. Zuo J, De Jager PL, Takahashi KA, Jiang W, Linden DJ, Heintz N. Neurodegeneration in Lurcher mice caused by mutation in delta2 glutamate receptor gene. *Nature* 1997; 388:769-73.
64. Norman DJ, Feng L, Cheng SS, Gubbay J, Chan E, Heintz N. The lurcher gene induces apoptotic death in cerebellar Purkinje cells. *Development* 1995; 121:1183-93.
65. Wullner U, Loschmann PA, Weller M, Klockgether T. Apoptotic cell death in the cerebellum of mutant weaver and lurcher mice. *Neurosci Lett* 1995; 200:109-12.
66. Wullner U, Weller M, Schulz JB, Krajewski S, Reed JC, Klockgether T. Bcl-2, Bax and Bcl-x expression in neuronal apoptosis: a study of mutant weaver and lurcher mice. *Acta Neuropathol (Berl)* 1998; 96:233-8.
67. Doughty ML, De Jager PL, Korsmeyer SJ, Heintz N. Neurodegeneration in Lurcher mice occurs via multiple cell death pathways. *J Neurosci* 2000; 20:3687-94.
68. Selimi F, Vogel MW, Mariani J. Bax inactivation in lurcher mutants rescues cerebellar granule cells but not purkinje cells or inferior olivary neurons. *J Neurosci* 2000; 20:5339-45.
69. Liang XH, Jackson S, Seaman M, Brown K, Kempkes B, Hibshoosh H, Levine B. Induction of autophagy and inhibition of tumorigenesis by beclin 1. *Nature* 1999; 402:672-6.
70. Selimi F, Lohof AM, Heitz S, Lalouette A, Jarvis CI, Bailly Y, Mariani J. Lurcher GRID2-induced death and depolarization can be dissociated in cerebellar Purkinje cells. *Neuron* 2003; 37:813-9.
71. Yuan J, Lipinski M, Degterev A. Diversity in the mechanisms of neuronal cell death. *Neuron* 2003; 40:401-13.
72. Kabeya Y, Mizushima N, Yamamoto A, Oshitani-Okamoto S, Ohsumi Y, Yoshimori T. LC3, GABARAP and GATE16 localize to autophagosomal membrane depending on form-II formation. *J Cell Sci* 2004; 117:2805-12.
73. Wang H, Bedford FK, Brandon NJ, Moss SJ, Olsen RW. GABA(A)-receptor-associated protein links GABA(A) receptors and the cytoskeleton. *Nature* 1999; 397:69-72.
74. Kashiwabuchi N, Ikeda K, Araki K, Hirano T, Shibuki K, Takayama C, Inoue Y, Kutsuwada T, Yagi T, Kang Y, et al. Impairment of motor coordination, Purkinje cell synapse formation, and cerebellar long-term depression in GluR delta 2 mutant mice. *Cell* 1995; 81:245-52.
75. Yankner BA. New clues to Alzheimer's disease: unraveling the roles of amyloid and tau. *Nat Med* 1996; 2:850-2.
76. Burke WJ, Chung HD, Huang JS, Huang SS, Haring JH, Strong R, Marshall GL, Joh TH. Evidence for retrograde degeneration of epinephrine neurons in Alzheimer's disease. *Ann Neurol* 1988; 24:532-6.
77. Coleman PD, Yao PJ. Synaptic slaughter in Alzheimer's disease. *Neurobiol Aging* 2003; 24:1023-7.
78. Morsch R, Simon W, Coleman PD. Neurons may live for decades with neurofibrillary tangles. *J Neuropathol Exp Neurol* 1999; 58:188-97.
79. Pearson RC, Powell TP. Anterograde vs. retrograde degeneration of the nucleus basalis medialis in Alzheimer's disease. *J Neural Transm Suppl* 1987; 24:139-46.
80. Scheff SW, Price DA. Synaptic pathology in Alzheimer's disease: a review of ultrastructural studies. *Neurobiol Aging* 2003; 24:1029-46.
81. Hashimoto M, Masliah E. Cycles of aberrant synaptic sprouting and neurodegeneration in Alzheimer's and dementia with Lewy bodies. *Neurochem Res* 2003; 28:1743-56.
82. Larner AJ. The cortical neuritic dystrophy of Alzheimer's disease: nature, significance, and possible pathogenesis. *Dementia* 1995; 6:218-24.
83. McKee AC, Kowall NW, Kosik KS. Microtubular reorganization and dendritic growth response in Alzheimer's disease. *Ann Neurol* 1989; 26:652-9.
84. Nixon RA. A "protease activation cascade" in the pathogenesis of Alzheimer's disease. *Ann N Y Acad Sci* 2000; 924:117-31.

85. Cataldo AM, Barnett JL, Pieroni, C, Nixon RA. Increased neuronal endocytosis and protease delivery to early endosomes in sporadic Alzheimer's disease: neuropathologic evidence for a mechanism of increased beta-amyloidogenesis. *J Neurosci* 1997; 17:6142-51.
86. Cataldo AM, Hamilton DJ, Barnett JL, Paskevich PA, Nixon RA. Properties of the endosomal-lysosomal system in the human central nervous system: disturbances mark most neurons in populations at risk to degenerate in Alzheimer's disease. *J Neurosci* 1996; 16:186-99.
87. Mathews PM, Guerra CB, Jiang Y, Grbovic OM, Kao BH, Schmidt SD, Dinakar R, Mercken M, Hille-Rehfeld A, Rohrer J, Mehta P, Cataldo AM, Nixon RA. Alzheimer's disease-related overexpression of the cation-dependent mannose 6-phosphate receptor increases Abeta secretion: role for altered lysosomal hydrolase distribution in beta-amyloidogenesis. *J Biol Chem* 2002; 277:5299-307.
88. Mathews PM, Jiang Y, Schmidt SD, Grbovic OM, Mercken M, Nixon RA. Calpain activity regulates the cell surface distribution of amyloid precursor protein. Inhibition of calpains enhances endosomal generation of beta-cleaved C-terminal APP fragments. *J Biol Chem* 2002; 277:36415-24.
89. Ginsberg SD, Hemby SE, Lee VM, Eberwine JH, and Trojanowski JQ. Expression profile of transcripts in Alzheimer's disease tangle-bearing CA1 neurons. *Ann Neurol* 2000; 48:77-87.
90. Cataldo A, Rebeck GW, Ghetti B, Hulette C, Lippa C, Van Broeckhoven C, van Duijn C, Cras P, Bogdanovic N, Bird T, Peterhoff C, Nixon R. Endocytic disturbances distinguish among subtypes of Alzheimer's disease and related disorders. *Ann Neurol* 2001; 50:661-5.
91. Nixon RA, Weigel J, Kumar A, Yu WH, Peterhoff CM, Cataldo AM, Cuervo AM. Autophagy in the neurodegeneration of Alzheimer's disease. *J Neuropath Exp Neurol* 2004; Submitted.
92. Nixon RA, Yu WH, Cuervo AM, Kumar A, Yang DS, Peterhoff CM. Autophagy in Alzheimer's Disease: Failure of a neuroprotective mechanism in Alzheimer's Disease and Related Disorders: Research Advances. Vol. 10. I. Khalid and B. Winblad, editors. "Ana Aslan" International Academy of Aging, Bucharest. 2004; In press.
93. Yu WH, Cuervo AM, Kumar A, Peterhoff CM, Schmidt SD, Mohan PS, Mercken M, Farmer MR, Tjernberg L, Jiang Y, Weigel J, Lamb BT, Duff KD, Uchiyama Y, Naslund J, Mathews PM, Cataldo AM, Nixon RA. Macroautophagy—A Novel Amyloid- β (Ab) Peptide-generating Pathway Activated in Alzheimer's Disease. 2004; Submitted.
94. Cataldo AM, Paskevich PA, Kominami E, Nixon RA. Lysosomal hydrolases of different classes are abnormally distributed in brains of patients with Alzheimer disease. *Proc Natl Acad Sci U S A* 1991; 88:10998-1002.
95. Kawai M, Cras P, Richey P, Tabaton M, Lowery DE, Gonzalez-DeWhitt PA, Greenberg BD, Gambetti P, Perry G. Subcellular localization of amyloid precursor protein in senile plaques of Alzheimer's disease. *Am J Pathol* 1992; 140:947-58.
96. Suzuki K, Terry RD. Fine structural localization of acid phosphatase in senile plaques in Alzheimer's presenile dementia. *Acta Neuropathol (Berl)* 1967; 8:276-84.
97. Anglade P, Vyas S, Javoy-Agid E, Herrero MT, Michel PP, Marquez J, Mouatt-Prigent A, Ruberg M, Hirsch EC, Agid Y. Apoptosis and autophagy in nigral neurons of patients with Parkinson's disease. *Histol Histopathol* 1997; 12:25-31.
98. Kegel KB, Kim M, Sapp E, McIntyre C, Castano JG, Aronin N, DiFiglia M. Huntingtin expression stimulates endosomal-lysosomal activity, endosome tubulation, and autophagy. *J Neurosci* 2000; 20:7268-78.
99. Qin ZH, Wang Y, Kegel KB, Kazantsev A, Apostol BL, Thompson LM, Yoder J, Aronin N, DiFiglia M. Autophagy regulates the processing of amino terminal huntingtin fragments. *Hum Mol Genet* 2003; 12:3231-44.
100. Nixon RA, Cataldo AM. The lysosomal system in neuronal cell death: a review. *Ann N Y Acad Sci* 1993; 679:87-109.
101. Masliah E, Mallory M, DeTeresa R, Lamont S, Miller A, Terry RD, Carragher B, Ellisman M. Re-evaluation of the structural organization of the neuritic plaques in Alzheimer's disease. *J Neuropathology and Experimental Neurology* 1993; 52:619-32.
102. Schmidt ML, DiDario AG, Lee VM, Trojanowski JQ. An extensive network of PHF tau-rich dystrophic neurites permeates neocortex and nearly all neuritic and diffuse amyloid plaques in Alzheimer disease. *FEBS Lett* 1994; 344:69-73.
103. Benzing WC, Mufson EJ, Armstrong DM. Alzheimer's disease-like dystrophic neurites characteristically associated with senile plaques are not found within other neurodegenerative diseases unless amyloid beta-protein deposition is present. *Brain Res* 1993; 606:10-8.
104. Hollenbeck PJ. Products of endocytosis and autophagy are retrieved from axons by regulated retrograde organelle transport. *J Cell Biol* 1993; 121:305-15.
105. Overly CC, Hollenbeck PJ. Dynamic organization of endocytic pathways in axons of cultured sympathetic neurons. *J Neurosci* 1996; 16:6056-64.
106. Overly CC, Lee KD, Berthiaume E, Hollenbeck PJ. Quantitative measurement of intraorganelle pH in the endosomal-lysosomal pathway in neurons by using ratiometric imaging with pyranine. *Proc Natl Acad Sci U S A* 1995; 92:3156-60.
107. Faddis BT, Hasbani MJ, Goldberg MP. Calpain activation contributes to dendritic remodeling after brief excitotoxic injury in vitro. *J Neurosci* 1997; 17:951-9.
108. Obin M, Mescio E, Gong X, Haas AL, Joseph J, Taylor A. Neurite outgrowth in PC12 cells. Distinguishing the roles of ubiquitylation and ubiquitin-dependent proteolysis. *J Biol Chem* 1999; 274:11789-95.
109. Shea TB, Beermann ML, Nixon RA. Multiple proteases regulate neurite outgrowth in NB2a/dl neuroblastoma cells. *J Neurochem* 1991; 56:842-51.
110. Arendt T. Alzheimer's disease as a loss of differentiation control in a subset of neurons that retain immature features in the adult brain. *Neurobiol Aging* 2000; 21:783-96.
111. Yang Y, Mufson EJ, Herrup K. Neuronal cell death is preceded by cell cycle events at all stages of Alzheimer's disease. *J Neurosci* 2003; 23:2557-63.
112. Brunk UT, Terman A. The mitochondrial-lysosomal axis theory of aging: accumulation of damaged mitochondria as a result of imperfect autophagocytosis. *Eur J Biochem* 2002; 269:1996-2002.
113. Brunk UT, Dalen H, Roberg K, Hellquist HB. Photo-oxidative disruption of lysosomal membranes causes apoptosis of cultured human fibroblasts. *Free Radic Biol Med* 1997; 23:616-26.
114. Tolkovsky AM, Xue L, Fletcher GC, Borutaite V. Mitochondrial disappearance from cells: a clue to the role of autophagy in programmed cell death and disease? *Biochimie* 2002; 84:233-40.
115. Uchiyama Y. Autophagic cell death and its execution by lysosomal cathepsins. *Arch Histol Cytol* 2001; 64:233-46.
116. Levy-Lahad E, Wasco W, Poorkaj P, Romano DM, Oshima J, Pettingell WH, Yu CE, Jondro PD, Schmidt SD, Wang K, et al. 1995; Candidate gene for the chromosome 1 familial Alzheimer's disease locus [see comments]. *Science* 269:973-7.
117. Price DL, Sisodia SS. Mutant genes in familial Alzheimer's disease and transgenic models. *Annu Rev Neurosci* 1998; 21:479-505.
118. Sherrington R, Rogaev EI, Liang Y, Rogaeva EA, Levesque G, Ikeda M, Chi H, Lin C, Li G, Holman K, et al. Cloning of a gene bearing missense mutations in early-onset familial Alzheimer's disease. *Nature* 1995; 375:754-60.
119. Cataldo AM, Peterhoff CM, Schmidt SD, Terio NB, Duff K, Beard M, Mathews PM, Nixon RA. Presenilin mutations in familial Alzheimer's disease and transgenic mouse models accelerate neuronal lysosomal pathology. *J Neuropath Exp Neurol* 2004; In press.
120. Esselens C, Oorschot V, Baert V, Raemaekers T, Spittaels K, Serneels L, Zheng H, Saftig P, De Strooper B, Klumperman J, Annaert W. Presenilin 1 mediates the turnover of telencephalin in hippocampal neurons via an autophagic degradative pathway. *J Cell Biol* 2004; 166:1041-54.
121. Yu WH, Kumar A, Peterhoff C, Kulnane LS, Uchiyama Y, Lamb BT, Cuervo AM, Nixon RA. Autophagic vacuoles are enriched in amyloid precursor protein-secretase activities: implications for beta-amyloid peptide over-production and localization in Alzheimer's disease. *Int J Biochem Cell Biol* 2004; 36:2531-40.
122. Keller JN, Hanni KB, Markesbery WR. Impaired proteasome function in Alzheimer's disease. *J Neurochem* 2000; 75:436-9.
123. Keller JN, Huang FF, Markesbery WR. Decreased levels of proteasome activity and proteasome expression in aging spinal cord. *Neuroscience* 2000; 98:149-56.
124. Baehrecke EH. Autophagic programmed cell death in *Drosophila*. *Cell Death Differ* 2003; 10:940-5.
125. Boland B, Campbell V. Abeta-mediated activation of the apoptotic cascade in cultured cortical neurones: a role for cathepsin-L. *Neurobiol Aging* 2004; 25:83-91.
126. Florez-McClure ML, Linseman DA, Chu CT, Barker PA, Bouchard RJ, Le SS, Laessig TA, Heidenreich KA. The p75 neurotrophin receptor can induce autophagy and death of cerebellar Purkinje neurons. *J Neurosci* 2004; 24:4498-509.
127. Jellinger KA, Stadelmann C. Problems of cell death in neurodegeneration and Alzheimer's Disease. *J Alzheimers Dis* 2001; 3:31-40.
128. Pasternak SH, Bagshaw RD, Guiral M, Zhang S, Ackerley CA, Pak BJ, Callahan JW, Mahuran DJ. Presenilin-1, nicastrin, amyloid precursor protein, and gamma-secretase activity are co-localized in the lysosomal membrane. *J Biol Chem* 2003; 278:26687-94.
129. Askanas V, Engel WK. Does overexpression of betaAPP in aging muscle have a pathogenic role and a relevance to Alzheimer's disease? Clues from inclusion body myositis, cultured human muscle, and transgenic mice. *Am J Pathol* 1998; 153:1673-7.
130. Askanas V, Engel WK. Sporadic inclusion-body myositis and its similarities to Alzheimer disease brain. Recent approaches to diagnosis and pathogenesis, and relation to aging. *Scand J Rheumatol* 1998; 27:389-405.
131. Cupers P, Orlans I, Craessaerts K, Annaert W, De Strooper B. The amyloid precursor protein (APP)-cytoplasmic fragment generated by gamma-secretase is rapidly degraded but distributes partially in a nuclear fraction of neurones in culture. *J Neurochem* 2001; 78:1168-78.
132. Dunn WA Jr. Studies on the mechanisms of autophagy: maturation of the autophagic vacuole. *J Cell Biol* 1990; 110:1935-45.
133. Vassar R, Citron M. Abeta-generating enzymes: recent advances in beta- and gamma-secretase research. *Neuron* 2000; 27:419-22.
134. Smith DH, Uryu K, Saatman KE, Trojanowski JQ, McIntosh TK. Protein accumulation in traumatic brain injury. *Neuromolecular Med* 2003; 4:59-72.
135. Bursch W. The autophagosomal-lysosomal compartment in programmed cell death. *Cell Death Differ* 2001; 8:569-81.
136. Duda JE, Giasson BI, Mabon ME, Lee VM, Trojanowski JQ. Novel antibodies to synuclein show abundant striatal pathology in Lewy body diseases. *Ann Neurol* 2002; 52:205-10.
137. Braak H, Del Tredici K, Rub U, de Vos RA, Jansen Steur EN, Braak E. Staging of brain pathology related to sporadic Parkinson's disease. *Neurobiol Aging* 2003; 24:197-211.
138. Vekrellis K, Rideout HJ, Stefanis L. Neurobiology of alpha-Synuclein. *Mol Neurobiol* 2004; 30:1-22.
139. Dawson TM, Dawson VL. Molecular pathways of neurodegeneration in Parkinson's disease. *Science* 2003; 302:819-22.
140. Forno LS, Norville RL. Ultrastructure of Lewy bodies in the stellate ganglion. *Acta Neuropathol (Berl)* 1976; 34:183-97.
141. Zhu JH, Guo F, Shelburne J, Watkins S, Chu CT. Localization of phosphorylated ERK/MAP kinases to mitochondria and autophagosomes in Lewy body diseases. *Brain Pathol* 2003; 13:473-81.

142. Sulzer D, Bogulavsky J, Larsen KE, Behr G, Karatekin E, Kleinman MH, Turro N, Krantz D, Edwards RH, Greene LA, Zecca L. Neuromelanin biosynthesis is driven by excess cytosolic catecholamines not accumulated by synaptic vesicles. *Proc Natl Acad Sci U S A* 2000; 97:11869-74.
143. Hsu LJ, Sagara Y, Arroyo A, Rockenstein E, Sisk A, Mallory M, Wong J, Takenouchi T, Hashimoto M, Masliah E. γ -synuclein promotes mitochondrial deficit and oxidative stress. *Am J Pathol* 2000; 157:401-10.
144. Stefanis L, Larsen KE, Rideout HJ, Sulzer D, Greene LA. Expression of A53T but not wild-type α -synuclein in PC12 cells induces alterations of the ubiquitin-dependent degradation system, loss of dopamine release, and autophagic cell death. *J. Neurosci* 2001; 21:9549-60.
145. Cuervo AM, Stefanis L, Fredenburg R, Lansbury PT, Sulzer D. Impaired degradation of mutant α -synuclein by chaperone-mediated autophagy. *Science* 2004; 368:1292-5.
146. McNaught KS, Olanow CW, Halliwell B, Isacson O, Jenner P. Failure of the ubiquitin-proteasome system in Parkinson's disease. *Nature Rev Neurosci* 2001; 2:589-94.
147. Ding Q, Dimayuga E, Martin S, Bruce-Keller AJ, Nukala V, Cuervo AM, Keller JN. Characterization of chronic low-level proteasome inhibition on neural homeostasis. *J Neurochem* 2003; 86:489-97.
148. Wojcik C, Schroeter D, Wilk S, Lamprecht J, Paweletz N. Ubiquitin-mediated proteolysis centers in HeLa cells: indication from studies of an inhibitor of the chymotrypsin-like activity of the proteasome. *Eur J Cell Biol* 1996; 71:311-8.
149. Rideout HJ, Lang-Rollin I, Stefanis L. Involvement of macroautophagy in the dissolution of neuronal inclusions. *Int J Biochem Cell Biol* 2004; 36:2551-62.
150. Bennett MC, Bishop JF, Leng Y, Chock PB, Chase TN, Mouradian MM. Degradation of α -synuclein by proteasome. *J Biol Chem* 1999; 274:33855-8.
151. Imai Y, Soda M, Takahashi R. Parkin suppresses unfolded protein stress-induced cell death through its E3 ubiquitin-protein ligase activity. *J Biol Chem* 2000; 275:35661-4.
152. Liu Y, Fallon L, Lashuel HA, Liu Z, Lansbury PT Jr. The UCH-L1 gene encodes two opposing enzymatic activities that affect α -synuclein degradation and Parkinson's disease susceptibility. *Cell* 2002; 111:209-18.
153. Tofaris GK, Layfield R, Spillantini MG. α -Synuclein metabolism and aggregation is linked to ubiquitin-independent degradation by the proteasome. *FEBS Lett* 2001; 509:22-6.
154. Ancolio K, da Costa CA, Ueda K, Checler F. α -Synuclein and the Parkinson's disease-related mutant Ala53Thr- α -synuclein do not undergo proteasomal degradation in HEK293 and neuronal cells. *Neurosci Lett* 2000; 285:79-82.
155. Rideout HJ, Larsen KE, Sulzer D, Stefanis L. Proteasomal inhibition leads to formation of ubiquitin/ α -synuclein-immunoreactive inclusions in PC12 cells. *J Neurochem* 2001; 78:899-908.
156. Rideout HJ, Stefanis L. Proteasomal inhibition-induced inclusion formation and death in cortical neurons require transcription and ubiquitination. *Mol Cell Neurosci* 2002; 21:223-38.
157. Paxinou E, Chen Q, Weisse M, Giasson BI, Norris EH, Rueter SM, Trojanowski JQ, Lee VM, Ischiropoulos H. Induction of α -synuclein aggregation by intracellular nitrate insult. *J Neurosci* 2001; 21:8053-61.
158. Webb JL, Ravikumar B, Atkins J, Skepper JN, Rubinshtein DC. α -Synuclein is degraded by both autophagy and the proteasome. *J Biol Chem* 2003; 278:25009-13.
159. Lee HJ, Khoshaghideh F, Patel S, Lee SJ. Clearance of α -synuclein oligomeric intermediates via the lysosomal degradation pathway. *J Neurosci* 2004; 24:1888-96.
160. Li W, Lesuisse C, Xu Y, Troncoso JC, Price DL, Lee MK. Stabilization of α -synuclein protein with aging and familial parkinson's disease-linked A53T mutation. *J Neurosci* 2004; 24:9400-9.
161. Mishizen-Eberz AJ, Guttman RP, Giasson BI, Day GA, Hodara R, Ischiropoulos H, Lee VM, Trojanowski JQ, Lynch DR. Distinct cleavage patterns of normal and pathologic forms of α -synuclein by calpain I in vitro. *J Neurochem* 2003; 86:836-47.
162. Iwata A, Maruyama M, Akagi T, Hashikawa T, Kanazawa I, Tsuji S, Nukina N. α -Synuclein degradation by serine protease neurosin: Implication for pathogenesis of synucleinopathies. *Hum Mol Genet* 2003; 12:2625-35.
163. Wadsworth JDF, Hill AF, Beck JA, Collinge J. Molecular and clinical classification of human prion disease. *British Med Bulletin* 2003; 66:241-54.
164. Pan K-M, Baldwin M, Nguyen J, Gasset M, Serban A, Groth D, Mehlhorn I, Huang Z, Fletterick RJ, Cohen FE. Conversion of α -helices to β -sheets features in the formation of the scrapie prion proteins. *Proc Natl Acad Sci USA* 1999; 96:10962-6.
165. Brandner S. CNS Pathogenesis of prion diseases. *British Med Bulletin* 2003; 66:131-9.
166. Budka H. Neuropathology of prion diseases. *British Med Bulletin* 2003; 66:121-30.
167. Aguzzi AA, Polymenidou M. Mammalian prion biology: One century of evolving concepts. *Cell* 2004; 116:313-27.
168. Boellaard JW, Schlote W, Tateishi J. Neuronal autophagy in experimental Creutzfeldt-Jakob's disease. *Acta Neuropathol (Berl)* 1989; 78:410-8.
169. Boellaard JW, Kao M, Schlote W, Diringer H. Neuronal autophagy in experimental scrapie. *Acta Neuropathol (Berl)* 1991; 82:225-8.
170. Liberski PP, Gajdusek DC, Brown P. How do neurons degenerate in prion diseases or transmissible spongiform encephalopathies (TSEs): neuronal autophagy revisited. *Acta Neurobiol Exp* 2002; 62:141-7.
171. Jeffrey M, Scott JR, Williams A, Fraser H. Ultrastructural features of spongiform encephalopathy transmitted to mice from three species of bovidae. *Acta Neuropathol (Berl)* 1992; 84:559-69.
172. Sikorska B, Liberski PP, Giraud P, Kopp N, Brown P. Autophagy is a part of ultrastructural synaptic pathology in Creutzfeldt-Jakob disease: a brain biopsy study. *Int J Biochem Cell Biol* 2004; 36:2563-73.
173. Rubinshtein DC. Lessons from animal models of Huntington's disease. *Trends Genet* 2002; 18:202-9.
174. Sapp E, Schwarz C, Chase K, Bhide PG, Young AB, Penney J, Vonsattel JP, Aronin N, DiFiglia M. Huntingtin localization in brains of normal and Huntington's disease patients. *Ann Neurol* 1997; 42:604-12.
175. Petersen A, Larsen KE, Behr GG, Romero N, Przedborski S, Brundin P, Sulzer D. Expanded CAG repeats in exon 1 of the Huntington's disease gene stimulate dopamine-mediated striatal neuron autophagy and degeneration. *Hum Mol Genet* 2001; 10:1243-54.
176. Nagata E, Sawa A, Ross CA, Snyder SH. Autophagosome-like vacuole formation in Huntington's disease lymphoblasts. *Neuroreport* 2004; 15:1325-8.
177. Skinner PJ, Viera-Green CA, Clark HB, Zoghbi HY, Orr HT. Altered trafficking of membrane proteins in purkinje cells of SCA1 transgenic mice. *Am J Pathol* 2001; 159:905-13.
178. Taylor JP, Tanaka F, Robitschek J, Sandoval CM, Taye A, Markovic-Plese S, Fischbeck KH. Aggresomes protect cells by enhancing the degradation of toxic polyglutamine-containing protein. *Hum Mol Genet* 2003; 12:749-57.
179. Ravikumar B, Vacher C, Berger Z, Davies JE, Luo S, Oroz LG, Scaravilli F, Easton DF, Duden R, O'Kane CJ, Rubinshtein DC. Inhibition of mTOR induces autophagy and reduces toxicity of polyglutamine expansions in fly and mouse models of Huntington disease. *Nat Genet* 2004; 36:585-95.
180. Venkatraman P, Wetzel R, Tanaka M, Nukina N, Goldberg AL. Eukaryotic proteasomes cannot digest polyglutamine sequences and release them during degradation of polyglutamine-containing proteins. *Mol Cell* 2004; 14:95-104.
181. Hackam AS, Singaraja R, Wellington CL, Metzler M, McCutcheon K, Zhang T, Kalchman M, Hayden MR. The influence of huntingtin protein size on nuclear localization and cellular toxicity. *J Cell Biol* 1998; 141:1097-105.
182. Ravikumar B, Duden R and Rubinshtein DC. Aggregate-prone proteins with polyglutamine and polyalanine expansions are degraded by autophagy. *Hum Mol Genet* 2002; 11:1107-17.
183. Yamada M, Tsuji S, Takahashi H. Involvement of lysosomes in the pathogenesis of CAG repeat diseases. *Ann Neurol* 2002; 52:498-503.