

Computing Local Thickness of 3D Structures with ImageJ

Robert P. Dougherty* and Karl-Heinz Kunzelmann**

*OptiNav, Inc., 10914 NE 18 ST, Bellevue, WA, 98004

**Poliklinik für Zahnerhaltung und Parodontologie Ludwig-Maximilians-Universität-München, Goethestr. 70 D-80336, München, Germany

The local thickness described by Hildebrand and Ruesgsegger can characterize a 3D binary image of a complex structure such as a bone, a cell, or a paper fiber [1]. Such images are available from, for example, micro-computed tomography [2]. Let $\Omega \subset R^3$ be the structure. The local thickness at point $p \in \Omega$ is the diameter of the largest sphere that contains p and is completely inside the structure.

The problem is to transform the 3D binary image into the 3D local thickness image. Several steps to are given in [1]. The first is a Euclidean distance transformation (EDT) The distance map at $q \in \Omega$, $D_{\text{map}}(q)$, is the distance from q to the nearest background point. The sphere of radius $D_{\text{map}}(q)$ centered at q is completely inside the structure. If p is within this sphere, then the local thickness at p , $t(p)$, is at least $D_{\text{map}}(q)$. Points q are scanned to find $t(p)$ by maximizing this bound. A further simplification is to first remove many of the points whose spheres of radius $D_{\text{map}}(q)$ are completely contained within the corresponding spheres of other points. Removing all of the redundant points would produce the distance ridge (DR) which is related to the skeleton and the medial axis [1, 4].

A 3D local thickness procedure has been implemented in Java as a plugin for ImageJ [5]. The EDT uses the fast algorithm given by Saito and Toriwaki [6]. The DR computation is a simplified form of a template scheme [7]. Care was taken to optimize the use computational resources. No temporary 3D buffers are required. A related implementation in C was given previously by Coeurjolly [8]

An example using a microtomographic dataset of a root canal is shown in Figures 1-2. The complete computation ($342 \times 254 \times 669$ pixels) required 38.8 seconds PC.

- [1] T. Hildebrand and P. Ruesgsegger, "A new method for the model-independent assessment of thickness in three-dimensional images" *J. of Microscopy*, **185** (1996) 67-75.
- [2] O.A. Peters, A. Laib, P. Ruegsegger and F. Barbakow, "Three-dimensional analysis of root canal geometry by highresolution computed tomography". *J Dent Res* **79** (2000) 1405-1409.
- [3] P. E. Danielsson, "Euclidean Distance Mapping," *Comp. Gr. and Img. Proc.*, **14** (1980) 227-248.
- [4] T. Saito and J. Toriwaki, "Reverse distance transformation and skeletons based on the Euclidean metric for n-dimensional digital binary pictures" *IEICE Trans. Inf. & Syst.* **E77-D** (1994) 1005-16.
- [5] W. S. Rasband, ImageJ, National Institutes of Health, USA, <http://rsb.info.nih.gov/ij/>, 2007.
- [6] T. Saito and J. Toriwaki, "New algorithms for Euclidean distance transformation on an n-dimensional digitized picture with applications," *Pattern Recognition* **27** (1994) 1551-1565.
- [7] E. Remy and E. Thiel, "Exact medial axis with Euclidean distance" *Image and Vision Computing* **23** (2005) 167-175.
- [8] D. Coeurjolly, "Algorithmique et géométrie discrète pour la caractérisation des courbes et des surfaces," Ph.D. dissertation, Université Lumière Lyon 2, Bron, Laboratoire ERIC, Dec. 2002.
<david.coeurjolly@liris.cnrs.fr>, <http://liris.cnrs.fr/~dcoeurjo/Code/DistanceTransformation>



Figure 1. The original 16bit data from the micro-CT were segmented into enamel, dentin and the pulp tissue based on their different radiographic absorption. Enamel is rendered in cyan, dentin in yellow and the pulp is visualized in red with ParaView 2.4 (Kitware, www.kitware.com).




Figure 2. The segmented pulp volume rendered with the VolumeJ plugin for ImageJ (v. 1.7a, Michael Abramoff). The local thickness is mapped to the surface of the pulp. This information is used to develop optimized root canal instruments (diameter and taper). During a root canal treatment infected tooth tissue has to be removed. The goal of an endodontic treatment is to keep the original shape of the pulp canal, but instrument it down to the apex without obliteration and without iatrogenic damages.