

Perspective

Nitric oxide-mediated epigenetic mechanisms in developing neurons

Alexi Nott and Antonella Riccio*

MRC Laboratory for Molecular and Cell Biology; and Department of Neuroscience, Physiology and Pharmacology; University College London; London, UK

Key words: epigenetics, neuron, transcription, nitric oxide, histone deacetylase, neuronal development, plasticity, differentiation

Epigenetic changes of chromatin are increasingly recognized as key modifications that dictate the differentiation state of cells during development. Within the central nervous system, extracellular cues induce chromatin remodelling events that are essential for neuronal progenitor proliferation, cell differentiation and, later, plasticity. In this review, we discuss recent studies that show how extracellular and intranuclear signals influence chromatin remodelling and neuron-specific gene expression. The gaseous molecule Nitric Oxide (NO) has recently emerged as a new key player that mediates the epigenetic changes associated with cell cycle arrest and differentiation in neurons. Histone deacetylases (HDACs) are the first identified intranuclear targets of NO, but, due to its highly diffusible nature, it is likely that many other nuclear factors are directly regulated by NO.

Introduction

The development of the nervous system requires the generation of a large number of neuronal cell types from a single population of progenitor cells. Initial cell fate determination of neuronal precursors is followed by neuronal migration to target destinations, axonal elongation and establishment of contacts with specific target cells. All these events are associated with a tightly regulated temporal and spatial expression of transcription factors, which is mediated by both extrinsic signals, including Notch, ephrins and neurotrophins, and intrinsic factors, such as chromatin remodelling factors.¹⁻³ It is now recognized that epigenetic modifications of DNA and histone proteins influence the overall structure of chromatin, regulating the access of transcription factors to gene promoters and driving transcription.⁴⁻⁷

Adrian Bird has recently proposed a revised definition of epigenetics as “the structural adaptation of chromosomal regions so as to register, signal or perpetuate altered activity states”.⁸ DNA is packaged into chromatin through its association with protein complexes called nucleosomes. A nucleosomal unit comprises 146 base pairs of DNA wrapped around an octameric histone complex composed of one H3/H4 tetramer and two H2A/H2B dimers.⁹ Nucleosomes are

connected by 20–60 base pairs of DNA associated with the linker histone H1. Chromatin exists either as highly condensed heterochromatin or as the less compact euchromatin. Heterochromatin is late-replicating, rich in repetitive sequences, and transcriptionally silent, whereas euchromatin organisation is more dynamic and can change transiently, allowing for gene activation and repression in response to extracellular cues.

Epigenetic regulation of chromatin involves covalent modification of both DNA and histone proteins.^{4,7} Histones can undergo a large number of posttranslational covalent modifications at their N-terminal tails, including acetylation, methylation, phosphorylation, ADP-ribosylation, sumoylation and ubiquitination.⁴⁻⁷ Histone acetylation, the most widely studied epigenetic modification, involves the cleavage of acetyl groups from acetyl-coenzyme A and its addition to the ϵ -NH⁺ of lysine residues. Acetylation of histones is catalysed by histone acetyl-transferases (HATs), which are divided into three main families, CBP/p300, MYST and GNAT and exhibit virtually no preference for specific lysine residues.^{10,11} Histone acetylation directly influences chromatin structure by reducing the positive charge on lysine residues and decreasing the affinity of histones for negatively charged DNA. This allows a localized ‘unravelling’ of chromatin, making it more accessible for binding of further co-activators and the basal transcription machinery. Histone acetylation also provides a platform for chromatin remodelling proteins that contain bromo-domains, such as the SWI/SNF complex.^{7,12} The assembly of these large multiprotein complexes is important for the initiation of transcription, DNA replication and repair and further histone modifications.

Conversely, the removal of acetyl groups is catalysed by histone deacetylases (HDACs) and is associated with chromatin condensing and transcriptional repression.¹³⁻¹⁸ There are four classes of HDACs, class I (HDACs 1–3 and 8), class II (HDACs 4–7, 9 and 10) and class IV (HDAC 11) that are all zinc dependent, whereas class III HDACs (SIRT 1–7) require the cofactor NAD⁺.^{16,19} Class I HDACs show high homology to the yeast Reduced potassium dependency-3 (Rpd3) and, with the exception of HDAC3, are ubiquitously expressed and principally localized in the nucleus. Class II HDACs are homologous to yeast Histone deacetylase-1 (Hda-1). They exhibit a tissue-specific expression and shuttle between the nucleus and cytoplasm (as does HDAC3). HDAC11, the sole member of class IV HDACs identified so far, shows an intermediate homology to both Rpd3 and Hda1. Sirtuins (class III HDACs), are homologous to silent information regulator-2 (sir2) and are evolutionarily distinct from classes I, II and IV HDACs.

*Correspondence to: Antonella Riccio; MRC Laboratory for Molecular Cell Biology; University College London; Gower Street; London WC1E 6BT UK; Email: a.riccio@ucl.ac.uk

Submitted: 12/19/08; Revised: 01/08/09; Accepted: 01/08/09

Previously published online as a *Cell Cycle* E-publication:
<http://www.landesbioscience.com/journals/cc/article/7805>




Figure 1. Neuronal progenitor cells exist in a proliferative state, and terminally differentiate to postmitotic neurons upon exposure to extrinsic stimuli, such as neurotrophins, ephrins, notch signaling and synaptic activity (upper). In proliferating neuronal precursors, HDAC2 is recruited to neural gene promoters through interaction with repressive complexes (REP), which include the corepressors HDAC1, REST, CoREST and MeCP2. Recruitment of HDAC2 to chromatin maintains local histones in a deacetylated state and is associated with transcriptional repression. Following extracellular stimulation NO accumulates within the nucleus, and S-nitrosylated HDAC2 is released from repressive complexes (lower). Dissociation of HDAC2 from neural gene promoters, together with the recruitment of transcription factors and coactivators, such as CREB and CBP, induces histone acetylation and a shift of chromatin to an active state. This event initiates the expression of many neuronal genes, including *c-fos*, *VGF*, *Egr1*, *Nos1* and *Bcl2*.

Epigenetic Mechanisms in the Developing Nervous System

The expression of neural genes during development requires the turnover of large transcriptional complexes that often contain either HATs or HDACs.²⁰ During the early stages of neuronal development, progenitor cells are maintained in a proliferative state through expression of Notch effectors. These include inhibitory basic helix-loop-helix (bHLH) transcription factors, such as *Hes1*, that act in concert with co-repressive complexes to inhibit neurogenic bHLH genes, such as *Ascl1*.²¹ The repressive activity of *Hes1* is mediated by the recruitment of Sin3A/HDAC complexes to gene promoters that results in localized histone hypoacetylation and gene silencing.²² Exposure of progenitor cells to neurogenic stimuli leads to ADP-ribosylation of the *Hes1* complex, which induces the dissociation of Sin3A/HDAC complex from *Hes1*, recruitment of HATs and hyperacetylation of local histone proteins.²³ The switch of *Hes1* from a transcriptional repressor to an activator is dependent on the presence of HDACs and HATs within the complex, suggesting that the conventional distinction between nuclear repressors and activators is becoming obsolete. The nuclear factor CREST for example, was initially described as a calcium-responsive transcriptional activator that regulates activity-dependent dendritic development.²⁴ Recent studies however, have shown that CREST is tightly associated with the nBAF chromatin remodelling complexes, and can either activate or repress transcription by recruiting specific cofactors to gene promoters.²⁵ In resting neurons, CREST interacts with the nBAF subunit Brg1 and represses transcription by recruiting HDAC1 and Rb complexes.²⁶ Following exposure of cortical culture to depolarizing conditions, Rb and HDAC1 are released from gene promoters and the CREST/Brg1 complexes associate with CBP, inducing histone hyperacetylation and gene expression.

Many neuronal genes contain a conserved motif within their promoters called the Repressor Element 1 (RE1) that upon methylation binds the zinc-finger transcriptional repressor RE1 silencing transcription factor (REST).^{27,28} REST recruits Sin3A/HDAC, as well as the repressive cofactors CoREST and MeCP2, to sites of gene repression. CoREST is a co-repressor that mediates long-term gene silencing by recruitment of HDACs, H3K9 histone methyltransferases and the H3K4 demethylase LSD1.^{29,30} REST silences many neuron-specific genes in non-neuronal cells, including those encoding ion channels, neurotransmitter receptors, synaptic vesicle proteins and adhesion proteins.³¹ Transcriptional and translational regulation of REST allows a tight control of neuron-specific genes during development, and later influences the plasticity of mature neurons.³² High levels of REST in neuronal embryonic stem cells prevent their exit from the cell cycle and blocks premature neuronal differentiation.³³ As neuronal development proceeds, the *Rest* gene itself is transcriptionally and translationally repressed, allowing precursor cells to differentiate into mature neurons.³³ The absence of REST/HDAC/Sin3A/CoREST repressive complexes from gene promoters induces the expression of neuronal genes.³⁴ However, the transcription of certain genes, including *Bdnf* and calbindin, remains low due to CoREST and MeCP2 complex binding to additional methylation sites.³³ Following neuronal depolarisation, MeCP2 is phosphorylated, possibly decreasing its binding affinity for methylated DNA and *Bdnf* transcription is induced.³⁵ Regulation of the binding of MeCP2 repressive complexes to neural gene promoters in response to neuronal activity provides a platform for synaptic plasticity in mature neurons.³²

Differentiation of progenitor cells into a neuronal phenotype is also regulated by the nuclear receptor co-repressor (N-CoR).^{36,37}

Repressive complexes containing N-CoR silence gene promoters of down-stream effectors of Notch signaling through the recruitment of HDAC3.³⁶ The absence of N-CoR in cortical progenitors prevents FGF-dependent self-renewal of neuronal progenitors and promotes their differentiation into astrocytes.³⁸ Interestingly, Ciliary Neurotrophic Factor (CNTF) triggers glial differentiation and leads to Akt1 kinase-dependent phosphorylation of N-CoR and subsequent relocalization to the cytoplasm.³⁸ These data imply that N-CoR/HDAC3 complexes play an important role in regulating the differentiation of early-stage progenitor cells by selectively repressing gliogenic genes.

Nitric Oxide Regulates Neurogenesis and Neuronal Development

Nitric Oxide (NO) is an important mediator of two pivotal neuronal developmental processes, proliferation and differentiation. The primary source of NO in neurons is the neuronal isoform of nitric oxide synthase (nNOS). After the discovery that NO acts as a physiological signaling molecule, it became apparent that cysteine thiol groups can couple NO to form *S*-nitrosylated proteins, a modification akin to phosphorylation.³⁹ *S*-nitrosylation has been reported to regulate the function of many proteins, affecting catalytic activity, protein-protein interaction and/or subcellular localisation.⁴⁰⁻⁴² Anti-proliferative effects of NO have been demonstrated in cerebellar granule cells, as well as in several neuronal-derived cell lines, where NO induces cell cycle arrest without affecting cell viability or apoptosis.⁴³⁻⁴⁶ The observation that NO triggers growth arrest in PC12 cells provided the initial indication that NO is an important regulator of neuronal development.^{43,47} Similarly, reduced proliferation and increased differentiation was observed in neuroblastoma cell lines and in cerebellar neurons upon exposure to endogenous and exogenous sources of NO.^{44,45,48} Interestingly, stimulation of mouse cortical neuroepithelial cluster cultures with either NO donors or BDNF inhibits cell proliferation and promotes differentiation, possibly by increasing nNOS levels, thereby providing a positive feedback loop.⁴⁸ Experiments performed in vitro by using embryonic brain slice cultures, or in postnatal rats in vivo have shown that exposure to NOS inhibitors impairs cerebellar granule cell migration and increases proliferation.^{49,50}

In the adult brain, neurogenesis persists within two regions, the sub ventricular zone (SVZ) and the subgranular zone of the dentate gyrus in the hippocampus. Discrete populations of neurons expressing nNOS are found in close proximity to these highly proliferative regions.^{51,52} Neuroblasts from the SVZ migrate tangentially along the rostral migratory stream (RMS) to the olfactory bulb where they differentiate to form mature interneurons. NOS positive cells are found along the RMS, with abundant processes that intermingle with the migrating precursor cells.⁵¹ Interestingly, NOS-negative progenitors in the adult SVZ begin to express NOS as they start migrating to the olfactory bulb, implicating that NO may play a role in regulating this process. Genetic deletion of nNOS, as well as systemic and intraventricular infusion of nNOS inhibitors in adult mice, augmented the number of cells generated within the SVZ and the dentate gyrus.⁵³⁻⁵⁵ Primary cultures of the SVZ obtained from nNOS null mice have also demonstrated an inhibitory effect of NO on proliferation without influencing cell survival.^{54,56}

In the developing nervous system, nNOS is transiently expressed in restricted regions of the brain, including the embryonic cortical plate, and the olfactory epithelium.⁵⁷ Within these populations, nNOS is not detectable during the early phases of neurogenesis, when neural stem cells are actively proliferating. nNOS mRNA is first detected in neural precursor cells at E10, with levels progressively increasing as precursors later differentiate to form post-mitotic mature neurons.^{48,57,58} Neurotrophin-dependent precursors in sensory and sympathetic ganglia also express high levels of nNOS, as they exit the cell cycle and start to differentiate.^{57,59} However, nNOS expression dramatically decreases postnatally, once peripheral neuron axons have established connections with their targets.⁵⁹ Importantly, stimulation of cortical neurons with the neurotrophic factor BDNF increases expression of nNOS and induces NO accumulation within both cytoplasmic and nuclear compartments.^{48,60,61}

Transcriptional Regulatory Targets of NO

Several studies have demonstrated that NO modulates gene expression through the activation of multiple transcription factors including CREB, N-Myc, NFκB, p53 and in bacteria, OxyR.⁶²⁻⁶⁵ A role of NO in regulating gene expression was first demonstrated in the neuronal-derived cell line, PC12. Structure to function analysis of the *c-fos* promoter demonstrated that Ca²⁺ signaling induced NO synthesis and CREB-dependent *c-fos* expression.⁶⁶ Other studies have shown that NO activation of CREB regulates neuronal survival, differentiation and plasticity.^{60,67-70} However, the mechanisms by which NO regulates CREB activation remain unclear. In cerebellar granule cells, NO-dependent phosphorylation of CREB through the sGC/PKG pathway has been shown to be required for CREB activation and for promoting neuronal survival.^{69,71} In contrast, NO produced following neurotrophin stimulation of cortical neurons mediates CREB activity through mechanisms that are completely independent of CREB phosphorylation and involve the recruitment of active CREB to target gene promoters.⁶⁰ More importantly, no changes of CREB phosphorylation were observed in cortical neurons obtained from nNOS null mice and exposed to either BDNF or depolarizing stimuli.⁶⁰

An important target of NO is the oncogene N-Myc, a transcription factor that is essential for the expansion of neuronal precursors during normal brain development.⁷² NO signaling inhibits N-Myc expression through the cGMP/PKG pathway, decreasing proliferation rates of neonatal precursor cerebellar granule neurons, neuroblastoma cell lines and embryonic carcinoma stem cells.^{45,50,73} Despite the many evidences that NO regulates the transcriptional activity of CREB and N-Myc, they are not target of *S*-nitrosylation, and the molecular mechanisms that regulate their activation remain unclear.

S-nitrosylation, however, mediates NO-dependent regulation of various zinc-finger-containing transcription factors, including egr-1 and NFκB.^{74,75} *S*-nitrosylation of zinc-sulphur clusters of egr-1 and NFκB disrupts zinc binding, inhibiting their association with DNA.⁷⁶⁻⁷⁸ In mammalian cells, NFκB is constitutively expressed and is retained within the cytoplasm by its inhibitory cofactor IκB. Upon IκB phosphorylation and subsequent proteolysis, NFκB is free to translocate to the nucleus to bind target gene promoters.⁶⁴ *S*-nitrosylation and tyrosine nitration of NFκB inhibits its binding to DNA.^{79,80} Furthermore, *S*-nitrosylation of IκB prevents its phosphorylation thereby increasing cytoplasmic retention of NFκB.^{81,82}

Other studies however, have suggested that NO signaling induces the expression of NF κ B-regulated genes.^{83,84} Similar conflicting evidences have been observed for the transcription factor AP1. NO has been demonstrated to inhibit AP1 binding to DNA, through modification of its two components, *c-fos* and *c-jun*.^{85,86} In contrast, NO signaling induces expression of the *c-fos*-dependent gene, tyrosine hydroxylase, as well as *c-fos* itself, in a number of neuronal systems.^{60,87-89} This may serve as a negative feedback mechanism, as NO initially induces expression of *c-fos* and later inhibits AP1 activity through S-nitrosylation of *c-fos* and *c-jun*.

A common link between the physiological processes regulated by NO during the development of the nervous system is the requirement for profound and long-lasting changes of gene expression. Exposure of cells to NO donors induces the transcription of a large number of genes through the binding of transcription factors to various cis-activating elements, including CRE, AP1, AP2, BRN-3, EGR, E2F1 and Sp1.^{90,91} NO regulates a broad range of functionally distinct genes involved in cell signaling, cell cycle, transcription, stress response, protein degradation, oxidative stress and apoptosis.⁹⁰ The large number of transcription factors activated and the diversity of genes induced suggest that NO is likely to signal at an epigenetic level, reflecting the broad impact that NO has on neuronal physiology.

Regulation of HDAC Function by NO Signaling

Neurotrophin stimulation and NO signaling are important modulators of cortical neurogenesis. Both BDNF and NO inhibit proliferation and promote neuronal differentiation.^{48,53,55,92,93} BDNF stimulation of cortical neurons induces accumulation of NO within the cytoplasmic and nuclear compartments.⁶¹ Furthermore, acetylation of histones H3 and H4 increases following BDNF treatment of cortical neurons in a NO-dependent manner.^{60,61} Recently, the chromatin-modifying enzyme HDAC2 has been identified as a key nuclear target of NO.^{61,94} S-nitrosylation of HDAC2 occurs at two cysteine residues, Cys262 and Cys274, and does not affect its deacetylase activity in neurons. Instead, HDAC2 S-nitrosylation induces its dissociation from CREB-regulated gene promoters. This event leads to an increase of histone acetylation at specific promoter regions and transcription of many genes, including *c-fos*, *egr1*, *VGF* and *nNos*.^{60,61} Importantly, NO-dependent modulation of HDAC2 function is necessary for dendritic growth in vitro, suggesting a key role of HDAC2 S-nitrosylation in cortical development.⁶¹

NO-dependent inhibition of HDAC2 function has also been reported in muscle cells.⁹⁴ In a mouse model of muscular dystrophy, NO-dependent inhibition of HDAC2 has been exploited as a potential novel therapeutic approach for this genetic disease.⁹⁴ Interestingly, in muscle cells, S-nitrosylation of HDAC2 on yet unidentified cysteine residues decreases its deacetylase activity,⁹⁴ although in neurons HDAC2 enzymatic activity is unchanged.⁶¹ This discrepancy may reflect a difference of cysteine residue(s) targeted by NO in muscle cells and in neurons. For example, in muscle cells HDAC2 S-nitrosylation may occur at Cys152, which is located within the deacetylase active site. It should also be considered that the transcriptional profile between muscle cells and neurons differs greatly, possibly involving the recruitment of HDAC2 to repressive complexes of cell-type specific genes through distinct mechanisms. With the exception of HDAC8, the deacetylase activity of HDACs

is dependent on protein-protein interaction with corepressors.^{95,96} Similarly, HDACs do not directly bind DNA and relies on the recruitment by corepressors to gene promoters. An attractive hypothesis is that S-nitrosylation of HDAC2 results in its dissociation from cell-type specific repressive complexes,²⁶ which may influence its association with chromatin and/or deacetylase activity (Fig. 1).

Regulation of deacetylase function by NO is not restricted to class I HDACs. In endothelial cells, a shear-stress response enhances NO synthesis and inhibits gene expression.^{97,98} Treatment of Human Umbilical Vein Endothelial Cells (HUVECs) with NO donors induced nuclear localisation of HDAC4 and HDAC5 and class II HDAC-dependent hypoacetylation of histone H3.⁹⁹ Nuclear localisation of class II HDACs is inhibited following phosphorylation by calcium-calmodulin dependent kinases (CaMKs) and its association with the cytoplasmic chaperone, 14-3-3.^{100,101} Exposure of HDAC4 to NO donors increases both its association with the protein phosphatase PP2A⁹⁹ and the phosphatase activity of PP2A, allowing HDAC4 to translocate to the nucleus. Interestingly, in neurons, NO inhibits deacetylase activity of cytoplasmic HDACs, although nuclear HDAC activity remains unaffected.¹⁰² These results imply that NO signaling in neuronal cells specifically affects the enzymatic activity of cytoplasmic class II HDACs, possibly influencing deacetylation of non-histone proteins.

Conclusions

The finding that NO regulates many aspects of gene expression in mammalian neurons has implications that reach beyond the regulation of cortical development, and include novel therapeutic approaches for both neurodegenerative disorders and cancer. Regulation of histone acetylation has been implicated in the pathogenesis of several neurodegenerative disorders, including Huntington's, Parkinson's and Alzheimer's diseases.¹⁰³⁻¹⁰⁵ These neurodegenerative disorders are associated with the formation of cytoplasmic aggregates that sequester intracellular proteins, interfering with various signaling pathways.^{106,107} It is now apparent that inclusion of transcriptional cofactors within these aggregates profoundly affects histone modification and gene expression.^{104,108} Interestingly, dysregulation of HDAC2 binding to neuronal gene promoters further exacerbates the transcriptional changes observed in neurons obtained from a mouse model of Huntington's disease.^{109,110}

As HDACs play an important role in regulating cell proliferation, differentiation and apoptosis, it is not surprising that they have been a major focus for cancer therapeutics for the past 10 years.¹¹¹⁻¹¹³ HDAC2 is highly expressed in gastric carcinomas, colorectal carcinomas, cervical dysplasias and endometrial stromal sarcomas.¹¹⁴⁻¹¹⁶ Dysregulated expression of HDAC2 in cancers is associated with reduced transcription of antiproliferative genes such as the cyclin-dependent kinase inhibitor p21.^{117,118} NO-dependent regulation of HDAC2 binding affinity for DNA provides an alternative avenue for future drug design that does not require inhibition of intrinsic deacetylase activity. Novel therapeutic approaches may target the chromatin-remodelling function of HDAC2, leaving deacetylation of non-chromatin associated factors intact, thereby providing an avenue for the development of more specific drugs.

The finding that a gaseous molecule, such as NO, has such a profound effect on the transcriptional profile of mammalian, yeast and bacterial cells,^{62,76,90,91,119} by directly modifying a large number

of nuclear proteins,⁶¹ suggest that NO may act as a master regulator of gene expression. Future research will reveal to which extent chromatin remodelling factors and nuclear events are functionally regulated by NO.

Acknowledgements

This work was supported by the Medical Research Council (Research Grant G0500792) and the European Research Council (Marie Curie International Reintegration Grant MCIRG-CT-2005-016501). A.R. is the recipient of an MRC Career Development Award (G117/533).

References

- Bertrand N, Castro DS, Guillemot F. Proneural genes and the specification of neural cell types. *Nat Rev Neurosci* 2002; 3:517-30.
- Gupta A, Tsai L, Wynshaw-Boris A. Life is a journey: a genetic look at neocortical development. *Nat Rev Genet* 2002; 3:342-55.
- Hevner RF. From radial glia to pyramidal-projection neuron: transcription factor cascades in cerebral cortex development. *Mol Neurobiol* 2006; 33:33-50.
- Strahl BD, Allis CD. The language of covalent histone modifications. *Nature* 2000; 403:41-5.
- Zhang Y, Reinberg D. Transcription regulation by histone methylation: interplay between different covalent modifications of the core histone tails. *Genes Dev* 2001; 15:2343-60.
- Berger SL. Histone modifications in transcriptional regulation. *Curr Opin Genet Dev* 2002; 12:142-8.
- Kouzarides T. Chromatin modifications and their function. *Cell* 2007; 128:693-705.
- Bird A. Perceptions of epigenetics. *Nature* 2007; 447:396-8.
- Luger K, Mäder AW, Richmond RK, Sargent DF, Richmond TJ. Crystal structure of the nucleosome core particle at 2.8 Å resolution. *Nature* 1997; 389:251-60.
- Roth SY, Denu JM, Allis CD. Histone acetyltransferases. *Annu Rev Biochem* 2001; 70:81-120.
- Carrozza MJ, Utley RT, Workman JL, Côté J. The diverse functions of histone acetyltransferase complexes. *Trends Genet* 2003; 19:321-9.
- Zeng L, Zhou MM. Bromodomain: an acetyl-lysine binding domain. *FEBS Lett* 2002; 513:124-8.
- Cress WD, Seto E. Histone deacetylases, transcriptional control, and cancer. *J Cell Physiol* 2000; 184:1-16.
- Ng HH, Bird A. Histone deacetylases: silencers for hire. *Trends Biochem Sci* 2000; 25:121-6.
- Grozinger CM, Schreiber SL. Deacetylase enzymes: biological functions and the use of small-molecule inhibitors. *Chem Biol* 2002; 9:3-16.
- Thiagalingam S, Cheng KH, Lee HJ, Mineva N, Thiagalingam A, Ponte JF. Histone deacetylases: unique players in shaping the epigenetic histone code. *Ann NY Acad Sci* 2003; 983:84-100.
- Verdin E, Dequiedt F, Kasler HG. Class II histone deacetylases: versatile regulators. *Trends Genet* 2003; 19:286-93.
- Yang X, Seto E. Collaborative spirit of histone deacetylases in regulating chromatin structure and gene expression. *Curr Opin Genet Dev* 2003; 13:143-53.
- Yang X, Grégoire S. Class II histone deacetylases: from sequence to function, regulation and clinical implication. *Mol Cell Biol* 2005; 25:2873-84.
- Allen ND. Temporal and epigenetic regulation of neurodevelopmental plasticity. *Philos Trans R Soc Lond B Biol Sci* 2008; 363:23-38.
- Cau E, Gradwohl G, Casarosa S, Kageyama R, Guillemot F. Hes genes regulate sequential stages of neurogenesis in the olfactory epithelium. *Development* 2000; 127:2323-32.
- Nuthall HN, Joachim K, Stifani S. Phosphorylation of serine 239 of Groucho/TLE1 by protein kinase CK2 is important for inhibition of neuronal differentiation. *Mol Cell Biol* 2004; 24:8395-407.
- Ju BG, Solum D, Song EJ, Lee KJ, Rose DW, Glass CK, et al. Activating the PARP-1 sensor component of the groucho/TLE1 corepressor complex mediates a CaMK kinase II-dependent neurogenic gene activation pathway. *Cell* 2004; 119:815-29.
- Aizawa H, Hu SC, Bobb K, Balakrishnan K, Ince G, Gurevich I, et al. Dendrite development regulated by CREST, a calcium-regulated transcriptional activator. *Science* 2004; 303:197-202.
- Wu JI, Lessard J, Olave IA, Qiu Z, Ghosh A, Graef IA, et al. Regulation of dendritic development by neuron-specific chromatin remodeling complexes. *Neuron* 2007; 56:94-108.
- Qiu Z, Ghosh A. A calcium-dependent switch in a CREST-BRG1 complex regulates activity-dependent gene expression. *Neuron* 2008; 60:775-87.
- Chong JA, Tapia-Ramírez J, Kim S, Toledo-Aral JJ, Zheng Y, Boutros MC, et al. REST: a mammalian silencer protein that restricts sodium channel gene expression to neurons. *Cell* 1995; 80:949-57.
- Schoenherr CJ, Anderson DJ. The neuron-restrictive silencer factor (NRSF): a coordinate repressor of multiple neuron-specific genes. *Science* 1995; 267:1360-3.
- Shi Y, Sawada J, Sui G, Affar el B, Whetstone JR, Lan F, et al. Coordinated histone modifications mediated by a CtBP co-repressor complex. *Nature* 2003; 422:735-8.
- Shi Y, Lan F, Matson C, Mulligan P, Whetstone JR, Cole PA, et al. Histone demethylation mediated by the nuclear amine oxidase homolog LSD1. *Cell* 2004; 119:941-53.
- Johnson DS, Mortazavi A, Myers RM, Wold B. Genome-wide mapping of in vivo protein-DNA interactions. *Science* 2007; 316:1497-502.
- Ballas N, Mandel G. The many faces of REST oversee epigenetic programming of neuronal genes. *Curr Opin Neurobiol* 2005; 15:500-6.
- Ballas N, Grunseich C, Lu DD, Speh JC, Mandel G. REST and its corepressors mediate plasticity of neuronal gene chromatin throughout neurogenesis. *Cell* 2005; 121:645-57.
- Schoenherr CJ, Paquette AJ, Anderson DJ. Identification of potential target genes for the neuron-restrictive silencer factor. *Proc Natl Acad Sci USA* 1996; 93:9881-6.
- Zhou Z, Hong EJ, Cohen S, Zhao WN, Ho HY, Schmidt L, et al. Brain-specific phosphorylation of MeCP2 regulates activity-dependent Bdnf transcription, dendritic growth and spine maturation. *Neuron* 2006; 52:255-69.
- Kao HY, Ordentlich P, Koyano-Nakagawa N, Tang Z, Downes M, Kintner CR, et al. A histone deacetylase corepressor complex regulates the Notch signal transduction pathway. *Genes Dev* 1998; 12:2269-77.
- Jepsen K, Rosenfeld MG. Biological roles and mechanistic actions of co-repressor complexes. *J Cell Sci* 2002; 115:689-98.
- Hermanson O, Jepsen K, Rosenfeld MG. N-CoR controls differentiation of neural stem cells into astrocytes. *Nature* 2002; 419:934-9.
- Stamler JS, Simon DI, Osborne JA, Mullins ME, Jaraki O, Michel T, et al. S-nitrosylation of proteins with nitric oxide: synthesis and characterization of biologically active compounds. *Proc Natl Acad Sci USA* 1992; 89:444-8.
- Stamler JS. Redox signaling: nitrosylation and related target interactions of nitric oxide. *Cell* 1994; 78:931-6.
- Lipton S, Choi YB, Takahashi H, Zhang D, Li W, Godzik A, et al. Cysteine regulation of protein function—as exemplified by NMDA-receptor modulation. *Trends Neurosci* 2002; 25:474-80.
- Hess D, Matsumoto A, Kim SO, Marshall HE, Stamler J. Protein S-nitrosylation: purview and parameters. *Nat Rev Mol Cell Biol* 2005; 6:150-66.
- Peunova N, Enikolopov G. Nitric oxide triggers a switch to growth arrest during differentiation of neuronal cells. *Nature* 1995; 375:68-73.
- Murillo-Carretero M, Ruano MJ, Matarredona ER, Villalobo A, Estrada C. Antiproliferative effect of nitric oxide on epidermal growth factor-responsive human neuroblastoma cells. *J Neurochem* 2002; 83:119-31.
- Ciani E, Severi S, Contestabile A, Bartsaghi R, Contestabile A. Nitric oxide negatively regulates proliferation and promotes neuronal differentiation through N-Myc downregulation. *J Cell Sci* 2004; 117:4727-37.
- Peunova N, Scheinker V, Ravi K, Enikolopov G. Nitric oxide coordinates cell proliferation and cell movements during early development of *Xenopus*. *Cell Cycle* 2007; 6:3132-44.
- Phung YT, Bekker JM, Hallmark OG, Black SM. Both neuronal NO synthase and nitric oxide are required for PC12 cell differentiation: a cGMP independent pathway. *Brain Res Mol Brain Res* 1999; 64:165-78.
- Cheng A, Wang S, Cai J, Rao MS, Mattson MP. Nitric oxide acts in a positive feedback loop with BDNF to regulate neural progenitor cell proliferation and differentiation in the mammalian brain. *Dev Biol* 2003; 258:319-33.
- Tanaka M, Yoshida S, Yano M, Hanaoka F. Roles of endogenous nitric oxide in cerebellar cortical development in slice cultures. *Neuroreport* 1994; 5:2049-52.
- Ciani E, Calvanese V, Crochemore C, Bartsaghi R, Contestabile A. Proliferation of cerebellar precursor cells is negatively regulated by nitric oxide in newborn rat. *J Cell Sci* 2006; 119:3161-70.
- Moreno-López B, Noval JA, González-Bonet LG, Estrada C. Morphological bases for a role of nitric oxide in adult neurogenesis. *Brain Res* 2000; 869:244-50.
- Islam AT, Kuraoka A, Kawabuchi M. Morphological basis of nitric oxide production and its correlation with the polysialylated precursor cells in the dentate gyrus of the adult guinea pig hippocampus. *Anat Sci Int/Japanese Association of Anatomists* 2003; 78:98-103.
- Moreno-López B, Romero-Grimaldi C, Noval JA, Murillo-Carretero M, Matarredona ER, Estrada C. Nitric oxide is a physiological inhibitor of neurogenesis in the adult mouse subventricular zone and olfactory bulb. *J Neurosci* 2004; 24:85-95.
- Matarredona ER, Murillo-Carretero M, Moreno-López B, Estrada C. Nitric oxide synthesis inhibition increases proliferation of neural precursors isolated from the postnatal mouse subventricular zone. *Brain Res* 2004; 995:274-84.
- Packer MA, Stasiv Y, Benraiss A, Chmielnicki E, Grinberg A, Westphal H, et al. Nitric oxide negatively regulates mammalian adult neurogenesis. *Proc Natl Acad Sci USA* 2003; 100:9566-71.
- Torroglosa A, Murillo-Carretero M, Romero-Grimaldi C, Matarredona E, Campos-Caro A, Estrada C. Nitric oxide decreases subventricular zone stem cell proliferation by inhibition of epidermal growth factor receptor and phosphoinositide-3-kinase/Akt pathway. *Stem Cells* 2007; 25:88-97.
- Bredt DS, Snyder SH. Transient nitric oxide synthase neurons in embryonic cerebral cortical plate, sensory ganglia and olfactory epithelium. *Neuron* 1994; 13:301-13.
- Ogura T, Nakayama K, Fujisawa H, Esumi H. Neuronal nitric oxide synthase expression in neuronal cell differentiation. *Neurosci Lett* 1996; 204:89-92.
- Thippeswamy T, McKay JS. Neuronal nitric oxide synthase and nerve growth factor expression in the enteric nervous system. *Cell Mol Biol (Noisy-le-grand)* 2005; 51:293-8.
- Riccio A, Alvania R, Lonze B, Ramanan N, Kim T, Huang Y, et al. A nitric oxide signaling pathway controls CREB-mediated gene expression in neurons. *Mol Cell* 2006; 21:283-94.

61. Nott A, Watson P, Robinson J, Crepaldi L, Riccio A. S-nitrosylation of histone deacetylase 2 induces chromatin remodelling in neurons. *Nature* 2008; 455:411-5.
62. Hausladen A, Privalle CT, Keng T, DeAngelo J, Stamler JS. Nitrosative stress: activation of the transcription factor OxyR. *Cell* 1996; 86:719-29.
63. Marshall HE, Merchant K, Stamler JS. Nitrosation and oxidation in the regulation of gene expression. *FASEB J* 2000; 14:1889-900.
64. Bogdan C. Nitric oxide and the regulation of gene expression. *Trends Cell Biol* 2001; 11:66-75.
65. Contestabile A. Regulation of transcription factors by nitric oxide in neurons and in neural-derived tumor cells. *Prog Neurobiol* 2008; 84:317-28.
66. Peunova N, Enikolopov G. Amplification of calcium-induced gene transcription by nitric oxide in neuronal cells. *Nature* 1993; 364:450-3.
67. Lu YF, Kandel ER, Hawkins RD. Nitric oxide signaling contributes to late-phase LTP and CREB phosphorylation in the hippocampus. *J Neurosci* 1999; 19:10250-61.
68. Puzzo D, Vitolo O, Trinchese F, Jacob JP, Palmeri A, Arancio O. Amyloid-beta peptide inhibits activation of the nitric oxide/cGMP/cAMP-responsive element-binding protein pathway during hippocampal synaptic plasticity. *J Neurosci* 2005; 25:6887-97.
69. Ciani E, Guidi S, Bartesaghi R, Contestabile A. Nitric oxide regulates cGMP-dependent cAMP-responsive element binding protein phosphorylation and Bcl-2 expression in cerebellar neurons: implication for a survival role of nitric oxide. *J Neurochem* 2002; 82:1282-9.
70. Crepaldi L, Riccio A. Chromatin learns to behave. *Epigenetics* 2009; In press.
71. Nagai-Kusuhara A, Nakamura M, Mukuno H, Kanamori A, Negi A, Seigel GM. cAMP-responsive element binding protein mediates a cGMP/protein kinase G-dependent anti-apoptotic signal induced by nitric oxide in retinal neuro-glial progenitor cells. *Exp Eye Res* 2007; 84:152-62.
72. Knoepfler PS, Kenney AM. Neural precursor cycling at sonic speed: N-Myc pedals, GSK-3 brakes. *Cell Cycle* 2006; 5:47-52.
73. Park SW, Li J, Loh HH, Wei LN. A novel signaling pathway of nitric oxide on transcriptional regulation of mouse kappa opioid receptor gene. *J Neurosci* 2002; 22:7941-7.
74. Thiriet N, Esteve L, Aunis D, Zwiler J. Immediate early gene induction by natriuretic peptides in PC12 pheochromocytoma and C6 glioma cells. *Neuroreport* 1997; 8:399-402.
75. Cibelli G, Policastro V, Rössler OG, Thiel G. Nitric oxide-induced programmed cell death in human neuroblastoma cells is accompanied by the synthesis of Egr-1, a zinc finger transcription factor. *J Neurosci Res* 2002; 67:450-60.
76. Kröncke KD, Fehsel K, Schmidt T, Zenke FT, Dasting I, Wesener JR, et al. Nitric oxide destroys zinc-sulfur clusters inducing zinc release from metallothionein and inhibition of the zinc finger-type yeast transcription activator LAC9. *Biochem Biophys Res Commun* 1994; 200:1105-10.
77. Misra RR, Hochadel JF, Smith GT, Cook JC, Waalkes MP, Wink DA. Evidence that nitric oxide enhances cadmium toxicity by displacing the metal from metallothionein. *Chem Res Toxicol* 1996; 9:326-32.
78. Aravindakumar CT, Ceulemans J, De Ley M. Nitric oxide induces Zn²⁺ release from metallothionein by destroying zinc-sulphur clusters without concomitant formation of S-nitrosothiol. *Biochem J* 1999; 344:253-8.
79. Marshall HE, Stamler JS. Inhibition of NFkappaB by S-nitrosylation. *Biochemistry* 2001; 40:1688-93.
80. Park SW, Huq MD, Hu X, Wei LN. Tyrosine nitration on p65: a novel mechanism to rapidly inactivate nuclear factor-kappaB. *Mol Cell Proteomics* 2005; 4:300-9.
81. Marshall H, Stamler J. Nitrosative stress-induced apoptosis through inhibition of NFkappaB. *J Biol Chem* 2002; 277:34223-8.
82. Reynaert NL, Ckless K, Korn SH, Vos N, Guala AS, Wouters EF, et al. Nitric oxide represses inhibitory kappaB kinase through S-nitrosylation. *Proc Natl Acad Sci USA* 2004; 101:8945-50.
83. Simpson CS, Morris BJ. Activation of nuclear factor kappaB by nitric oxide in rat striatal neurons: differential inhibition of the p50 and p65 subunits by dexamethasone. *J Neurochem* 1999; 73:353-61.
84. Miscusi M, Ebner F, Ceccariglia S, Menegazzi M, Mariotto S, Berra L, et al. Early nuclear factor-kappaB activation and inducible nitric oxide synthase expression in injured spinal cord neurons correlating with a diffuse reduction of constitutive nitric oxide synthase activity. *J Neurosurg Spine* 2006; 4:485-93.
85. Nikitovic D, Holmgren A, Spyrou G. Inhibition of AP-1 DNA binding by nitric oxide involving conserved cysteine residues in Jun and Fos. *Biochem Biophys Res Commun* 1998; 242:109-12.
86. Klatt P, Molina EP, Lamas S. Nitric oxide inhibits c-Jun DNA binding by specifically targeted S-glutathionylation. *J Biol Chem* 1999; 274:15857-64.
87. Wu J, Fang L, Lin Q, Willis WD. Fos expression is induced by increased nitric oxide release in rat spinal cord dorsal horn. *Neurosci* 2000; 96:351-7.
88. Najimi M, Robert JJ, Mallet J, Rostène W, Forgez P. Neurotensin induces tyrosine hydroxylase gene activation through nitric oxide and protein kinase C signaling pathways. *Mol Pharmacol* 2002; 62:647-53.
89. Chan SH, Chang KF, Ou CC, Chan JY. Nitric oxide regulates c-fos expression in nucleus tractus solitarius induced by baroreceptor activation via cGMP-dependent protein kinase and cAMP response element-binding protein phosphorylation. *Mol Pharmacol* 2004; 65:319-25.
90. Hemish J, Nakaya N, Mittal V, Enikolopov G. Nitric oxide activates diverse signaling pathways to regulate gene expression. *J Biol Chem* 2003; 278:42321-9.
91. Dhakshinamoorthy S, Sridharan S, Li L, Ng P, Boxer L, Porter A. Protein/DNA arrays identify nitric oxide-regulated cis-element and trans-factor activities some of which govern neuroblastoma cell viability. *Nucleic Acids Res* 2007; 35:5439-51.
92. Barnabé-Heider F, Miller FD. Endogenously produced neurotrophins regulate survival and differentiation of cortical progenitors via distinct signaling pathways. *J Neurosci* 2003; 23:5149-60.
93. Bartkowska K, Paquin A, Gauthier A, Kaplan D, Miller F. Trk signaling regulates neural precursor cell proliferation and differentiation during cortical development. *Development* 2007; 134:4369-80.
94. Colussi C, Mozzetta C, Gurtner A, Illi B, Rosati J, Straino S, et al. HDAC2 blockade by nitric oxide and histone deacetylase inhibitors reveals a common target in Duchenne muscular dystrophy treatment. *Proc Natl Acad Sci USA* 2008; 105:19183-7.
95. Hu E, Chen Z, Fredrickson T, Zhu Y, Kirkpatrick R, Zhang GF, et al. Cloning and characterization of a novel human class I histone deacetylase that functions as a transcription repressor. *J Biol Chem* 2000; 275:15254-64.
96. Lee H, Rezai-Zadeh N, Seto E. Negative regulation of histone deacetylase 8 activity by cyclic AMP-dependent protein kinase A. *Mol Cell Biol* 2004; 24:765-73.
97. Boo YC, Jo H. Flow-dependent regulation of endothelial nitric oxide synthase: role of protein kinases. *Am J Physiol, Cell Physiol* 2003; 285:499-508.
98. Braam B, de Roos R, Bluyssen H, Kemmeren P, Holstege F, Joles JA, et al. Nitric oxide-dependent and nitric oxide-independent transcriptional responses to high shear stress in endothelial cells. *Hypertension* 2005; 45:672-80.
99. Illi B, Dello Russo C, Colussi C, Rosati J, Pallaoro M, Spallotta F, et al. Nitric oxide modulates chromatin folding in human endothelial cells via protein phosphatase 2A activation and class II histone deacetylases nuclear shuttling. *Circ Res* 2008; 102:51-8.
100. McKinsey TA, Zhang CL, Olson EN. Identification of a signal-responsive nuclear export sequence in class II histone deacetylases. *Mol Cell Biol* 2001; 21:6312-21.
101. Sucharov CC, Langer S, Bristow M, Leinwand L. Shuttling of HDAC5 in H9C2 cells regulates YY1 function through CaMKIV/PKD and PP2A. *Am J Physiol Cell Physiol* 2006; 291:1029-37.
102. Watson P, Riccio A. Nitric oxide and histone deacetylases. A new relationship between old molecules. *Commun Integr Biol* 2009; 2:1-3.
103. Steffan JS, Bodai L, Pallos J, Poelman M, McCampbell A, Apostol BL, et al. Histone deacetylase inhibitors arrest polyglutamine-dependent neurodegeneration in *Drosophila*. *Nature* 2001; 413:739-43.
104. Hughes RE. Polyglutamine disease: acetyltransferases awry. *Curr Biol* 2002; 12:141-3.
105. Abel T, Zukin RS. Epigenetic targets of HDAC inhibition in neurodegenerative and psychiatric disorders. *Curr Opin Pharmacol* 2008; 8:57-64.
106. Perez MK, Paulson HL, Pendse SJ, Saionz SJ, Bonini NM, Pittman RN. Recruitment and the role of nuclear localization in polyglutamine-mediated aggregation. *J Cell Biol* 1998; 143:1457-70.
107. Kazantsev A, Preisinger E, Dranovsky A, Goldgaber D, Housman D. Insoluble detergent-resistant aggregates form between pathological and nonpathological lengths of polyglutamine in mammalian cells. *Proc Natl Acad Sci USA* 1999; 96:11404-9.
108. Butler R, Bates GP. Histone deacetylase inhibitors as therapeutics for polyglutamine disorders. *Nat Rev Neurosci* 2006; 7:784-96.
109. Zuccato C, Ciammola A, Rigamonti D, Leavitt BR, Goffredo D, Conti L, et al. Loss of huntingtin-mediated BDNF gene transcription in Huntington's disease. *Science* 2001; 293:493-8.
110. Zuccato C, Tartari M, Crotti A, Goffredo D, Valenza M, Conti L, et al. Huntingtin interacts with REST/NRSF to modulate the transcription of NRSE-controlled neuronal genes. *Nat Genet* 2003; 35:76-83.
111. Ginsburg E, Salomon D, Sreevalsan T, Freese E. Growth inhibition and morphological changes caused by lipophilic acids in mammalian cells. *Proc Natl Acad Sci USA* 1973; 70:2457-61.
112. Altenburg BC, Via DP, Steiner SH. Modification of the phenotype of murine sarcoma virus-transformed cells by sodium butyrate. Effects on morphology and cytoskeletal elements. *Exp Cell Res* 1976; 102:223-31.
113. Boffa LC, Vidali G, Mann RS, Allfrey VG. Suppression of histone deacetylation in vivo and in vitro by sodium butyrate. *J Biol Chem* 1978; 253:3364-6.
114. Huang Y, Man HY, Sekine-Aizawa Y, Han Y, Juluri K, Luo H, et al. S-nitrosylation of N-ethylmaleimide sensitive factor mediates surface expression of AMPA receptors. *Neuron* 2005; 46:533-40.
115. Song J, Noh JH, Lee JH, Eun JW, Ahn YM, Kim SY, et al. Increased expression of histone deacetylase 2 is found in human gastric cancer. *APMIS* 2005; 113:264-8.
116. Hrzenjak A, Moinfar F, Kremser ML, Strohmeier B, Staber PB, Zatloukal K, et al. Valproate inhibition of histone deacetylase 2 affects differentiation and decreases proliferation of endometrial stromal sarcoma cells. *Mol Cancer Ther* 2006; 5:2203-10.
117. Lagger G, O'Carroll D, Rembold M, Khier H, Tischler J, Weitzer G, et al. Essential function of histone deacetylase 1 in proliferation control and CDK inhibitor repression. *EMBO J* 2002; 21:2672-81.
118. Glazak MA, Seto E. Histone deacetylases and cancer. *Oncogene* 2007; 26:5420-32.
119. Kim SO, Merchant K, Nudelman R, Beyer WF, Keng T, DeAngelo J, et al. OxyR: a molecular code for redox-related signaling. *Cell* 2002; 109:383-96.